SUPPLEMENTAL LABELING

TOLFENPYRAD	GROUP	21A	INSECTICIDE
	GROUP	39	FUNGICIDE



Pome Fruit (Crop Group 11-10)

apple; azarole; crabapple; loquat; mayhaw; Hook. & Arn., and *C. rufula* Sarg; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

This supplemental labeling expires 06/18/2021 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for BEXAR insecticide before applying. Use of BEXAR insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for BEXAR insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

APPLICATION RATE CHART FOR BEXAR INSECTICIDE

Pest	Rate/Acre	Use Directions
Katydid Leafhoppers Aphids (excluding woolly apple aphid)	14.0 to 21.0 fl oz/acre (0.15 to 0.22 lb ai/acre) 17.0 to 21.0 fl oz/acre (0.18 to 0.22 lb ai/acre)	 USE RESTRICTIONS Apply by ground only, using a minimum of 100 gallons of water per acre. Do not apply by air on pome fruit. Do not apply more than 53.5 fl oz (0.56 lb ai) per acre per year. Do not make more than 2 applications per year.
Apple maggot Leafrollers Mealybugs Pear psylla Pear rust mite Plum curculio Thrips Codling moth*	21.0 to 27.0 fl oz/acre (0.22 to 0.28 lb ai/acre)	 Allow at least 14 days between applications. Preharvest Interval (PHI): 14 days
Powdery mildew* *suppression only		

©2018 Nichino America, Inc. BEXAR and Nichino America logo are registered trademarks of Nichino America, Inc.

NICHINO
AMERICA*
Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700