SUPPLEMENTAL LABELING

TOLFENPYRAD	GROUP	21A	INSECTICIDE
	GROUP	39	FUNGICIDE



Citrus Fruit (Crop Group 10-10)

Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin orange, clementine); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these

This supplemental labeling expires 06/18/2021 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for BEXAR insecticide before applying. Use of BEXAR insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for BEXAR insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

APPLICATION RATE CHART FOR BEXAR INSECTICIDE

Pest Rate/Acre		Use Directions	
Asian citrus psyllid	14.0 to 27.0 fl oz/acre	• For high air velocity, low volume or air	
Katydid	(0.15 to 0.28 lb ai/acre)	curtain sprayers, apply using a minimum of 30 gallons of water per acre	
Aphids	17.0 to 27.0 fl oz/acre	USE RESTRICTIONS	
	(0.18 to 0.28 lb ai/acre)	Do not apply by air on citrus fruit.	
Citrus bud mite		 Apply by ground only, using a minimum of 100 gallons of water per acre. 	
Citrus mealybug	21.0 to 27.0fl oz/acre	 Do not use on nursery stock. 	
Citrus thrips	(0.22 to 0.28 lb ai/acre)	 Do not apply more than 27.0 fluid ounces 	
Soft scale insects, including		(0.28 lb ai) per acre per growing season.	
citricola scale, barnacle		, , , , , , , , , , , , , , , , , , , ,	
scale, brown soft scale		Do not make more than 1 application per growing coacon.	
Citrus peelminer		growing season.	
Citrus red mite	24.0 to 27.0 fl oz/acre	Do not make more than 2 applications per	
Citrus rust mite (silver mite) Mealybugs	(0.25 to 0.28 lb ai/acre)	Preharvest Interval (PHI): 3 days	
Citrus orangedog*			
Citrus weevil*			
Cutworms*			
*suppression only			

©2018 Nichino America, Inc. BEXAR and Nichino America logo is registered trademark of Nichino America, Inc.

NICHINO
AMERICA*
Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700