
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1480A-S1 1220

AZOXYSTROBIN	GROUP	11	FUNGICIDE
BENZOVINDIFLUPYR	GROUP	7	FUNGICIDE

Elatus® Fungicide

SOLATENOL® Technology*

This supplemental label expires on 12/30/2023 and must not be used or distributed after this date.

Active Ingredients:	
Azoxystrobin**	30.0%
Benzovindiflupyr***	15.0%
<hr/>	
Other Ingredients:	55.0%
Total:	100.0%

*Technology denotes the active ingredient, Benzovindiflupyr.

**CAS No. 131860-33-8

***CAS No. 1072957-71-1

Elatus Fungicide is formulated as a wettable granule (WG) and contains 0.30 lb ai azoxystrobin and 0.15 lb ai benzovindiflupyr per pound.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1480

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Elatus Fungicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



DIRECTIONS FOR USE

Ginseng

Crops (Including all cultivars, varieties, and/or hybrids of these) (Not for use in California)			
Ginseng			
Target Disease	Rate (oz/A)	Application Timing	Use Directions
Alternaria blight (<i>Alternaria panax</i>)	7.3*	For foliar diseases, make an application at the onset of disease or when conditions are conducive for disease.	Apply by ground or chemigation. Apply in a minimum spray volume of 50 gallons per acre. See Section 4.4.5
*7.3 oz product/A is equivalent to 0.137 lb ai azoxystrobin and 0.068 lb ai benzovindiflupyr.			
Resistance Management: • Refer to Section 3.1 .			
USE RESTRICTIONS			
1. Refer to Section 6.1 for additional product use restrictions. 2. Maximum Single Application Rate: 7.3 oz/A 3. Minimum Application Interval: 14 days 4. Maximum Annual Rate: 29.2 oz/A/year a. DO NOT exceed 0.272 lb ai/A/year of benzovindiflupyr-containing products. b. DO NOT exceed 2.0 lb ai/A/year of azoxystrobin-containing products. 5. DO NOT exceed 4 applications per year. 6. DO NOT apply by air. 7. Pre-Harvest Interval (PHI): 15 days			

Sugar Beet

Crops (Including all cultivars, varieties, and/or hybrids of these) (Not for use in California)			
Sugar beet			
Target Disease	Rate (oz/A)	Application Timing	Use Directions
Soilborne Diseases: Circular Spot, Southern Blight <i>(Sclerotium rolfsii)</i> Pythium Root Rot <i>(Pythium aphanidermatum)</i> Rhizoctonia Stem Canker, Crown Rot <i>(Rhizoctonia solani)</i>	0.3 – 0.6 oz/1000 linear row feet	Apply in-furrow at planting.	Apply by ground. Do not apply as a dribble application over the seed row. If cool soil conditions are expected after planting which could result in an extended period of plant emergence, do not apply Elatus Fungicide in-furrow. If using Elatus Fungicide at the time of planting, do not use a starter fertilizer with it.
Circular Spot, Southern Blight <i>(Sclerotium rolfsii)</i> Pythium Root Rot <i>(Pythium aphanidermatum)</i> Rhizoctonia Stem Canker, Crown Rot <i>(Rhizoctonia solani)</i>	7.1	Apply at the 2- to 8-leaf stage.	Apply by ground using 3-7 inch banded applications in a minimum of 10 gallons per acre.
Precaution: • Tank mixtures of Elatus Fungicide with crop oil concentrates (COC) or methylated spray oil (MSO) may result in crop injury.			
Resistance Management: • Refer to Section 3.1 .			
USE RESTRICTIONS			
1. Refer to Section 6.1 for additional product use restrictions. 2. Maximum Single Application Rate: 7.1 oz/A 3. Minimum Application Interval: 5 days 4. Maximum Annual Rate: 14.6 oz/A/year a. DO NOT exceed 0.136 lb ai/A/year of benzovindiflupyr-containing products. b. DO NOT exceed 2.0 lb ai/A/year of azoxystrobin-containing products. 5. DO NOT exceed 2 applications per year. 6. Apply as an in-furrow spray in a minimum of 10 gallons per acre. 7. Pre-Harvest Interval (PHI): Apply up to the BBCH 31 growth stage.			

©2021 Syngenta

Elatus® and the Syngenta logo are trademarks of a Syngenta Group Company