

PENOX SULAM	GROUP	2	HERBICIDE
OXYFLUORFEN	GROUP	14	HERBICIDE

# FIFRA Section 24(c) Special Local Need (SLN) Label

**Corteva Agriscience LLC      9330 Zionsville Road      Indianapolis, IN 46268-1054 USA**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON**

## **Cleantraxx™**

EPA Reg. No. 62719-702

EPA SLN No. OR-220014

**This label is valid until December 31, 2027, or until otherwise amended, withdrawn, canceled, or suspended**

### **For Control of Herbaceous Broadleaf Weeds and Grasses in Christmas Tree Plantations**

**Active Ingredient:**

penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazolo[1,5c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide .....	0.85%
oxyfluorfen: 2-chloro-1-(3-ethoxyl-4-nitrophenoxy)-4-(trifluoromethyl) benzene .....	40.31%
Other Ingredients .....	58.84%
Total.....	100.00%

Contains 0.083 lb penoxsulam and 3.93 lb oxyfluorfen active ingredient per gallon

**Keep Out of Reach of Children**

## **CAUTION**

### **Environmental Hazards**

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. See Directions for Use for additional restrictions. Do not contaminate water when disposing of equipment wash water or rinsate.

**GROUNDWATER ADVISORY STATEMENT:** This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**NON-TARGET ORGANISM ADVISORY STATEMENT:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Read this SLN label and the label affixed to the container of Cleantraxx™ before applying. Carefully follow all precautionary statements and applicable use directions.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on this SLN label and the label affixed to the container of Cleantraxx.

## Product Information

Cleantraxx is a selective herbicide for preemergence and postemergence residual weed control of certain broadleaf and grass weeds in Christmas tree plantations. Apply Cleantraxx in fall through early spring, prior to germination of targeted weeds, as per labeled use directions. The best weed control is obtained by application to weeds either preemergence or early postemergence when weeds are small and actively growing. Any cultural practices that disturb or redistribute surface soil following treatment with Cleantraxx will reduce weed control effectiveness.

## Application Directions

- Make a single application of Cleantraxx at 3 to 4.5 pints per acre in fall or spring. Cleantraxx may be applied broadcast by aerial or ground application or as a spot or directed spray application over-the-top of dormant Christmas trees for preemergent and early postemergent control of weeds. When making an application to newly planted Christmas trees, ensure soil has settled before applying. Apply Cleantraxx only when trees are dormant and there is no new growth, as applications made during active growth stages may injure desirable trees. In general, applications made before bud break or after terminals harden off have the least potential for injury to Christmas trees. Users should determine through field trials whether applications of Cleantraxx give acceptable tree tolerance under local conditions. Apply the specified rate of Cleantraxx in a total spray volume of at least 10 gallons per acre.
- Preemergence control: Cleantraxx is stable on the soil surface for up to 21 days, but must be incorporated by moisture to provide effective preemergence control of susceptible weeds. A single rainfall of 0.5 inches or more, within 21 days after application, is necessary to activate Cleantraxx.
- Postemergence control: The addition of an adjuvant at 1 quart per 100 gallons of spray volume (0.25% v/v) of a crop oil concentrate, methylated seed oil or 80% active (minimum) nonionic surfactant is required for effective postemergence control of susceptible emerged weeds. For applications over the top of Christmas trees, use only adjuvants/surfactants for which the desired Christmas trees are known to be tolerant.

## Application Precautions

- For best control of emerged weeds, apply a tank mix of Cleantraxx with a postemergence herbicide registered for use on Christmas trees.
- Where rate ranges are given, use a lower rate in the rate range on coarse textured soils low in organic matter, lighter weed infestations and for reduced lengths of residual weed control. Use a higher rate in the rate range on medium to fine textured soils, soils containing higher organic matter, heavy weed infestations, or for extended residual preemergence weed control.
- Preemergence weed control is most effective when soil surfaces are clean (free of weed residues or clippings) and weed free.
- Postemergence applications of Cleantraxx are most effective when made to weeds at the seedling stage. Applications made later than the 4-inch or 4 leaf stage of susceptible weeds may result in partial

control or suppression. Make postemergence applications to seedling grasses not exceeding the 2-leaf stage.

- Any cultural practices, cultivation, or disturbance of the soil surface after application will decrease the weed control provided by Cleantraxx.

### **Use Restrictions**

- Do not apply more than 4.5 pints of Cleantraxx per acre per application.
- Do not make more than one application per year.
- Do not apply Cleantraxx until soil has been settled by packing or rainfall.
- Do not apply Cleantraxx when weather conditions favor drift. Avoid drift to all non-target crops and areas.
- Do not treat inside banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not contaminate water used for irrigation or domestic purposes.
- Do not feed or allow animals to graze on any areas treated with Cleantraxx.
- Do not apply to frozen soil or snow-covered soil.
- **BUFFER REQUIREMENTS:** A 25 foot vegetative buffer strip must be maintained between all areas treated with this product and lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- **Chemigation:** Do not apply this product through any type of irrigation system.

### **Conditions and Risks of Use for Special Local Need**

**USE OF CLEANTRAXX (THE “PRODUCT”) ON CHRISTMAS TREE PLANTATIONS (THE “CROP”) FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP.**

**This Product is available for use in the manner described in this Special Local Need (SLN) label on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this SLN Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.**

**By purchasing the Product for use, or using the Product, in the manner described in this SLN Labeling, you acknowledge and accept that to the extent permitted by law:**

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;**
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this SLN labeling;**
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;**
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;**

**(5) these Conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Corteva Agriscience, or its respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.**

**If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.**

**These Conditions and Risks of Use for Special Local Need are required by Corteva Agriscience and not specified by the US EPA or the State of Oregon.**

**24(c) Registrant:**

**Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

<sup>TM</sup>® Trademarks of Corteva Agriscience and its affiliated companies

R419-010  
Accepted: 12/07/22  
Initial Printing