

FIFRA Section 24(c) Special Local Need (SLN) Label

For Distribution and Use Only Within the State of Oregon

Fontelis[®]
EPA Reg. No. 352-834
EPA SLN No. OR-220004

This label is valid until December 31, 2024 or until otherwise amended, withdrawn, canceled, or suspended.

For Control of Foliar and Soil-borne Plant Diseases in Potatoes

<i>Active Ingredient</i>	<i>By Weight</i>
Penthiopyrad	20.4%
<i>Other Ingredients</i>	79.6%
TOTAL	100.0%

Contains 1.67 pounds of penthiopyrad per gallon of product

Keep Out of Reach of Children **CAUTION**

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and oysters. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this chemical from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Read this SLN label and the label affixed to the container for Fontelis fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the label affixed to the container for Fontelis.

Fontelis is a broad-spectrum fungicide, recommended for control of foliar and soil-borne plant diseases in many crops, including potatoes, and has preventive, curative, and locally systemic activity. Fontelis must be applied in a regularly scheduled protective spray program in rotation with other fungicides

Disease Controlled or Suppressed	Product Rate fl oz/acre	Comments
Alternaria early blight and brown spot (Alternaria solani, Alternaria alternata) Gray mold (Botrytis cinerea) Powdery mildew (Erysiphe spp.)	10 to 24 fl oz	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
Black dot (Colletotrichum coccodes)	14 to 24 fl oz	
Disease suppression White mold (Sclerotinia sclerotiorum)	14 to 24 fl oz	Make initial application at 100% full bloom of the primary inflorescence, or prior to row closure, and then again 14 days later.
Soil-borne diseases Rhizoctonia stem canker and black scurf (Rhizoctonia solani)	0.7 to 1.6 fl oz/1000 row-ft	At-plant, in-furrow application. Maximum rate per acre per application is 24 fl oz. For instructions, see "Soilborne/Seedling Disease Control for Tomatoes, Snap Beans and Bulb Onions" on the label affixed to the container for Fontelis and the "Soil Application Rates for Potatoes" below.
<p>Make no more than 2 sequential applications of Fontelis before switching to a fungicide with a different mode of action. Minimum time (PHI) between application and harvest is 7 days. Do not exceed 72 fl oz (0.94 lb active ingredient) of Fontelis/acre per year from any combination of seed, soil, or foliar treatments.</p> <p>Do not use Fontelis fungicide for an in-furrow application when penthiopyrad has been applied as a seed piece treatment for potatoes. Do not apply more than two in-season foliar potato applications of Fontelis at the maximum single application rate when penthiopyrad has been used as a seed piece treatment for potatoes. Do not apply more than 15 ounces of penthiopyrad active ingredient/acre/year to potatoes in total from any combination of seed, soil, or foliar treatments.</p>		

Soil Application Rates for Potatoes

Rate per 1000 row feet	Product per Acre (fl oz) ^a						
	22 " rows	30 " rows	32 " rows	34 " rows	36 " rows	38 " rows	40 " rows
0.7	16.7	12.2	11.4	10.9	10.2	9.6	9.2
1.2	28.6 ^b	20.9	19.6	18.5	17.4	16.5	15.7
1.6	-	27.9 ^c	26.1 ^d	24.6 ^e	23.0	22.0	21.1

^a Consult the maximum rate per acre allowed for the crop, and do not exceed that rate when using this application method.

^b In 22 inch rows, the highest rate for crops with 24 fl oz/acre maximums is 1.0 fl oz/1000 ft row, and for crops with 30 fl oz/acre maximums is 1.26 fl oz/1000 ft row.

^c In 30 inch rows, the highest rate for crops with 24 fl oz/acre maximums is 1.38 fl oz/1000 ft row.

^d In 32 inch rows, the highest rate for crops with 24 fl oz/acre maximums is 1.47 fl oz/1000 ft row.

^e In 34 inch rows, the highest rate for crops with 24 fl oz/acre maximums is 1.56 fl oz/1000 ft row.

Use Restrictions

- - Use this product only in commercial and farm plantings.
- - Do not use for home plantings.
- - Do not formulate this product into other end-use products.

24(c) Registrant:

E. I. du Pont de Nemours & Company
Chestnut Run Plaza
974 Centre Road
Wilmington, DE 19805

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R613-012

Accepted Date: 02/27/22

Initial Printing