

FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Stinger®

EPA Reg. No. 62719-73

EPA SLN No. NY-220001

For Distribution and Use Only Within the State of New York

This label is valid until December 31, 2026, or until otherwise amended, withdrawn, canceled, or suspended.

Perennial Sowthistle Control in Dry Bulb Onion

Use of Stinger® herbicide on dry bulb onion may result in crop injury, crop yield reduction and/or crop loss. Read and understand the Special Conditions and Risks of Use section on this label before using this product in accordance with this FIFRA Section 24(c) Special Local Need.

Active Ingredient:	% by Weight
clopyralid: 3,6-dichloro-2-pyridinecarboxylic acid, monoethanolamine salt.....	40.9%
Other Ingredients.....	59.1%
Total.....	100.0%

Acid Equivalent: clopyralid: 3,6-dichloro-2-pyridinecarboxylic acid - 31% (3 lb/gal)

Keep Out of Reach of Children

CAUTION

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Read this SLN label and the label affixed to the container for Stinger herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this SLN label, use of Stinger according to this SLN is subject to all use directions, precautions and limitations imposed by the label affixed to the container for Stinger.

Application Rate: Apply 1/4 pint of Stinger per acre for the first application (0.094 pound acid equivalent per acre) as a foliar broadcast application when dry bulb onions are at the 4-leaf stage to control perennial sowthistle. Apply 1/2 pint of Stinger per acre for the second foliar broadcast application (0.188 pound acid equivalent per acre). A nonionic surfactant may be added at a rate of 0.25% v/v. Apply Stinger in a minimum spray volume of 10 gallons per acre.

Restrictions:

- Do not apply more than a total of 3/4 pint (0.282 pounds acid equivalent per acre) of Stinger per acre per year per crop.
- The minimum retreatment interval is 14 days.
- Do not make more than two applications per crop per year (0.094 pound acid equivalent per acre rate for the first application and 0.188 pound acid equivalent per acre rate for the second application).

Precautions:

- Stinger is a residual herbicide and applications should be made based upon accurate rate per acre calibrations. Applications of Stinger can injure the dry bulb onion plant and significantly reduce yields depending upon rates used, timing of application, and environmental conditions.

Special Conditions and Risks of Use

USE OF Stinger® herbicide (THE “PRODUCT”) ON DRY BULB ONION (THE “CROP”) MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Corteva Agriscience makes the Product available for use in the manner described in this Special Local Need (SLN) on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this SLN Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this SLN Labeling, you acknowledge and accept that:

- (1) You assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this SLN Labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agriscience and not specified by the U.S. EPA or the State of New York.

24(c) Registrant:
Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054 USA

™® Trademarks of Corteva Agriscience and its affiliated companies

R043-124

Accepted Date: 03/01/22

Initial Printing