



A systemic fungicide for the control of apple, almond, asparagus*, barley*, bean* (except succulent), corn*, grape, grasses grown for seed, hop*, leafy brassica green*, onion, peanut*, pecan, pistachio, soybean*, stone fruit, sunflower, walnut, watermelon, and wheat diseases

ACTIVE INGREDIENTS:

Potassium phosphite¹ 49.0%
Tebuconazole: α -[2-(4-chlorophenyl)ethyl]- α -(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol² 3.3%

OTHER INGREDIENTS: 47.7%

TOTAL: 100.0%

¹ 30.4% (3.66 lbs/gal) phosphorous acid equivalent; ² 0.42 lbs/gal tebuconazole

*Not for use in California

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical information.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. For additional information, call: 1-800-424-9300.	
SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND FOR DIRECTIONS FOR USE	

SHAKE WELL BEFORE USE

FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE CALL TOLL FREE: 1-800-424-9300.

AGRICULTURAL CHEMICAL: DO NOT SHIP OR STORE WITH FOODS, FEEDS, DRUGS OR CLOTHING.

EPA Reg. No. 42519-35-5905 EPA Est. No. 5905-GA-1

NET CONTENTS: 2.5 Gallons

Manufactured For

HELENA AGRI-ENTERPRISES, LLC

225 SCHILLING BOULEVARD, SUITE 300

COLLIERVILLE, TENNESSEE 38017 U.S.A. • (901) 761-0050

5043-9/19025 1/19
AD 020618HAE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Natural rubber and polyvinyl chloride are chemical-resistant to this product.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- chemical-resistant gloves (nitrile, butyl, neoprene, and/or barrier laminate)
- shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Tebuconazole is toxic to mammals, fish, and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas.

GROUND WATER ADVISORY

Tebuconazole is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

SURFACE WATER ADVISORY

Tebuconazole may contaminate water through drift of spray in wind. It has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or tribal agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

5043-9/190625_2/19
AD 020618HAE

Failure to follow the **USE DIRECTIONS, RESTRICTIONS** and **PRECAUTIONS** on this label may result in crop injury or poor disease control and/or illegal residues.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all crops, except apples and stone fruit. The REI for apples and stone fruit is 5 days.

PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- coveralls
- chemical-resistant gloves (nitrile, butyl, neoprene, natural rubber, and/or barrier laminate)
- shoes plus socks

CHEMIGATION

Do not apply this product through any type of irrigation system.

REQUIREMENTS FOR GROUND APPLICATIONS

For ground applications, adjust nozzle height and droplet size with wind speed according to the following table:

Wind speed	Nozzle height	Droplet size for standard nozzles (ASAE standard 572)
Less than 10 mph	Up to 2 feet	medium or coarser
	2-4 feet	coarse or coarser
	4-6 feet	very coarse or coarser
10 to 15 mph	0-2 feet	coarse or coarser
	2-4 feet	very coarse or coarser
	4-6 feet	extremely coarse

RESTRICTIONS: Do not apply when the wind speed exceeds 15 mph. Do not apply at a nozzle height of greater than 6 feet above the ground or crop canopy. Apply as a medium or coarser spray (ASAE standard 572).

REQUIREMENTS FOR AERIAL APPLICATIONS

For aerial applications, apply only when the wind speed is less than or equal to 15 mph using a release height of no more than 10 feet above the ground or crop canopy. If the wind speed is less than 10 mph, apply as a medium or coarser spray (ASAE standard 572). If the wind speed is between 10 mph and 15 mph, apply as a coarse or coarser spray (ASAE standard 572). The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. When aerial applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. No aerial application in New York State.

RESTRICTIONS: Do not make aerial applications into temperature inversions.

5043-9/190625 3/19
AD 020618HAE

For the following row crops (barley, corn, dry beans, soybeans, and wheat), make applications by air in a minimum of 2 gallons of water per acre. For fruit and vegetable crops specified on this label, as well as grasses grown for seed, use a minimum of 5 gallons of water per acre for aerial applications. For almond, pistachio, and walnut, use a minimum of 10 gallons of water per acre for aerial applications. Approved adjuvants and other additives may be added to the spray solution to improve spray coverage. **VIATHON®** can be tank-mixed with most recommended fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants and other additives. Under some conditions, the use of additives or adjuvants may improve the performance of **VIATHON®**. However, all varieties and cultivars have not been tested with all possible tank-mix combinations. Physical incompatibility, reduced disease control, or crop injury may result from mixing **VIATHON®** with other products. Therefore, before using any tank-mix, test the combination on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of the tank-mix application.

SPRAY DRIFT MANAGEMENT

Aerial applications: Mount the spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use the minimum practical boom length, which must not exceed 75 percent of the wingspan or rotor diameter. Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 2 gallons of spray solution per acre by aircraft spray equipment. Release spray at the lowest possible height consistent with good pest control and flight safety. **RESTRICTION:** Do not apply more than 10 feet above the crop canopy. Make aerial and ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). **RESTRICTION:** Do not apply when wind velocity exceeds 15 mph (miles per hour). Avoid applications when wind gusts approach 15 mph. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area. Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. **RESTRICTION:** Do not spray during conditions of low humidity and/or high temperature. **RESTRICTION:** Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Rotational Crop Restrictions:

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any other crop may be planted into treated areas 120 days after last application.

PRODUCT INFORMATION

VIATHON® is a systemic product. When used in conjunction with good cultural management practices and as part of a complete disease control program, **VIATHON®** will control/suppress specified diseases associated with apples, almond, asparagus, barley, beans, corn, grapes, grasses grown for seed, hops, leafy brassica greens, onions, peanuts, pecans, pistachio, soybeans, stone fruit, sunflowers, walnut, watermelons, and wheat. In order to achieve maximum results in controlling/suppressing the diseases, carry out spraying before the appearance of the diseases or when diseases are first observed. Initiate applications when environmental conditions are favorable for disease development. **Not approved for use on corn, apples and soybeans in New York State.**

Resistance Management: As **VIATHON®** contains a demethylation inhibitor (DMI), it is best applied in alternation with a non-DMI fungicide as a resistance management strategy.

5043-9/190625_4/19
AD 020618HAE

MIXING AND APPLICATION INSTRUCTIONS

For barley, corn, dry beans, soybeans, and wheat: Apply **VIATHON**[®] in a minimum of 10 gallons of water per acre by ground sprayer or in a minimum of 2 gallons of spray solution per acre by aircraft.

For fruit and vegetable crops, as well as grasses grown for seed: **VIATHON**[®] is best applied in 50-100 gallons of water per acre by ground sprayer or in a minimum of 5 gallons or more of water per acre by aircraft.

For almonds, pistachios, and walnuts: Apply **VIATHON**[®] in a minimum of 50 gallons per acre by ground sprayer or in a minimum of 10 gallons per acre by aircraft. Ground applications are best applied after petal fall.

Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the volume of water in the spray tank for complete crop coverage.

RESTRICTION: Do not exceed specified application rates. Fill spray tank halfway with clean water. Add the required amount of **VIATHON**[®] to the tank, as indicated in the application instructions. While agitating, add the remaining amount of water. Apply solution directly to the foliage. For optimum control, thorough coverage is required. **VIATHON**[®] spray must have a minimum of 2 hours of drying time on plant foliage for the active ingredients to move into the plant tissue before rain or irrigation occurs.

Compatibility:

Conduct a tank spray compatibility test when you plan to mix this product with other products. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, and then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank. Follow the more restrictive labeling requirements of any tank mix partner.

RESTRICTION: Do not tank mix with products whose label prohibits tank mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Phytotoxicity indications:

The components of this product have been evaluated for phytotoxicity on a large variety of crops under various normal field conditions. However, testing all plant varieties, in all mixtures and combinations is not feasible. Test a small portion of the area to be treated for phytotoxicity, before treating the entire area. Test tank-mix combinations on a small number of plants before treating large areas, as crop sensitivity to these mixtures may vary. Do not apply to plants under stress.

DISEASE CONTROL AND APPLICATION INSTRUCTIONS

VIATHON® is for use on the following crops, all of which have an REI of 12 hours, except apples and stone fruit. Apples and stone fruit have an REI of 5 days.

CROP	DISEASE	RATE	INSTRUCTIONS
ALMONDS, WALNUTS	Brown rot blossom blight (<i>Monilinia</i> spp.) Anthracnose (<i>Colletotrichum gleosporoides</i>) Powdery mildew* (<i>Podosphaera tridactyla</i>) Scab (<i>Cladosporium carpophilus</i>) Leaf rust* (<i>Tranzschelia discolor</i>)	2-4 pints/acre	<p>Begin applications before infection. Apply on a 7-14 day spray interval. Use the higher specified rate and shorter interval when disease pressure is severe. Refer to guidelines below for the best application timing for each disease.</p> <p>Brown rot blossom blight: Treat at pink bud (5-10% bloom) and full bloom. One application at full bloom is sufficient in most orchards in most years if there is no rain. If brown rot has been severe or in years of extended bloom accompanied by rainfall, a second or third application near full bloom may be necessary.</p> <p>Anthracnose: Apply VIATHON® sprays beginning at 5-10% bloom or pink bud and repeat every 10-14 days if rains persist. Late spring rains may necessitate additional applications.</p> <p>Powdery mildew: Make applications at jacket-split (late petal fall) and mid-spring.</p> <p>Scab: One application up to 5 weeks after petal fall is effective, but an earlier application (2 weeks after petal fall) is best.</p> <p>Leaf rust: Apply 5 weeks after petal fall and follow 4 to 5 weeks later in late spring and summer. Apply before rust symptoms are visible.</p>
	Alternaria	2.5-4 pints/acre or 2 pints VIATHON® + 4 pints chlorothalonil/acre	Alternaria: Make applications in May-June-July. Use the higher specified rate with increased disease pressure. For VIATHON® + chlorothalonil applications, make the first application in May, followed by VIATHON® applications alone in June and July.
	Botryosphaeria (walnuts only)	4 pints/acre	Botryosphaeria: Most effective when applied in May, June, July, and August.
<p>Specific Use Restrictions:</p> <p>1) Do not make more than 4 applications per season.</p> <p>2) Do not apply more than 16.5 pints per acre per season.</p> <p>3) Do not cut the cover crop in treated areas, or allow grazing by livestock in treated areas.</p> <p>4) The PHI is 35 days.</p> <p>* Not for use in California for powdery mildew and leaf rust.</p>			

5043-9/190625 6/19
AD 020618HAE

CROP	DISEASE	RATE	INSTRUCTIONS
APPLES Not approved for use on apples in New York State	Scab	2-3 pints/acre	Apply on a 7-10 day interval from greentip through 2 nd cover spray. Apply post-infection sprays within 72 hours after the beginning of a scab infection period.
	Cedar apple rust		Apply at 7-10 day intervals from pink bud through 2 nd cover spray.
	Powdery mildew		Apply at 7-10 day intervals from tight clusters through 2 nd cover spray. Tank mix the lowest specified rate of a spray surfactant to improve coverage. Alternate sprays with a non-DMI fungicide to address fungicide resistance management.
Specific Use Restrictions: 1) Do not make more than 6 applications per season. 2) Do not apply more than 12 pints per acre per season. 3) The PHI is 75 days.			
CROP	DISEASE	RATE	INSTRUCTIONS
ASPARAGUS*	Rust (<i>Puccinia</i> spp.)	2-3 pints/acre	Apply as a spray to the developing ferns after harvest. Apply at the earliest sign of disease or when weather is conducive to disease. Apply VIATHON ® in alternation with another effective fungicide. Repeat on a 14 day interval.
Specific Use Restrictions: 1) Do not make more than 4 applications per season. 2) Do not apply more than 9 pints per acre per season. 3) The PHI is 180 days. * Not for use in California			

CROP	DISEASE	RATE	INSTRUCTIONS
BARLEY*	Rusts Head blight suppression	2 pints/acre	Make a single application per season by air or ground at the earliest sign of rust (<i>Puccinia</i> spp.). Optimal timing for Fusarium head blight suppression is when heads are fully emerged on 50% of the plants (growth stage 10.5).
Specific Use Restrictions: 1) Straw cut after harvest may be fed or used for bedding. 2) Grazing or feeding forage to livestock is permitted 6 days after the last application. 3) Do not make more than 1 application per season. 4) Do not apply more than 2 pints per acre per season. 5) Do not apply within 30 days of harvest. * Not for use in California			
CROP	DISEASE	RATE	INSTRUCTIONS
BEANS* (fresh & dry, except succulent shelled)	Rust	2-3 pints/acre	Apply on a protective spray schedule or when weather is conducive for rust. Repeat applications on a 10-14 day interval, or as necessary to maintain control. VIATHON® may be applied with a low specified rate of surfactant.
Specific Use Restrictions: 1) On fresh beans, do not make more than 4 applications per season. Do not apply more than 12 pints per acre per season. The PHI is 7 days. 2) On dry beans, do not make more than 2 applications per season. Do not apply more than 6 pints per acre per season. The PHI is 14 days. * Not for use in California			

CROP	DISEASE	RATE	INSTRUCTIONS
CORN* Not approved for use on corn in New York State Field corn, Sweet corn, Seed corn, Popcorn	Rust (<i>Puccinia</i> spp.) Northern leaf blight (<i>Exserohilum turcicum</i>) Southern leaf blight (<i>Bipolaris maydis</i>) Northern leaf spot (<i>Bipolaris zeicola</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>)	2-3 pints/acre	Apply the higher specified rate (3 pts/acre) when conditions favor severe disease development or VIATHON® may be mixed with a strobilurin fungicide to improve control when conditions favor severe disease. Apply preventatively when weather favors disease development. Repeat on a 7-14 day interval. If a strobilurin mixing partner is applied, determine the physical compatibility of this product with VIATHON® . Use a jar test as described in the Compatibility section of this label. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.
Specific Use Restrictions: 1) Do not make more than 6 applications per season. 2) Do not apply more than 12 pints VIATHON® per acre per season. 3) The PHI is 7 days for applications of this product alone. 4) When tank mixing, no label dosage rate may be exceeded, and the most restrictive label precautions and restrictions must be followed. * Not for use in California, New York and Washington.			
CROP	DISEASE	RATE	INSTRUCTIONS
GRASSES GROWN FOR SEED	Rust (<i>Puccinia</i> spp.) Powdery mildew	2-4 pints/acre	Apply when weather conditions favor disease development. Repeat on a 14-day interval. Apply the higher rate under heavier disease pressure. Thorough coverage is important for disease control.
Specific Use Restrictions: 1) Do not make more than 4 applications per season. 2) Do not apply more than 8 pints per acre per season. 3) The PHI is 4 days. 4) Chaff, screenings and straw from treated areas may be used for feed purposes; do not forage, cut green crop, or use seed for feed purposes. 5) Regrowth may be grazed starting at 17 days after the last application.			

CROP	DISEASE	RATE	INSTRUCTIONS
GRAPES	Black rot Powdery mildew	2 pints/acre	Black rot: Make the first application when the new shoots are 1-3 inches. Continue on a 7-10 day interval until veraison is complete. Powdery mildew: Apply preventatively, starting before bloom and continue on a 14-21 day interval depending on weather and disease pressure.
	Downy mildew	2.5-4 pints/acre	Downy mildew: Apply preventatively using 2.5 pints in 50 gallons water/acre. Increase the rate and volume of spray with increasing disease pressure to obtain the necessary coverage and control.
	Ripe rot	4 pints/acre	Ripe rot: Apply at bunch closure and again at the veraison stage. As VIATHON® contains a DMI fungicide, it is best alternated in a spray schedule to address fungicide resistance management.

Specific Use Restrictions:
1) Do not make more than 4 applications per season.
2) Do not apply more than 16 pints per acre per season.
3) The PHI is 14 days.
4) Do not exceed spray solution concentration of 0.5% potassium phosphite.
5) Do not apply to Venifera grapes by back pack sprayer.

CROP	DISEASE	RATE	INSTRUCTIONS
HOPS*	Downy mildew Powdery mildew	2-4 pints/acre	Apply on a preventative spray schedule when the shoots are 6-12 inches. Repeat on a 10-14 day interval. Best applied with a low rate of surfactant.

Specific Use Restrictions:
1) Do not make more than 8 applications per season.
2) Do not apply more than 16 pints per acre per season.
3) The PHI is 14 days.
* Not for use in California

CROP	DISEASE	RATE	INSTRUCTIONS
LEAFY BRASSICA GREENS* Broccoli raab, Chinese cabbage (bok choy), Collards, Kale, Mizuna, Mustard greens, Mustard spinach, Rape greens, Turnip greens	Cercospora Powdery mildew Alternaria leaf spot Downy mildew	2 pints/acre	Apply on a 10-14 day schedule as a preventative treatment.
Specific Use Restrictions: 1) Do not make more than 4 applications per season. 2) Do not apply more than 8 pints per acre per season. 3) The PHI is 7 days. 4) Application to turnip greens is limited to East of the Rockies. * Not for use in California.			
CROP	DISEASE	RATE	INSTRUCTIONS
ONIONS Green onions Dry bulb onions Green onions include: leek, spring onion, scallion, Japanese bunching onions, green shallots, green eschalots Dry bulb onions include: garlic, elephant garlic, Welsh onion, shallot	Rust* (<i>Puccinia</i> spp.) Purple blotch (<i>Alternaria porii</i>) Downy mildew (<i>Peronospora destructor</i>)	2-3 pints/acre	Use VIATHON® as a preventative treatment. Repeat applications on a 10-14 day interval.
Specific Use Restrictions: 1) Do not make more than 3 applications per season. 2) Do not apply more than 6 pints per acre per season. 3) The PHI is 7 days. * Not for use in California for rust.			

CROP	DISEASE	RATE	INSTRUCTIONS
PEANUTS*	Soil-borne: White mold, southern blight (<i>Sclerotium rolfsii</i>) Rhizoctonia limb rot Pythium pod rot Foliar: Early and Late Leaf spot Leaf rust Web blotch	3-3.5 pints/acre	Apply in a seven application program in sprays 3, 4, 5, and 6 on a preventative 14-day schedule. Make applications of chlorothalonil prior to and following VIATHON® (sprays 1, 2, and 7) to control foliar diseases. Irrigate to carry the spray into the root and pod zone for soil-borne diseases. Use VIATHON® in conjunction with IPM practices to reduce severity of soil-borne diseases.
Specific Use Restrictions: 1) Do not make more than 4 applications per year. 2) Do not apply more than 14 pints per acre per year. 3) The PHI is 14 days. * Not for use in California.			
CROP	DISEASE	RATE	INSTRUCTIONS
PECANS	Anthracnose* Brown leaf spot* Downy spot* Liver spot* Scab Vein spot* Zonate leaf spot* Scab Vein spot* Zonate leaf spot*	Early Season: 2 pints/acre Post pollination: 2-2.5 pints/ acre	Apply in 100 gallons of water by ground or 5 gallons or more diluent by air in a preventative spray schedule beginning at budbreak and continue on a 10-14 day spray schedule through the pollination period. Alone or combination with labeled rates of Super Tin 80WP or Elast 400F. Apply the higher specified rate to scab-susceptible varieties or when weather conditions favor severe scab disease.
Specific Use Restrictions: 1) Do not make more than 4 applications per year. 2) Do not apply after shucks split. 3) Do not apply more than 16.5 pints per acre per year. 4) Do not feed or graze cover crops in treated areas to livestock. 5) Do not add a surfactant when tank mixing. * Not for use in California for anthracnose, brown leaf spot, downy spot, liver spot, vein spot, and zonate leaf spot.			

CROP	DISEASE	RATE	INSTRUCTIONS
PISTACHIOS	Botryosphaeria Panicle and shoot Blight Alternaria late blight	4 pints/acre	Apply in 100 gallons of water per acre by ground or 5 gallons or more diluent by air. Begin applications at 25-50% bloom and repeat on a 10-14 day interval to protect flowers and fruit. Make two additional pre-harvest sprays 50 and 35 days prior to harvest. Use good sanitation measures and control insect vectors to limit the spread of Botryosphaeria. Alternaria: Start applications in June and continue on a 21 day schedule. Continue until early August.
Specific Use Restrictions: 1) Do not make more than 4 applications per season. 2) Do not apply more than 16 pints per acre per season. 3) The PHI is 35 days.			

CROP	DISEASE	RATE	INSTRUCTIONS
SOYBEAN*	Soybean rust (<i>Phakopsora pachyrizi</i>)	1.5-2 pints/acre	Apply the higher specified rate (2 pts/acre) when conditions favor severe disease development or VIATHON® may be mixed with a strobilurin fungicide to improve control when conditions favor severe disease.
	Downy mildew (<i>Peronospora manshurica</i>) Powdery mildew (<i>Microsphaera diffusa</i>) Frogeye leafspot (<i>Cercospora sojina</i>) Aerial blight (<i>Rhizoctonia solani</i>) Brown Leaf spot (<i>Septoria glycines</i>)	2 pints/acre	<p>For rust, apply preventively when weather conditions favor disease development. Repeat applications on a 7-14 day interval.</p> <p>Make applications between R1 and R5 (flowering through visible seed in the pod in the upper 4 nodes of the plant). Make a second application 21 days later if necessary.</p> <p>Use spray volumes of 2 gallons or more by air, and 10 gallons or more by ground.</p> <p>To prevent mixing issues, add BLENDX® VHC at 0.25% (1 quart/100 gallons). Keep under constant agitation during spraying. Use adjuvants at their lowest labeled rate.</p> <p>If a strobilurin mixing partner is applied, determine the physical compatibility of this product with VIATHON®. Use a jar test as described in the Compatibility section of this label. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.</p>
<p>Specific Use Restrictions:</p> <p>1) Do not make more than 2 applications per year.</p> <p>2) Do not apply more than 4 pints/0.225 pounds of the active ingredient tebuconazole per acre per year.</p> <p>3) The PHI is 21 days.</p> <p>4) When tank mixing, no label dosage rate may be exceeded, and the most restrictive label precautions and restrictions must be followed.</p> <p>* Not for use in California, New York and Washington.</p>			

CROP	DISEASE	RATE	INSTRUCTIONS
STONE FRUIT Apricot, sweet cherry, tart cherry, nectarine, peach, plum, chickasaw plum, damson plum, Japanese plum, plumcot, fresh prune	Blossom blight (<i>Monilinia laxa</i> , <i>Monilinia fructicola</i>)	2-4 pints/acre	Blossom blight: Start applications at white bud on cherry, pink bud on peach and nectarines, and at green tip on plums and prunes. Make applications at 50% bloom and petal fall.
	Cherry leaf spot	4 pints/acre	Cherry leaf spot, powdery mildew, rust: Begin applications at petal fall and continue on a 7 day interval early in the season during rapid growth, then move to a 14 day interval. A post-harvest application may be made to control overwintering inoculum.
	Powdery mildew	2-4 pints/acre	
	Rust	2.5-4 pints/acre	On peach, prune, and plum, begin applications at canker emergence and continue on a 14 day interval as needed.
	Brown rot	2-4 pints/acre	Brown rot: Begin applications 3 weeks before harvest and rotate with other registered fungicides on 7 day intervals through harvest. As VIATHON ® contains a DMI fungicide, it is best alternated to address fungicide resistance development.
Specific Use Restrictions: 1) Do not make more than 6 applications per year. 2) Do not apply more than 12 pints per acre per year. 3) There is a 0-day PHI. 4) Do not apply by air.			

CROP	DISEASE	RATE	INSTRUCTIONS
SUNFLOWERS	Rust	2-3 pints/acre	<p>Apply at the earliest sign of infection, or when weather conditions favor rust development. Apply the higher specified rate on susceptible varieties and/or severe disease conditions. Application may be repeated at 14 days to maintain control. Apply in 20 gallons spray by ground and 5 gallons spray by air.</p> <p>For optimum disease control, use the lowest rate of an approved spray surfactant in the spray. The spray must have two to four hours of drying time on the plant for VIATHON® to move into the plant.</p>

Specific Use Restrictions:
 1) Do not make more than 4 applications per year.
 2) Do not apply more than 8 pints per acre per year.
 3) The PHI is 50 days.

CROP	DISEASE	RATE	INSTRUCTIONS
WATERMELON	Powdery mildew*	2-3 pints/acre	<p>Apply the specific dose in a protective spray schedule to foliage and fruit. Repeat on a 10-14 day interval.</p>
	Gummy stem blight (<i>Didymella bryonae</i>) suppression	4 pints/acre	
	Downy mildew*	4 pints/acre	

Specific Use Restrictions:
 1) Do not make more than 4 applications per year.
 2) Do not apply more than 12 pints/acre per season.
 3) The PHI is 7 days.
 * Not for use in California for powdery mildew and downy mildew.

CROP	DISEASE	RATE	INSTRUCTIONS
WHEAT	<p>Rusts, leaf, stem*, and stripe (<i>Puccinia</i> spp.)</p> <p>Head blight* or scab* (<i>Fusarium</i> spp.) suppression</p> <p>Septoria Glume Blotch* (<i>Staganospora nodorum</i>)</p> <p>Powdery mildew* (<i>Erysiphe graminis</i>/ <i>Blumeria graminis</i>)</p> <p>Tan spot* (<i>Pyrenophora tritici-repentis</i>) Septoria Leaf Spot* (Septoria complex)</p>	2 pints/acre	<p>Scout fields for symptoms. Apply at earliest sign of disease. For head blight, application is typically best made at growth stage 10.5.</p> <p>Rusts, leaf, stem, and stripe: Apply at first sign of pustules.</p> <p>Head blight (suppression): Apply at beginning of flowering (Feekes 10.51).</p> <p>Septoria Glume Blotch: Apply when 50% of the heads on the main stem are in flower.</p> <p>Powdery mildew: Apply at the onset of disease.</p> <p>Tan spot and Septoria Leaf Spot: Apply at flag leaf emergence, or at the first sign of disease until end of flowering.</p> <p>Monitor fields carefully for disease, especially during times of weather conducive for disease. Apply a surfactant at the lowest label rate to optimize control.</p>
<p>Specific Use Restrictions:</p> <p>1) Grazing or feeding forage to livestock is permitted 6 days after the last application.</p> <p>2) Do not apply within 30 days of harvest. Straw may be used for bedding.</p> <p>3) Do not make more than 1 application per season.</p> <p>4) Do not apply more than 2 pints per acre per season.</p> <p>* Not for use in California for stem rust, head blight or scab, Septoria glume blotch, powdery mildew, tan spot, or Septoria leaf spot.</p>			

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry, secure, cool place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture or dispose of in a sanitary landfill, or by incineration. Do not burn unless allowed by state and local ordinances. If burned, stay out of smoke.

WARRANTY LIMITATIONS AND DISCLAIMER

To the extent consistent with applicable law, Helena Agri-Enterprises, LLC warrants that at the time of delivery, the product will conform to its chemical description on the label, that it will pass without objection in the trade under the contract description, that seller will convey good title thereto, and that such product will be delivered free from any lawful security interest, lien or encumbrance.

This is the only warranty made on this product. Helena Agri-Enterprise, LLC EXPRESSLY DISCLAIMS ANY IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE AND, EXCEPT AS SET FORTH IN THE ABOVE PARAGRAPH, ANY OTHER EXPRESS OR IMPLIED WARRANTIES. To the extent consistent with applicable law, buyer assumes all risk of use of this material when such use is contrary to label instructions. Read and follow the label directions carefully.

VIATHON® and **BLENDEX®** are registered trademarks of Helena Holding Company.

Super Tin (triphenyltin hydroxide) is a registered trademark of United Phosphorus Ltd.

Elast (dodine) is a registered trademark of Aceto Agricultural Corp.



A systemic fungicide for the control of apple, almond, asparagus*, barley*, bean* (except succulent), corn*, grape, grasses grown for seed, hop*, leafy brassica green*, onion, peanut*, pecan, pistachio, soybean*, stone fruit, sunflower, walnut, watermelon, and wheat diseases

ACTIVE INGREDIENTS:

Potassium phosphite¹ 49.0%

Tebuconazole: α -[2-(4-chlorophenyl)ethyl]- α - (1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol² 3.3%

OTHER INGREDIENTS : **47.7%**

TOTAL: **100.0%**

¹ 30.4% (3.66 lbs/gal) phosphorous acid equivalent

² 0.42 lbs/gal tebuconazole

*Not for use in California

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical information.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. For additional information, call: 1-800-424-9300.	

SHAKE WELL BEFORE USE
FOR CHEMICAL SPILL, LEAK, FIRE OR EXPOSURE CALL TOLL FREE: 1-800-424-9300.

AGRICULTURAL CHEMICAL: DO NOT SHIP OR STORE WITH FOODS, FEEDS, DRUGS OR CLOTHING.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

USER SAFETY RECOMMENDATIONS:

Users should: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or tribal agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry, secure, cool place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture or dispose of in a sanitary landfill, or by incineration. Do not burn unless allowed by state and local ordinances. If burned, stay out of smoke.

EPA Reg. No. 42519-35-5905

EPA Est. No. 5905-GA-1 **NET CONTENTS: 2.5 Gallons**

Manufactured For
HELENA AGRI-ENTERPRISES, LLC
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TENNESSEE 38017 U.S.A. • (901) 761-0050

5043-9/190625 19/19
AD 020618HAE

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING