### SUPPLEMENTAL LABELING

# RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

TO BE USED BY CERTIFIED APPLICATORS ONLY - NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR. EXCEPT THAT UNCERTIFIED PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

# Syngenta Crop Protection, LLC

P. O. Box 18300 Greensboro, North Carolina, 27419-8300 SCP 1652A-S8A 0822

> PARAQUAT DICHLORIDE GROUP 22 HERBICIDE

Gramoxone® SL 3.0 HERBICIDE

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

### **Active Ingredients:**

Paraquat dichloride\*..... Other Ingredients: 56.1% 100.0%

Total:

\*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

# DIRECTIONS FOR USE RESTRICTED USE PESTICIDE

Cucurbits and Cabbage, OH Only

Crops (Including all cultivars, varieties, and/or hybrids of these)					
Cabbage	Melons				
Cucumber	Squash				
Target	Rate	Application Timing	Use Directions		

Cucumber			quasn
Target Weed	Rate pt/A	Application Timing	Use Directions
Refer to Section 7.0.	1.5	Postemergence (Directed/Shielded Spray)	For use in Ohio Only  Use for control or suppression of emerged weeds between rows after crop establishment.  Apply as a ground directed spray in a minimum of 20 gal/A spray volume when weeds are succulent and weed growth is less than 6".  Use precision directed spray equipment adjusted to prevent spray contact with crop plants.  Always use a high quality non-ionic surfactant.  Do not exceed 30 psi nozzle pressure or spray under conditions which may cause excessive drift.  See Section 4.4.4 of the Gramoxone SL 3.0 label for adjuvant recommendations.

### **Tank Mix Options:**

 Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on cucurbits and cabbage applied by the same methods and at the same timings. Refer to Section 4.4.2 for tank-mix precautions in the Gramoxone SL 3.0 label.

# **Resistance Management:**

Refer to Section 3.1.

### Precaution:

Do not allow spray to contact green stems or foliage, as injury may result.

# **USE RESTRICTIONS**

- 1. Refer to **Section 6.1** for additional product use restrictions.
- 2. Maximum Single Application Rate: 1.5 pt/A
- 3. **REI:** 48 hours
- 4. Minimum Application Interval: 7 days
- 5. Maximum Annual Rate: 3.0 pt/A
  - a. **DO NOT** exceed 1.1 lb cation/A/year of paraquat-containing products.
- 6. **DO NOT** apply after bloom.
- 7. **DO NOT** make more than 3 applications per year.
- 8. **DO NOT** use on soils with less than 5% clay.
- 9. **DO NOT** apply this product through any type of irrigation system.
- 10. **DO NOT** allow animals to graze in treated areas.
- 11. Pre-Harvest Interval (PHI): Harvest crop at normal harvest maturity.

Gramoxone®, and the SYNGENTA Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for: Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, North Carolina, 27419-8300