



Section 24(c) Special Local Need Label

S-METOLACHLOR GROUP **15** HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Dual Magnum® Herbicide

For Control of Weeds in Highbush Blueberry, Blackberry, and Red and Black Raspberry

EPA Reg. No. 100-816
EPA SLN No. OR-110005

This label for Dual Magnum Herbicide is valid until December 31, 2026 or until otherwise amended, withdrawn, cancelled, or suspended.

Active Ingredient:

S-metolachlor (CAS No. 87392-12-9) 83.7%

Other Ingredients: 16.3%

Total: 100.0%

Dual Magnum is formulated as an Emulsifiable Concentrate (EC) and contains the equivalent of 83.7% or 7.62 lb of active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**SYNGENTA'S SPECIAL CONDITIONS, RISKS OF USE AND DISCLAIMER FOR USE OF
DUAL MAGNUM ON CROPS ON THIS 24(c) LABEL**

IMPORTANT- READ BEFORE USE

**THESE CONDITIONS RISKS OF USE AND DISCLAIMER ARE REQUIRED BY SYNGENTA
CROP PROTECTION LLC AND NOT SPECIFIED BY U.S. EPA OR THE OF STATE OF
OREGON**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SYNGENTA CROP PROTECTION,
LLC INTENDS THAT THE PRODUCT THAT IS THE SUBJECT OF THIS SECTION 24(c)
LABEL BE PURCHASED ONLY BY END USERS WHO AGREE BY ELECTRONIC
SIGNATURE ON SYNGENTA CROP PROTECTION'S INTERNET SITE TO THE TERMS AND
CONDITIONS REQUIRED BY SYNGENTA CROP PROTECTION, LLC INCLUDING A WAIVER
AND RELEASE FROM ALL LIABILITY AND INDEMNIFICATION BY THE USER AND/OR
GROWER OF SYNGENTA AND OTHERS FOR FAILURE TO PERFORM AND FOR CROP
INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS FROM USE OF DUAL MAGNUM**

ON CROPS ON THIS 24(c) LABEL. IF SUCH TERMS AND CONDITIONS ARE UNACCEPTABLE, RETURN THE DUAL MAGNUM AT ONCE UNOPENED OR USE THE DUAL MAGNUM FOR A DIFFERENT APPROVED USE IN ACCORDANCE WITH THE LABEL AFFIXED TO THE PRODUCT CONTAINER.

USE OF DUAL MAGNUM (THE “PRODUCT”) ON CROPS LISTED (THE “CROP”) FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP. SYNGENTA RECOMMENDS THAT THE USER TEST THIS PRODUCT TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.

Syngenta Crop Protection, LLC makes the Product available for use in the manner described in this Section 24(c) Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss.

The decision to use this Product in the manner described in this Section 24(c) Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors. Syngenta recommends that the user test this Product to determine its suitability for such intended use.

By purchasing the Product for use, or using the Product in the manner described in this Section 24(c) Labeling, you acknowledge and accept that, to the extent consistent with applicable law:

- 1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- 2) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Section 24(c) labeling;
- 3) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- 4) Syngenta Crop Protection, LLC disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- 5) these conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Syngenta Crop Protection, LLC or their respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Environmental Hazards: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Groundwater Advisory: S-metolachlor is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Surface Water Advisory: S-metolachlor has the potential to contaminate surface water through ground spray drift. Under some conditions, the active ingredient may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

Mixing/Loading Precautions

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates. Check-valves or antisiphoning devices must be used on all mixing and/or irrigation equipment.

- This product must not be mixed or loaded within 50 ft of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs.
- This product must not be mixed/loaded or used within 50 ft of all wells, including abandoned wells, drainage wells, and sink holes.
- Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad.
 - Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad.
 - Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained.
 - The pad shall be sloped to facilitate material removal.
 - An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad.
 - A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad.
- Containment capacities as described above shall be maintained at all times.

The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This FIFRA Section 24(c) label and the main EPA-registered Dual Magnum Herbicide label must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label.
- The risk of crop injury increases when adjuvants (e.g., non-ionic surfactants, crop oils, etc.), nitrogen sources (e.g., AMS, UAN), fertilizers or other pesticides are applied with Dual Magnum.
- **Rotational Crops:** Do not rotate to food or feed crops other than those listed on the main EPA-registered label.
- **Chemigation:** For use under this SLN label, do not apply this product through any type of irrigation system.

USE DIRECTIONS

Dual Magnum may be used for pre-emergence suppression or control of annual grass and small seeded broadleaf weeds. Dual Magnum will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

Rate Calculation for Banded Applications

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Dual Magnum rate per acre for a broadcast treatment} = \text{Amount of Dual Magnum needed per acre for a banded treatment}$$

HIGHBUSH BLUEBERRY

Apply Dual Magnum as a banded treatment to both sides of the blueberry row at 0.67 – 1.3 pt/A (0.64 – 1.24 lb ai/A) prior to weed emergence. The application should be directed to the soil surface in a 3 foot band on each side of the blueberry row to avoid direct contact with the crop foliage or crop injury may occur. When making the banded treatment, apply proportionally less spray mixture on the area actually treated. Use the lower end of the Dual Magnum rate range for soils that are relatively coarse textured and higher rates on fine textured soils.

Precautions: (1) Dual Magnum will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means. (2) Not all blueberry cultivars have been tested so injury may occur on some types of blueberry. Growers are encouraged to treat a few plants as a test and evaluate crop injury. (3) The risk of crop injury from Dual Magnum is greater at the higher use rates. (4) Blueberry plants that have been established for less than one year may be more sensitive to applications of Dual Magnum than those plants established for more than one year. For this reason, the lower rates are suggested on newer plantings. (5) Tank mixing Dual Magnum with other products (pesticides, adjuvants, etc.) will increase the risk for foliar crop injury.

Restrictions: (1) Do not make more than one application per crop per season. (2) Do not apply more than 1.3 pt/A (1.24 lb ai/A) of Dual Magnum per crop per season. (3) Do not harvest for 28 days following the Dual Magnum application. (4) Do not use on lowbush blueberry.

BLACKBERRY, RED AND BLACK RASPBERRY

Apply Dual Magnum in the early spring as a banded application to both sides of the plant row at 1.0 – 2.0 pt/A (0.95 – 1.91 lb ai/A) prior to weed emergence. The application should be directed to the soil surface to avoid direct contact with the crop foliage or crop injury may occur. When making the banded treatment, apply proportionally less spray mixture on the area actually treated. Use the lower end of the Dual Magnum rate range for soils that are relatively coarse textured and the higher rates on fine textured soils.

Precautions: (1) Dual Magnum will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means. (2) Not all caneberry types, cultivars or hybrids have been tested so injury from Dual Magnum may occur on some crops. (3) The risk of crop injury from Dual Magnum is greater at the higher use rates. (4) Application of Dual Magnum to crop foliage may result in crop injury and illegal residues. (5) Caneberry plants that have been established for less than one year may be more sensitive to applications of Dual Magnum than those plants established for more than one year. For this reason, the lower rates are suggested on newer plantings.

Restrictions: (1) Do not make more than one application of Dual Magnum per crop per season. (2) Do not apply more than 2.0 pt/A (1.91 lb ai/A) of Dual Magnum per crop per season. (3) Do not harvest for 28 days following the Dual Magnum application.

©2021

Dual Magnum® and the Syngenta logo are trademarks of a Syngenta Group Company.

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: OR0816136CA1121