



GUARANTEED ANALYSIS
Magnesium (Mg).....6.00%

Derived from: Magnesium Acetate.

KEEP OUT OF REACH OF CHILDREN



WARNING

Avoid breathing fertilizer mists. Avoid prolonged or repeated skin contact.
Avoid eye contact. Wash hands thoroughly after handling this product.

F1548

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL 1-866-944-8565.

GENERAL INFORMATION

FULLBACK® 6% MAGNESIUM is generally compatible with most pesticides and fertilizer solutions. Add to water before adding pesticides. Test for compatibility by means of a jar test prior to large scale tank mixing. FULLBACK 6% MAGNESIUM is intended for use as a plant nutrient supplement for the correction of magnesium deficiencies. FULLBACK 6% MAGNESIUM is intended for use on all agricultural, horticultural and ornamental crops that require and respond to magnesium nutritional supplements. FULLBACK 6% MAGNESIUM is to be applied directly to the soil as a spray or through drip or injection systems, or foliar application through conventional, low-volume, or aerial sprayers. FULLBACK 6% MAGNESIUM is compatible with a wide range of fertilizer materials, pesticides, herbicides and fungicides. FULLBACK 6% MAGNESIUM may be formulated into NPK fertilizers as a source of magnesium. This product should be stored in a cool, dry area out of direct sunlight.

NOTE: When used as directed this product does not supply all nutrients required by plants and is to supplement a soil fertility program based on soil tests.

EQUIPMENT

Brass, copper and galvanized metals may be subject to mild corrosion from prolonged or repeated contact with this product.

COMPATIBILITY

FULLBACK can be applied alone or with other compatible spray materials. Ammonium polyphosphate fertilizers tend to be more compatible than orthophosphate types. Including some polyphosphate in the mixture often improves compatibility of orthophosphates.

Proposed mixtures should be evaluated in a jar test before full scale use. Fertilizer based mixtures that also contain pesticides should be sprayed as soon as possible.

Note: When combining FULLBACK with phosphate fertilizers vigorous agitation is required for good product mixing. See mixing directions below for more information.

CROP SAFETY

This product has been tested on a number of crop varieties with no phytotoxicity observed at labeled rates; however, this product has not been tested on all varieties and cultivars. Due to the large number of plant species and their associated varieties or cultivars, and due to variable growing conditions, it is impossible to test every plant for tolerance to this product. It is the responsibility of the user to determine if this product can be used safely prior to commercial use. In a small test area, test the specified rates on plants for phytotoxicity prior to general use. Before using any tank mix, test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Treat a small group of target crop including plant cultivars and evaluate for 3 to 7 days prior to making an application to your entire crop.

APPLICATION DIRECTIONS

For best results, foliar applications of FULLBACK 6% MAGNESIUM should be made in early morning or late evening. Some foliar burning can result if applied during hot, dry daytime conditions. The recommended rates have been extremely successful for foliar usage, though phytotoxic symptoms may occasionally occur on sensitive crops. Consult your state Agricultural Experiment Station or Extension Agent for additional information on rates, timing of application and sensitivity of varieties or hybrids to phytotoxicity.

MIXING INSTRUCTIONS

1. Add FULLBACK 6% MAGNESIUM to water and start agitation.
2. Add pesticides or other nutrients.
3. Fill tank and agitate thoroughly.

RATES: DIRECTIONS FOR USE

Crop	Soil Applied Rate	Foliar Applied Rate	Application Directions and Timing
(01) Root and Tuber Vegetables including Potato, Sugarbeet, Carrot, etc.	1.0 to 8.0 pts/A	1.0 to 4.0 pts/A	1st application at tuber initiation. Can repeat as needed every 14 to 21 days.
(03) Bulb Vegetables including Onion, etc.	1.0 to 8.0 pts/A		1st application after thinning, or 3- to 4-leaf stage. Can repeat as needed every 14 to 21 days.
(04) Leafy Vegetables including Celery, Head and Leaf Lettuce, etc.	1.0 to 8.0 pts/A		1st application after thinning, or 3- to 4-leaf stage. Can repeat as needed every 14 to 21 days.
(05) Brassica Vegetables including Broccoli, Cabbage, Cauliflower, Collards, Brussels Sprouts, etc.	1.0 to 8.0 pts/A	1.0 to 4.0 pts/A	1st application after thinning or 3- to 4-leaf stage. Can repeat as needed every 14 to 21 days.
(06) Legume Vegetables including all Beans, Soybeans, etc.	1.0 to 8.0 pts/A		1st application at v3 to v4. Can repeat as necessary every 14 to 21 days until senescence.
(08) Fruiting Vegetables including Peppers, Tomato, Eggplant, etc.	1.0 to 8.0 pts/A		1st application at crop establishment. Can repeat every 14 to 21 days until harvest.
(09) Cucurbits including all Melons, Cucumber, Squash, etc.	1.0 to 8.0 pts/A		1st application at 3- to 4-leaf stage. Can repeat as necessary every 14 to 21 days until harvest.
(10) Citrus Fruits including Oranges, Lemons, Limes, etc.	1.0 to 8.0 pts/A		1st application at 1st bloom. Can repeat as necessary every 14 to 21 days until harvest.
(11) Pome Fruit including Apple, Pear, etc.	1.0 to 8.0 pts/A	1.0 to 4.0 pts/A	1st application at first bloom. Can repeat every 14 to 21 days until harvest.
(12) Stone Fruit including Cherry, Peach, Plum, etc.	1.0 to 8.0 pts/A	1.0 to 4.0 pts/A	1st application at first bloom. Can repeat as necessary every 14 to 21 days until harvest.
(13) Berry and Small Fruit including Strawberry, Blueberry, Caneberries (Blackberry, Raspberry)	1.0 to 8.0 pts/A	1.0 to 4.0 pts/A	1st application at first bloom. Can repeat as necessary every 14 to 21 days until harvest.
(13) Berry and Small Fruit including Grape	1.0 to 8.0 pts/A		1st application when 4 to 6 inches of new shoot growth

(14) Tree Nuts including Almond, Hazelnut, Pistachio, Pecan, Walnut	1.0 to 8.0 pts/A	1.0 to 8.0 pts/A	1st application at first bloom. Can repeat as necessary every 14 to 21 days until harvest. Spray canopy "to wet"
(15) Cereal Grains including Corn, Wheat, Grain Sorghum, Rice, etc.	1.0 to 8.0 pts/A		1st application at v3 to v5. Can repeat as necessary every 14 to 21 days until pollination. Small grains: 1st application timing at 4- to 6-leaf (tillering). 2nd application may be made up to the full flag leaf stage.

Directions for Use cont'd.

Crop	Soil Applied Rate	Foliar Applied Rate	Application Directions and Timing
(16) Forage Fodder and Straw of Cereal Grains	1.0 to 8.0 pts/A	1.0 to 8.0 pts/A	Begin on emergence of spring growth or deficiency symptoms. Repeat every 14 to 21 days as needed.
(17) Grass forage Fodder and Hay including Grasses Grown for Seed	1.0 to 8.0 pts	1.0 to 8.0 pts/A	Begin on emergence of spring growth or deficiency symptoms. Repeat every 14 to 21 days as needed.
(18) Animal Feed, Non Grass including all Alfalfa, Clover, etc.	1.0 to 8.0 pts		Apply 1st application at 2nd to 4th true leaf stage and 1 to 2 subsequent applications after cutting.
(20) Oilseed including Cotton, Sunflower, Canola	1.0 to 8.0 pts		Begin applications at 2- to 4-leaf stage. Repeat applications at 14- to 21-day intervals as needed.
(99) Misc including Avocado, Artichoke, Hops, Flax, Mint, Olive, Peanut, Tobacco, etc.	1.0 to 8.0 pts		1st application at first bloom. Can repeat as necessary every 14 to 21 days until harvest.

STORAGE AND DISPOSAL

STORAGE: Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. Do not contaminate water, food, or feed by storage or disposal.
DISPOSAL: Do not reuse containers. Triple rinse (or equivalent) adding rinse water to spray tank, then offer for recycling at an ACRG site (go to <http://www.acrecycle.org/> for locations) or by reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by appropriate authorities. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food or feed storage or disposal.

HANDLING: Avoid breathing fertilizer mists. Avoid prolonged or repeated skin contact. Avoid eye contact. Wash hands thoroughly after handling this product.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary. LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in accordance with such Directions for Use under normal conditions of use. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.regulatory-info-lpi.com>

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

Fullback is a registered trademark of Loveland Products, Inc.



WARNING

This product can expose you to chemicals including nickel, which are known to the State of California to cause cancer. For more information go to www.P65Warnings.ca.gov.

Density: 10.12 lbs/gallon @ 68 °F

Liquid Measure: 265 gallons (1,003.13 liters)

Net Weight: 2,681.80 lbs (1,216.44 kg)

Product Code: 1000174486

Density: 10.12 lbs/gallon @ 68 °F

Liquid Measure: gallons (liters)

Net Weight: lbs (kg)

Product Code: 1000174488

GUARANTEED BY: LOVELAND PRODUCTS, INC.® P.O. BOX 1286 GREELEY, COLORADO 80632-1286