

FIFRA Section 24(c) Special Local Need Label



Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid 2F

EPA Reg. No. 62719-442

EPA SLN No. CA-070015

For Distribution and Use Only in the State of California

Control of Citrus Leafminer in Non-Bearing Citrus (Including Nurseries)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this FIFRA 24(c) Special Local Need labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Directions for Use

Refer to product labeling for Use Precautions and Restrictions, Mixing and Application instructions.

Application Rates, Application Timing and Restrictions:

Application Rate (fl oz/acre)	Application Timing	Restrictions
6 - 12 (0.09 – 0.18 lb ai/acre)	Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Best control is attained with two consecutive applications of Intrepid 2F at 7- to 14-day intervals. After two successive applications of Intrepid 2F, rotate to another insecticide in a different chemical class for at least 30 days before making additional applications of Intrepid 2F. After switching to an alternative MOA for 30 days or more, two more applications of Intrepid 2F may be made at 7- to 14-day intervals.	<ul style="list-style-type: none">• Do not apply more than 12 fl oz of Intrepid 2F (0.18 lb ai) per acre per application.• Do not make more than 4 application per calendar year or apply more than 48 fl oz of Intrepid 2F (0.72 lb ai) per calendar year.• Do not use on fruit bearing trees.

Ground Application: Apply a minimum of 50 gallons per acre by conventional ground equipment to trellised trees or trees 10 feet tall or less. For trees greater than 10 feet tall, use a minimum of 100 gallons per acre. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 20 gallons per acre. Intrepid 2F can be aerially applied when conditions warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Resistance Management: Do not apply Intrepid 2F more than two times in succession to any single generation of citrus leafminer or more than two times within any 30-day period. After use of Intrepid 2F (up to two times) in any 30-day period, rotate to another class of chemistry with a different mode of action or use no insecticide for the next 30 days. Do not make more than four applications of Intrepid 2F to non-bearing citrus in a calendar year.

Spray Adjuvants: The use of 1% NR-415 oil or similar product has been shown to improve penetration, coverage and retention of the spray material.

Note: Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

™®Trademarks of Corteva Agriscience and its affiliated companies

R846-114

Approved: 10/19/2023

Replaces R846-050