

# Supplemental Labeling

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Intrepid® 2F

EPA Reg. No. 62719-442

This supplemental label expires on 10/31/2025 and must not be used or distributed after this date.

### Control of Armyworm in Rice For Distribution and Use Only in the State of California

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid® 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

#### Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

#### Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
armyworm	8 - 10 (0.12 – 0.16 lb ai/A)	Begin applications when first signs of feeding damage appear or when threshold levels of feeding damage occur. Use a higher rate for heavier infestations and under conditions in which thorough coverage is more difficult	<ul style="list-style-type: none"> <li>• <b>Preharvest Interval:</b> Do not apply within 14 days of harvest</li> <li>• Do not apply more than a total of 20 fl oz of Intrepid 2F (0.32 lb ai) per acre per year</li> <li>• Do not make more than two applications per year</li> <li>• Hold treated water for at least 7 days after treatment</li> <li>• <b>Minimum Re-treatment Interval:</b> 10 days</li> </ul>

**Ground Application:** Apply in a minimum of 5 gpa by conventional ground equipment. Use a spray volume that assures uniform coverage of the infested portions of the treated crop. Calibrate equipment to the desired spray volume.

**Aerial Application:** Apply in a minimum of 5 gpa. Use a higher carrier volume for heavy infestations and in situations where thorough coverage is difficult to achieve. Aerial applicators should evaluate conditions existing at the time of application and make appropriate adjustments to reduce drift.

### **Spray Adjuvants**

The addition of agricultural adjuvants to sprays of Intrepid 2F may improve initial spray deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. When an adjuvant is to be used with this product, Corteva Agriscience recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Always add adjuvants last in the mixing process.

**Note:** Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

™®Trademarks of Corteva Agriscience and its affiliated companies

R846-106

EPA accepted 10/12/2022

Initial printing.