

MONSANTO COMPANY  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167 USA

# Roundup PowerMAX® II Herbicide

## Recommendations Prior to Transplanting Cucurbits and Fruiting Vegetables in Sandy Soils

EPA Reg. No. 524-537

### ***FIFRA 2(ee) Recommendation***

## **FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, FLORIDA AND GEORGIA**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin is not product labeling, but is issued to clearly describe use recommendations as permitted under FIFRA Section 2(ee). Always read and follow label directions. The applicable labeling for this product must be in the possession of the user at the time of application.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Read the product labeling affixed to the container of Roundup PowerMAX II Herbicide before applying. Use of Roundup PowerMAX II Herbicide according to this product bulletin is subject to the use precautions and limitations imposed by the labeling affixed to the container.

**CROPS:** Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Gherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all), Muskmelon, Persian melon, Pumpkin, Squash (summer, winter), Watermelon, Eggplant, Groundcherry, Okra, Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato.

When applying Roundup PowerMAX II Herbicide prior to transplanting these crops in bare ground production soil with over 85% sand and/or less than 0.5% organic matter, if the soil is not tilled after application and before planting, apply no more than 32 fluid ounces of this product per acre in a single application, and allow for a minimum accumulation of 0.5 inch of rainfall or overhead irrigation and wait 7 or more days between application and transplanting. Make no more than 1 application of this product within 2 weeks before transplanting.

When applying Roundup PowerMAX II Herbicide prior to transplanting these crops in mulch production where soil is over 85% sand and/or less than 0.5% organic matter, wait 3 or more days before transplanting following a single application of this product up to 32 fluid ounces per acre, or wait 10 or more days following a single application between 32 and 64 fluid ounces per acre, AND allow for a single rainfall or irrigation event of at least 0.5 inch between application of this product at any rate and transplanting. Punch new transplant holes and place plants a minimum of 3 inches from old holes or torn mulch.

Single Application Rate of Roundup PowerMAX II Herbicide	Interval Between Application and Transplanting	Rainfall/Irrigation Before Transplanting
Up to 32 fluid ounces	3 or more days before transplanting	0.5 inch or more
32 to 64 fluid ounces	10 or more days before transplanting	0.5 inch or more

For product information or assistance using this product, call toll-free 800-332-3111.  
In case of an emergency involving this product, call collect, day or night, 314-694-4000.

Roundup PowerMAX is a registered trademark of Monsanto Technology LLC.

©2020 MONSANTO COMPANY

1/27/2020

63045U2-13