FIFRA §2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH DAKOTA

Sonalan® HFP

For aerial applications on dry beans, sunflower and rapeseed (canola varieties only)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

DRY BEANS

(Does not include soybean)

Sonalan HFP - Alone

Apply and incorporate Sonalan HFP in the spring before planting or in the fall. Fall application may be made only with dry bulk fertilizer. See instructions for fall application of Sonalan HFP under the heading Application Timing in the Product Information section of this label. Follow soil preparation, application, and incorporation procedures in the Product Information section of this label.

Broadcast Application Rates for General Weed Control Except Groundcherry and Nightshade:

Soil Texture	Sonalan HFP (pt/acre)
coarse	1 ½ - 2
medium	2 – 2 ½
fine	2 ½ - 3

Broadcast Application Rates for General Weed Control Including Groundcherry and Nightshade From Seed:

Soil Texture	Sonalan HFP (pt/acre)
coarse	3 - 3 1/2
medium	3 ½ – 4
fine	4 – 4 ½

Two incorporation passes are required for control of ground cherry and eastern black nightshade.

Precaution: When high rates for control of nightshade and groundcherry are applied, rotate only to crops listed on this label.

Chemigation: Sonalan HFP may be applied to dry beans by chemigation at labeled rates. Refer to Chemigation Application section for application guidelines for chemigation.

Aerial applications: Sonalan HFP may be applied to dry beans by air at labeled rates.

SUNFLOWER

Sonalan HFP - Alone

Apply and incorporate Sonalan® HFP herbicide in the spring before planting or in the fall. Fall application may be made only with dry bulk fertilizer. See instructions for fall application of Sonalan HFP under the heading Application Timing in the Product Information section of this label. Follow soil preparation, application, and incorporation procedures specified in the Product Information section of this label.

Broadcast Application Rates for General Weed Control Except Groundcherry and Nightshade:

Soil Texture	Sonalan HFP (pt/acre)
coarse	1 ½ - 2
medium	2 - 2 ½
fine	2 ½ - 3

Broadcast Application Rates for General Weed Control Including Groundcherry and Nightshade From Seed:

broadcast Application Nates for Ceneral Weed Control including Croundenerry and Hightshade From Ceed.	
Soil Texture	Sonalan HFP (pt/acre)
coarse	3 – 3 ½
medium	3 ½ - 4
fine	4 – 4 ½

Chemigation: Sonalan HFP may be applied to sunflower by chemigation at labeled rates. Refer to Chemigation Application section for application guidelines for chemigation.

Aerial applications: Sonalan HFP may be applied to sunflowers by air at labeled rates.

Precautions and Restrictions:

- Do not graze or forage crop grown in treated soil or cut for hay or silage.
- · When high rates for control of nightshade and groundcherry are applied, rotate only to crops listed on this label.

RAPESEED (CANOLA VARIETIES ONLY)

Sonalan HFP - Alone

Apply and incorporate Sonalan® HFP herbicide before planting. Application may be made in the fall or in the spring prior to planting canola or crambe. See instructions for fall application, application directions and soil incorporation directions in the Product Information section of this label.

Use the correct rate for each soil type to avoid crop injury. Prolonged wet or cold soils, deep seed placement, soil crusting, or application overlaps may combine to cause slowed or reduced emergence.

Broadcast Application Rates:

Soil Texture	Sonalan HFP (pt/acre)
coarse	1 ½

medium	2
fine	2 ½

Tank Mix or Overlay Recommendations

For broader spectrum weed control, other products registered for use in canola or crambe may be applied in tank mix combination or as a sequential overlay Consult the manufacturer's label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the Product Information section of this label.

Aerial applications: Sonalan HFP may be applied to rapeseed (canola varieties only) by air at labeled rates.

Restriction:

• Do not graze or forage crop grown in treated soil.

19-05-02



Produced For: Gowan Company P.O. Box 5569 Yuma, AZ 85366

EPA Reg. No. 10163-356 EPA Est. No.