

AZOXYSTROBIN	GROUP	11	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE



Active Ingredients:	
Azoxystrobin*	19.8%
Difenoconazole**	19.8%
Other Ingredients:	60.4%
Total:	100.0%

*CAS No. 131860-33-8

**CAS No. 119446-68-3

Formulated as a suspension concentrate (SC) containing 1.88 lb of azoxystrobin active ingredient and 1.88 lb of difenoconazole active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN. CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOTLINE NUMBER	
In case of emergency, call ChemTrec at 1-800-424-9300, available 24/7.	

See additional precautionary statements and directions for use inside booklet.

EPA REG. NO. 5905-646

AD 111424HAE

EPA EST. NO. First Letters of Product Batch Code Indicate Producing Establishment,
5905-AR-1=WA • 5905-GA-1=CG • 5905-CA-1=KC • 5905-IA-1=DI

Manufactured For
HELENA AGRI-ENTERPRISES, LLC
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TENNESSEE 38017 U.S.A. • (901) 761-0050

Job 245120

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION/PRECAUCION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Socks and shoes, and
- Chemical resistant gloves such as; Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq 14 mils, Polyvinyl Chloride (PVC) \geq 14 mils, and Viton \geq 14 mils.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Physical and Chemical Hazards

Do not mix or allow coming into contact with oxidizing agents. Hazardous chemical reaction may occur.

Environmental Hazards

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer.

Difenoconazole is toxic to fish, mammals and aquatic invertebrates. Drift and runoff may be hazardous to estuarine/ marine organisms in water adjacent to treated areas.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of Azoxystrobin and a degradate of Azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Helena immediately if you observe any adverse environmental effects due to use of this product.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants,
- Socks and shoes, and
- Chemical resistant gloves such as: Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq 14 mils, Polyvinyl Chloride (PVC) \geq 14 mils, and Viton \geq 14 mils.

PRODUCT INFORMATION

MAKO is a broad-spectrum product containing two systemic fungicides. It has preventative and curative properties and is recommended for the control of many important plant diseases. MAKO provides excellent disease control of many leaf spots and powdery mildews. MAKO is applied as a foliar spray and can be used in block, alternating spray or tank-mix programs with other crop protection products. All applications must be made according to the use directions that follow.

USE PRECAUTIONS AND RESTRICTIONS

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

ATTENTION

MAKO is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray MAKO where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State Extension Agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply MAKO to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

USE INFORMATION

Application: Thorough coverage is necessary to provide good disease control. Make no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Adjuvants: When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is recommended.

Use of Adjuvants: Under certain weather conditions (particularly high temperatures) MAKO in combination with high rates of silicone-based or oil containing (petroleum or crop) additives or adjuvants may cause injury. Do not exceed 0.125% adjuvant (v/v). Consult a Helena representative for more information concerning additives or adjuvants.

A tank mixture with Dimethoate may cause crop injury.

On fresh market tomatoes, do not use adjuvants or tank mix MAKO with any EC product.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if the maximum amount of MAKO has been used. If resistant isolates to Group 11 or group 3 fungicides are present, efficacy can be reduced for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, with highly susceptible varieties, or when environmental conditions are conducive to disease.

Integrated Pest Management (IPM): MAKO should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established for your area. MAKO may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

AZOXYSTROBIN	GROUP	11	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE

For resistance management, please note that MAKO contains both azoxystrobin, a strobilurin fungicide in Group 11 and difenoconazole, a triazole fungicide in Group 3. Any fungal population may contain individuals naturally resistant to either or both of the active ingredients in MAKO and other Group 11 or Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

As part of a resistance management strategy:

- Apply a maximum of 4 sprays during one crop cycle.
- Apply no more than 2 sequential applications unless otherwise stated in the crop section.
- Rotate the use of MAKO or other Group 11 and Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Helena Agri-Enterprises, LLC at 1-901-761-0050. You can also contact your pesticide distributor or university extension specialist to report resistance.

Rotational Crops: Please see the following table for the crop rotational restrictions:

Rotational Crops	Planting Time From Last MAKO Application
Berry, Low Growing, Subgroup 13-07G Brassica (Cole) Leafy Vegetables Crop Group 5 Bulb Vegetables, bulb onion Subgroup 3-07A and green onion Subgroup 3-07B Carrots Chickpeas Citrus Fruit Crop Group 10-10 Cotton Subgroup 20C Cucurbit Vegetables Crop Group 9 Fruit, small, vine climbing Subgroup 13-07F, except fuzzy kiwifruit Fruiting Vegetables Crop Group 8-10 Guava Papaya Okra Pepper Potatoes Rice Soybeans Stone Fruit Crop Group 12-12 Strawberries Sugar Beets Tomatoes Tree Nuts Crop Group 14-12 Tuberous & Corm Vegetable Subgroup 1C Watercress Wild Rice	0 days
Cereals (Wheat, Barley, Oats, Rye, Triticale) Root and Tuber Vegetables, Crop Group 1 (except Carrot, Sugar Beet, and Tuberous Corm Vegetable Subgroup 1C)	30 days
Buckwheat Millet	365 days
All Other Crops Intended for Food and Feed	60 days

Crop Tolerance: Plant tolerance has been found to be acceptable for all crops on the label, however, not all possible tank-mix combinations have been tested under all conditions. When possible, it is recommended to test the combinations on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application. See USE PRECAUTIONS AND RESTRICTIONS regarding apple phytotoxicity.

Greenhouse Use: For resistance management, do not use MAKO for transplant production.

Spray Drift Management: To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

MIXING AND APPLICATION METHODS

Spray Equipment

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Nozzles should be the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Screens placed on suction side of pump should be 16-mesh or coarser.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's recommendations.

Pump

- Use a pump with capacity to:
 - Maintain 35-40 psi at nozzles.
 - Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturer's and state recommendations.

For specific local directions and spray schedules, consult the current state agricultural recommendations.

Mixing Instructions

- MAKO is a suspension concentrate (SC) formulation.
- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

MAKO Alone (No Tank Mix)

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add MAKO to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after MAKO has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

MAKO + Tank Mixtures: MAKO is usually compatible with all tank-mix partners listed on this label. To determine the physical compatibility of MAKO with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and MAKO to the spray tank.
- Allow MAKO to completely disperse.
- Spray the mixture with the agitator running.
- Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label.

- Label dosage rate must not be exceeded, and the most restrictive label precautions and limitations must be followed.
- This product must not be mixed with any product which prohibits such mixing.
- It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Application Instructions

MAKO may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

Ground Application

- Apply in a minimum of 10 gal of water per acre, unless specified otherwise.
- Do not apply through any ultra-low volume (ULV) spray system.
- Thorough coverage is necessary to provide good disease control.

Aerial Application

- Use only on crops where aerial applications are indicated.
- Thorough coverage is necessary to provide good disease control.
- Apply in a minimum of 5 gallons of water per acre unless specified otherwise.
- Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.
- Do not apply directly to humans or animals.
- Do not apply through any ultra-low volume (ULV) spray system.

ATTENTION

MAKO is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray MAKO where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State Extension Agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply MAKO to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.1-0.25 inches/acre. Excessive water may reduce efficacy.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.

3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Center Pivot Irrigation Equipment

Notes: (1) Use only with drive systems which provide uniform water distribution. (2) Do not use end guns when chemigating MAKO through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply $\frac{1}{8}$ – $\frac{1}{2}$ inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. When applying MAKO through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of MAKO required to treat the area covered by the irrigation system.
- Add the required amount of MAKO and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the MAKO solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the MAKO solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20- to 30-minute interval. When applying MAKO through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of MAKO required to treat the area covered by the irrigation system.
- Add the required amount of MAKO into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the MAKO solution has cleared the last sprinkler head.

SPECIFIC INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

SPECIFIC CROP DIRECTIONS FOR USE

Cotton (Subgroup 20C)		
Target Diseases	Use Rate fl oz product/A	Remarks
Aerolate mildew <i>(Ramularia gossypii)</i> Alternaria leaf spot <i>(Alternaria spp)</i> Anthracnose <i>(Glomerella gossypii)</i> Ascochyta blight <i>(A.gossypii)</i> Boll rots <i>(Ascochyta gossypii,</i> <i>Alternaria spp.,</i> <i>Diplodia spp., Phoma spp.)</i> Cotton rust <i>(Puccinia schedonnardi)</i> Diplodia boll rot <i>(Diplodia spp.)</i> Hardlock <i>(Fusariumverticillioides)</i> Leafspots and blights <i>(Alternaria spp.,</i> <i>Ascochyta gossypii,</i> <i>Cercospora spp.,</i> <i>Stemphyllium spp.)</i> Southwestern cotton rust <i>(Puccinia cacabata,</i> <i>Puccinia spp.)</i> Stemphyllium leaf spot <i>(Stemphyllium spp.)</i> Target spot <i>(Corynespora cassiicola)</i>	6.0 -7.7*	For best activity, apply MAKO prior to or early in the disease development. An adjuvant may be added at specified rates. For foliar disease control, the first application should be targeted approximately at pin-head square to first bloom or when conditions are conducive for disease development. For best control of target spot, adjust the GPA to ensure coverage of upper and lower leaves. Subsequent applications may be made on a 14-21 day interval. Do not apply more than two sequential applications before alternating to a fungicide with a different mode of action.
*6.0 - 7.7 fl oz product/A contains 0.88 - 0.113 lb azoxystrobin/A and 0.88 - 0.113 lb difenoconazole/A.		
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. MAKO can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy. Applicators should use care in making applications near non-target aquatic habitats.</p>		
<p>Specific Use Restrictions:</p> <ul style="list-style-type: none"> • Do not apply more than 23.1 fl oz/A/year of MAKO (0.34 lb azoxystrobin and 0.34 lb difenoconazole). • Do not apply more than 0.45 lb ai/A/year of azoxystrobin-containing products. • Do not apply more than 0.34 lb ai/A/year of difenoconazole-containing products. • Do not apply MAKO within 45 days of harvest (45-day PHI). 		

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Soybean	Aerial Blight (<i>Rhizoctonia solani</i>) Alternaria Leaf Spot (<i>Alternaria spp.</i>) Anthracnose (<i>Colletotrichum truncatum</i>) Brown Spot (<i>Septoria glycines</i>) Cercospora Blight and Leaf Spot (<i>C. kikuchii</i>) Frogeye Leaf Spot (<i>Cercospora sojina</i>) Pod and Stem Blight (<i>Diaporthe phaseolorum</i>) Powdery Mildew (<i>Microsphaera diffusa</i>) Rust (<i>Phakopsora spp.</i>)	7.0 - 7.5*	Begin applications prior to disease onset when conditions are conducive for disease. Apply MAKO on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. The addition of a spreading/penetrating type adjuvant such as a non-ionic based surfactant or crop oil concentrate or blend is recommended. If disease pressure is high, use the shortest interval and highest rate.

*7 - 7.5 fl oz product/A contains 0.103-0.11 lb azoxystrobin/A and 0.103-0.11 lb difenoconazole/A.

Application: For best results, sufficient water volume must be used to provide thorough coverage. MAKO can be applied by ground, chemigation, or aerial application. For aerial applications, apply in a minimum of 2 gallons/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

- Do not apply more than 14.8 fl oz/A/year of MAKO (0.22 lb azoxystrobin and 0.22 lb difenoconazole).
- Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- Do not apply more than 0.22 lb ai/A/year of difenoconazole-containing products.
- Do not feed soybean hay, forage and silage to livestock.
- Do not apply within 14 days of harvest (14-day PHI).

MAKO Rate Conversion Table for Food Use

Fl oz product/acre	Lb ai azoxystrobin	Lb ai difenoconazole
4.5	0.067	0.067
5.6	0.082	0.082
6.5	0.096	0.096
7.0	0.103	0.103
7.5	0.110	0.110
7.7	0.113	0.113
8.0	0.118	0.118
8.3	0.122	0.122
8.5	0.125	0.125

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original container only. Store in a cool, dry and well-ventilated place. Protect from excessive heat. Keep container closed when not in use. Do not store near food or feed.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling (less than or equal to 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

**CONDITIONS OF SALE - LIMITED WARRANTY
AND LIMITATIONS OF LIABILITY AND REMEDIES**

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and must be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Agri-Enterprises, LLC (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. To the extent consistent with applicable law, the Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Agri-Enterprises, LLC's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent consistent with applicable law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2025.

AZOXYSTROBIN	GROUP	11	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE



MAKO™

Active Ingredients:	
Azoxystrobin*	19.8%
Difenoconazole**	19.8%
Other Ingredients:	60.4%
Total:	100.0%

*CAS No. 131860-33-8

**CAS No. 119446-68-3

Formulated as a suspension concentrate (SC) containing 1.88 lb of azoxystrobin active ingredient and 1.88 lb of difenoconazole active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN. CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOTLINE NUMBER	
In case of emergency, call ChemTrec at 1-800-424-9300, available 24/7.	

See additional precautionary statements and directions for use inside booklet.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION/PRECAUCION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.



Manufactured For
HELENA AGRI-ENTERPRISES, LLC
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TENNESSEE 38017 U.S.A. • (901) 761-0050

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original container only. Store in a cool, dry and well-ventilated place. Protect from excessive heat. Keep container closed when not in use. Do not store near food or feed.

Pesticide Disposal

Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling (less than or equal to 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

EPA REG. NO. 5905-646

AD 111424HAE

EPA EST. NO. First Letters of Product Batch Code Indicate Producing Establishment, 5905-AR-1=WA • 5905-GA-1=CG • 5905-CA-1=KC • 5905-IA-1=DI

NET CONTENTS:

PF 245120

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

OPM# 134192

PROOF

THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors** and **Dieline**.

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

PLEASE NOTE: Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors.

THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.

Dieline does not print.



4201 North Westport Ave. • Sioux Falls, SD 57107
Phone: (605) 978-0451 • Fax: (605) 978-0463

DATE	JOB NUMBER	CUSTOMER
2/27/25	245120	Helena
LABEL SIZE	BOOKLET SIZE	
6.375" X 6"	5.75" X 5.25"	
LABEL COLORS	BOOKLET OUTSIDE COLORS	BOOKLET INSIDE COLORS

PATTERN VARNISH: YES NO

Form: CS 006B - 3/29/2017

ARTWORK IS APPROVED

REVISED PROOF NEEDED

WE CANNOT PROCESS THIS ORDER WITHOUT AN AUTHORIZED SIGNATURE

Signed _____ Date _____