

Sale, use, and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.

SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 1420A-S2 1020

CYANTRANILIPROLE	GROUP	28	INSECTICIDE
-------------------------	--------------	-----------	--------------------

Fortenza®

Insecticide

A seed treatment product for protection against rice water weevil and suppression of grape colaspis on rice (wet seeded)

This supplemental label expires on 10/13/2023 and must not be used or distributed after this date.

Active Ingredient:

Cyantraniliprole ¹	48.8%
-------------------------------------	-------

Other Ingredients:	51.2%
--------------------	-------

Total:

100.0%

¹CAS No. 736994-63-1

Fortenza is formulated as a flowable suspension.

Fortenza contains 5.0 lb/gal of cyantraniliprole (FS). One fluid ounce of Fortenza contains 17.72 grams of cyantraniliprole.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1420

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Fortenza as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with Fortenza (cyantraniliprole) insecticide seed treatment.
- Do not use treated seed for food, feed or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with Fortenza (cyantraniliprole):

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets, or livestock to have access to treated seed.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Use of Fortenza as a seed treatment must be communicated to all personnel involved in seasonal insect control recommendations.
- This product is highly toxic to bees exposed to direct treatment or residues on blooming crop or weeds. Ensure that the planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops and weeds.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading.
- Dispose of all excess treated seed by burying seed away from bodies of water. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate bodies of water when disposing of excess treated seed or washwaters of planting equipment.
- Excess treated seed may be used for ethanol production only if (a) ethanol production by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.
- Dispose of seed packaging in accordance with local requirements.
- For rice (wet-seeded): Do not apply a total of more than 0.4 lb ai per acre of cyantraniliprole-containing products per calendar year; this is the total cyantraniliprole applied by seed treatment, soil application and foliar application.
- For rice (wet seeded): This seed has been treated with a maximum of 0.03 mg cyantraniliprole per seed. Calculate actual amount of cyantraniliprole applied per acre based on seeds planted per acre.
- Including the Fortenza seed treatment, make no more than two applications of cyantraniliprole or other Group 28 products per generation to the same insect species on a crop or within a 30-day period (count planting date as day 1 if using treated seed). Application(s) to the next generation of target pest(s) must be with an effective product with a different mode of action (non-Group 28 insecticide).

CROP USE DIRECTIONS

RICE (WET SEEDED) – NOT FOR USE IN CALIFORNIA

To provide early-season protection against damage caused by rice water weevil and suppression of grape colaspis on rice (wet seeded).

Fortenza at labeled rates provides protection against the damage caused by rice water weevil during early-season growth and development of rice. Fortenza treated rice seed may be broadcast (aerial) into flooded fields. Fortenza treated rice seed cannot be soaked or pre-germinated before seeding.

Full-season protection or complete reduction of damage from the use of Fortenza is not expected against the listed insects.

Apply Fortenza at a maximum rate of 0.03 milligrams active cyantraniliprole per seed (3.47 fl oz per 100 lb of seed*) (each milliliter of Fortenza contains 0.6 grams of cyantraniliprole). Do not apply more than 0.17 lb active cyantraniliprole per acre as a seed treatment application. Do not apply more than 0.4 lb ai of cyantraniliprole-containing products per acre per calendar year including all types of applications (seed treatment, soil, foliar).

*Based on an average of 21,000 seeds per pound

It is recommended to apply Fortenza with compatible and registered seed treatment fungicides such as Apron XL, Dynasty or Maxim 4FS which are proven to provide early-season protection from seed and seedling diseases, and Cruiser 5FS for additional early-season insect pest protection.

Fortenza® and the Syngenta Logo are trademarks of a Syngenta Group Company

©2020 Syngenta