

FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

MYCLOBUTANIL

GROUP

3

FUNGICIDE

For Distribution and Use Only in the State of Oregon

Rally® 40WSP

EPA Reg. No. 62719-410

EPA SLN No. OR-230005

Blackberry Rust Control in Evergreen Blackberry (Thornless Evergreen)

This label valid until December 31, 2027 or until otherwise amended, withdrawn, canceled, or suspended.

Active Ingredient:

myclobutanil: a-butyl-a-(4-chlorophenyl)-1H-
1,2,4, triazole-1-propanenitrile 40%
Other Ingredients 60%
Total 100%

Keep Out of Reach of Children

CAUTION

Environmental Hazards

For terrestrial uses, do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

GROUNDWATER ADVISORY

Myclobutanil is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This SLN label and the federal label for this product must be in the possession of the user at the time of pesticide application.

Read this SLN and the label affixed to the container for Rally 40WSP before applying.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the container label for Rally 40WSP.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

General Information

Rally 40WSP is a systemic, protectant and curative fungicide. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program.

Blackberry rust (*Phragmidium violaceum*) spores are carried by the wind. When *P. violaceum* infects leaves, purple spots develop on the top of leaves with corresponding yellow and black pustules underneath the leaves. Other plant tissues can become infected, including floral bracts. Some defoliation and die-back of the cane can also occur. As part of a comprehensive IPM program, susceptible clones of Himalayan blackberry plants, which may serve as a source of inoculum, should be controlled.

Application Directions

Application Timing: Initiate applications when fully expanded leaves are present. Consult your Corteva Agriscience representative, extension specialist, or certified crop advisor for any additional local use recommendations for your area.

Application Rate: Apply as a foliar spray at the rate indicated for the target disease.

Disease	Rally 40WSP (oz/acre)	Use Recommendations	Restrictions
blackberry rust (<i>Phragmidium violaceum</i>)	5 (2 oz ai/acre)	Apply at 10- to 18-day intervals. Use the shorter spray interval under heavy disease pressure.	<ul style="list-style-type: none">• Do not apply more than 20 oz of Rally 40WSP (0.5 lb active) per acre per calendar year.• Do not make more than 4 applications per calendar year. One of the four applications may be made after harvest.• For use under this SLN, do not apply Rally 40WSP using irrigation equipment.• Pre-harvest Interval: Do not apply within 5 days of harvest.

Section 24(c) Registrant:
Corteva Agriscience
9330 Zionsville Road
Indianapolis, IN 46268
1-800-992-5994 (option #4)

™®Trademarks of Corteva Agriscience and its affiliated companies

R182-081
Accepted: 02/29/2023
Replaces R182-067