

# NoMate®

## PBW MEC

### PHEROMONE COMMUNICATION DISRUPTANT

NoMate® PBW MEC is a pheromone communication disruptant for control of Pink bollworm (*Pectinophora gossypiella*)

#### Active Ingredients:

(Z,Z)-7,11- Hexadecadien-1-ol acetate.....10.0%  
(Z,E)-7,11- Hexadecadien-1-ol acetate.....10.0%

Inert Ingredients .....80.0%  
TOTAL.....100.0%

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

#### STATEMENT OF PRACTICAL TREATMENT

**If in Eyes:** Flush with plenty of water. Get medical attention if irritation persists.

**If on Skin:** Wash affected areas with soap and water. Get medical attention if irritation persists.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes or clothing. Do not breathe vapors or spray mist.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

EPA EST. No. 36638-MT-01  
EPA REG. No. 36638-12

Net contents: 1 Pint (16 fl. ozs.)

NoMate is a trademark of Ecogen Inc.  
ver. 3/93 • Copyright ©1993 Ecogen Inc.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a safe manner. Store in original container only, in a cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### APPLICATIONS INSTRUCTIONS

This product is intended for use in controlling Pink bollworm damage in cotton by preventing male moths from finding and mating females. Its use results in reduced crop damage by suppression of fertile egg laying and larval infestation. Apply 1/3 to 1 2/3 fluid ounces (2 to 10 grams active ingredient equivalent), per acre. Apply as a tank mix with other pesticides or in 1 to 3 gallons of water per acre. Repeat applications may be necessary depending upon the extent of infestation.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

### WARRANTY AND CONDITIONS OF SALE

Ecogen warrants that this product conforms to the description on this label and is reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under normal conditions of use.

ECOGEN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

If this product is defective, Buyer's exclusive remedy shall be the replacement of the product, or if replacement is impractical, refund of the purchase price. In no case will Ecogen be liable for incidental, consequential or special damages resulting from the handling, storage or use of this product.

**SCENTRY**  
BIOLOGICALS INC.

610 Central Avenue  
Billings, Montana 59102  
(406) 248-5856

## **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE). The requirements in this section only apply to uses of this product that are covered by the Worker Protection Standard. This product does not have a restricted-entry interval.

## **INSTRUCTIONS FOR USE**

NOMATE PBW MEC is intended for use in preventing pink bollworm damage in cotton. The product functions by interfering with mating communication between adult moths. Its use results in reduced crop damage by suppression of fertile egg laying and larval infestation. Apply 1/3 to 1 2/3 fluid ounces (2 to 10 grams active ingredient equivalent), per acre. Apply as a tank mix with other pesticides or in 1 to 3 gallons of water per acre. Repeat applications may be necessary depending upon the extent of infestation.

**DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.**

## **WARRANTY AND CONDITIONS OF SALE**

Scentry Biologicals warrants that this product conforms to the description on this label and is reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under normal conditions of use.

SCENTRY BIOLOGICALS MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

If this product is defective, Buyer's exclusive remedy shall be the replacement of the product, or if replacement is impractical, refund of the purchase price. In no case will Scentry Biologicals be liable for incidental, consequential or special damages resulting from the handling, storage or use of this product.