GROUP 10A ACARICIDE

This supplemental label expires on December 31, 2021 and must not be used or distributed after this date.

Onager Optek Miticide

EMULSIFION FOR AGRICULTURAL USE ONLY

> Onager Optek contains 1.0 lb. active ingredient per gallon Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

PREHARVEST INTERVAL

The required days between the last application and harvest are given in () after each crop name.

APPLICATION DIRECTIONS

APPLICATION DIRECTIONS			
CROP	PEST	RATE	COMMENTS
		Oz./acre	
WHEAT, including emmer wheat and triticale (14 day PHI) (West of the Rockies)	Banks grass mite	10 – 24	Apply at the first sign of mites before populations begin to build. Ground Application: Apply the recommended rate of Onager by ground equipment at 15 – 20 GPA. Air Application: Apply the recommended rate of Onager by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: see details under Chemigation Systems.
	Do not make more than one application per year.		
	Do not apply after the boot stage of growth.		
	Do not harvest hay or forage for at least 14 days after application.		
	Do not apply more than 24 oz formulated product per acre per year.		
GH TOMATOES (1 day	Tetranychid Mites	12 - 24	Use sufficient volume of water to obtain complete coverage but not less than
PHI)	(such as European		20 gallons per acre.
	Red Mite and Two-		
	spotted Spider		
	Mite)	L .	
	Do not make more than one application per year.		
	Do not apply more than 24 ounces per acre per year.		
CITRUS (7 day PHI)	Tetranychid Mites	12 - 24	Application by ground equipment only. Use sufficient volume of water to
(CA, AZ AND TX ONLY)	(such as European		obtain complete coverage but not less than 20 gallons per acre.
	Red Mite and Two-		
	spotted Spider		
	mite)		
	Do not make more than one application per year. Paratase the same than 0.4 approximately appro		
	Do not apply more than 24 ounces per acre per year.		

NET CONTENTS



Produced For: Gowan Company P.O. Box 5569 Yuma, AZ 85366