



Ag Independent Retailers



CNI Agri-Oil

Crop Oil Concentrate

PRINCIPAL FUNCTIONING AGENTS:

Paraffin base petroleum oil	83.0%
Alkylphenol ethoxylate, alcohol ethoxylate	17.0%
TOTAL:	100.0%

Ingredients exempt from tolerance under 40 CFR 180.

Contains Petroleum distillates.

KEEP OUT OF REACH OF CHILDREN

WARNING

Causes eye irritation.
Causes skin irritation.
Harmful if swallowed.



See **FIRST AID** and **Additional Precautionary Statements Below**

Read Entire Label Before Using This Product

Net Content:
2.5 Gals. (9.46 L)

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

WARNING: Causes eye and skin irritation. Harmful if swallowed or if in contact with skin. Do not get on skin, in eyes or on clothing. Do not eat, drink or smoke when using this product. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Take off contaminated clothing and wash before reuse.

Avoid release to the environment.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear eye protection, face protection, protective gloves and protective clothing.

FIRST AID

IF IN EYES:

- Rinse cautiously with water for 15 to 20 minutes.
- Remove contact lenses, if present and easy to do so. Continue rinsing.
- If eye irritation persists, get medical attention.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- If skin irritation occurs, get medical advice.

IF SWALLOWED:

- Call a poison control center or doctor if you feel unwell.
- Rinse mouth.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call CHEMTREC at 1-800-424-9300 for emergency.

ENVIRONMENTAL HAZARDS

Do not contaminate water sources by cleaning of equipment or disposal of washwater.

USE INFORMATION

CNI AGRI-OIL is a superior blend of highly refined paraffinic oil and emulsifier-surfactant system. It is designed for use with a broad range of pesticides including herbicides, insecticides, fungicides and/or fertilizers and micronutrients. This product increases contact activity of the spray, enhances penetration and provides uniform coverage of the leaf surface. This product contains an antifoam to suppress foam development in tank-mix combinations.

DIRECTIONS FOR USE

Read and follow all information on the label of the pesticide, fertilizer or micronutrient with which this product is used.

Always refer to the pesticide and additive label(s) for additional recommendations or precautions. Follow the most restrictive label.

To prepare agricultural sprays containing this product, fill the spray tank half full with water, add the proper amount of pesticide, fertilizer or micronutrient as directed by its label and then add this product. Maintain agitation during the entire process. Continue agitation while filling the tank with water and agitate before and during each application to ensure a uniform spray.

Note: Care should be exercised when sprays containing this product are applied to new crop varieties or highly inbred crop lines not previously treated with this product. It is recommended that a small test area be treated before spraying large areas. Use care during periods when pesticides are more active, such as during periods of hot or humid conditions. Not for aquatic use.

Manufactured For:
CNI

800 Business Park Drive, Hwy 82 W
Leesburg, GA 31763
229-883-7050

Crop	Problem/Situation	Pesticide*	Rate of This Product
Citrus	Annual and Perennial grass weeds	Fusilade®, Gramoxone®, Poast®, Prism®	1% v/v or 1 to 2 pts. per acre
Corn (Field, Seed, Silage, Sweet)	Annual and Perennial grasses, Broadleaf weeds	Accent®, Atrazine 4L or 90 DF, Atra-5™, Banvel®, Basagran®, Basis Gold®, Clarity®, Guardsman®, Resource®	1 to 1.25% v/v or 1 pt. up to 1 qt. per acre
Cotton	Annual and Perennial grasses, Broadleaf weeds	Cobra®, Diuron 4L & 80 Herbicide, Fusilade, Poast, Poast Plus, Select®, Staple®	0.5 to 1% v/v or 1 to 2 pts. per acre
Cotton	Post-emergent grass (Spot or small area treatment)	Fusilade, Poast	1 to 2 pts. per acre
Cotton	Boll Openers	Def® 6, Harvade®	1 to 2 pts. per acre
Cotton	Defoliation	Defol® 5, Defol 6, Defol 750 (sodium chlorate), Def 6, Harvade	1 to 2 pts. per acre
Peanuts	Annual and Perennial grasses, Broadleaf weeds	Basagran, Poast, Pursuit®	1.25% v/v or 1 pt. to 1 qt. per acre
Peanuts	Ascochyta web blotch, Blackhull, Cercospora leaf spot, Rust	Benlate® with Manzate® (Do not use Benlate alone)	8 to 16 fl. ozs. per acre
Pecans	Brown leaf spot, Liverspot, Powdery mildew, Scab	Benlate Fungicide (Do not use Benlate alone)	8 to 16 fl. ozs. per acre
Pome and Stone fruits	Black rot, Fly speck, Fruit rots, Powdery mildew, Scab, Sooty blotch	Benlate Fungicide (Do not use Benlate alone)	8 to 16 fl. ozs. per acre
Potatoes	Annual and Perennial grasses	Gramoxone, Poast	1 to 2 pts. per acre
Potatoes	Vine desiccation	Defol 5, Defol 6, Defol 750	1 to 2 pts. per acre
Rice	Annual and Perennial grass weeds	Basagran, Propanil	1 to 2 pts. per acre
Sorghum or Milo	Annual and Perennial grass weeds	Atrazine 4L or 90 DF, Atra-5	1 to 2 pts. per acre
Soybeans	Annual and Perennial grass weeds	Basagran, Classic®, Cobra, Fusilade, Poast, Pursuit, Resource, Synchrony® STS	0.5 to 1.25% v/v or 1 pt. to 1 qt. per acre
Soybeans	Post-emergent grass (Spot or small area treatment)	Fusilade, Poast	1 to 2 pts. per 25 gals.
Sugar beets	Annual and Perennial grass weeds	Poast, Prism	1 to 1.25% v/v or 1 qt. per acre
Sugarcane	Annual and Perennial grass weeds	Atrazine 4L or 90 DF, Atra-5, Diuron 4L, Diuron 80 Herbicide	8 fl. ozs. to 1 qt. per acre
Ornamental, Nursery and other Non-food crops	Post-emergent grass	Poast	2 pts. per acre

*Follow pesticide label(s) for use rates and instructions.

NON-CROP USE		
Application Site	High-Volume Ground (per 100 gals.)	Low-Volume Ground or Aerial (per acre)
Conifer Release* (Herbaceous and Woody)	—	4 fl. ozs. Herbaceous; 8 fl. ozs. Woody (Based on 10 gallons per acre total volume)
Ditchbank	1 to 4 pts.	2 to 4 pts.
Forestry Site Prep	1 to 4 pts.	2 to 4 pts.
Non-Crop Applications	1 to 4 pts.	2 to 4 pts.
Utility and Roadside R.O.W.	1 to 4 pts.	2 to 4 pts.

*Do not exceed 0.25% by spray volume for Herbaceous conifer release or 0.75% by spray volume for Woody conifer release.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PRODUCT STORAGE: Store in original container in a cool, dry place.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent) during mixing and loading. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state and local ACRC recycler visit the ACRC web page at www.acrecycle.org. Decontaminated containers may also be disposed of in a sanitary landfill.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically directed and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

All other brand names, product names, or trademarks belong to their respective holders.