

GUARANTEED ANALYSIS

Calcium (Ca)...... 6.0%

DERIVED FROM: Calcium Carbonate

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENT

Wash thoroughly after handling this product. Do not ingest. Avoid contact with eyes and skin. Do not inhale spray mist.

FIRST AID

IF ON SKIN: Take off contaminated clothing off and wash before reuse. Wash skin with plenty of soap and water. *IF IN EYES:* Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. *IF SWALLOWED:* Call a poison control center or doctor immediately for treatment advice. Rinse mouth. Do not induce vomiting.

Call a poison control center or doctor for treatment advice if symptoms persists.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm

NOTICE OF WARRANTY - Westbridge warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Westbridge. In no case shall Westbridge be liable for consequential, special or indirect damages resulting from the use or handling of this product. Westbridge makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

MANUFACTURED BY WESTBridge

1260 Avenida Chelsea, Vista, CA 92081 USA (800) 876-2767 • www.westbridge.com

NET CONTENTS/NET WEIGHT:

- □ 2.5 Gallons (9.7 L) / 24 Lbs (10.89 Kg)
- □ 30 Gallon Drum (114 L) / 288 Lbs (132 Kgs)
- ☐ 55 Gallon Drum (209 L) / 528 Lbs (242 Kgs)
- ☐ 275 Gallon Tote (1040 L) / 2640 Lbs (1210 Kgs)

Bulk density 9.6 lbs/gl at 68°F.

PRODUCT INFORMATION

Earthtrend® Cal Plus is a liquid plant food containing calcium, which is essential for membrane permeability and for promoting root and leaf development and overall plant vigor.

Earthtrend® Cal Plus is completely water soluble for improved nutrient availability and uptake by plant roots and leaves, and to provide a fast method of boosting plants under calcium deficiency stress.

Earthtrend® Cal Plus can be applied to all crops including vegetables, fruit trees and vines, tree nuts, flowering and ornamental trees and shrubs, flowers, containerized plants, lawns and turf. Apply as a soil drench or a foliar spray.

Earthtrend® Cal Plus may <u>not</u> be compatible with phosphate fertilizers.

Earthtrend® Cal Plus can also be used to leach sodium from soils and correct salinic irrigation water.

DIRECTIONS FOR USE

SHAKE WELL BEFORE USING. FOR COMMERCIAL USE ONLY. ALWAYS READ ENTIRE LABEL.

Aerial or Foliar Application: 1 to 2 quarts per acre. Apply to plant foliage to point of run-off, throughout growing season as needed.

Soil Application: 1/2 to 2 gallons per acre. Apply in irrigation water until ground is saturated. Continued applications every 7 to 14 days at rate of 1 quart per acre through growing season as needed. Alternatively, Earthtrend Cal Plus can also be applied as a pre-plant, side dress, fertigated, chemigated, top dressed or banded to the soil.

Nursery & Container: 1/2 to 2 ounces per gallon of water; apply to plant foliage to run-off or as a soil drench, continue through the season as needed.

Turf & Lawns: 3 to 6 ounces per 1,000 square feet; apply through irrigation water, continue applications at rate 1 ounce per 1,000 square feet every 7 to 14 days as needed.

Compatibility: Always add Earthtrend® Cal Plus last when mixing with sulfur products. Always conduct a compatibility jar test before mixing Earthtrend® Cal Plus with new products. Earthtrend® Cal Plus may not be compatible with phosphate fertilizers.

NOTICE: Phytotoxicity occasionally occurs on sensitive plants. Perform a phytotoxicity test on a small plot before large-scale application. To lower the risk of phototoxicity, applications should be made early morning or late evening.