

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA**

Carzol[®] SP
INSECTICIDE

**IN WATER SOLUBLE PACKAGING
For Agricultural Use Only**

EPA REG. NO. 10163-265

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

| CROP | PEST | APPLICATION |
|--|--|--|
| Oranges, Lemons, Limes, Tangerines, Tangelos and Grapefruit | Katydid | Apply 0.5 to 1.25 lbs. of CARZOL SP per acre as outside coverage spray when thrips appear. For aerial applications, use a minimum of 10 gallons of water per acre. |
| | <ul style="list-style-type: none">• Do not apply more than 1.25 lbs. CARZOL SP per acre in one crop season.• Do not apply after fruit reach a diameter of one inch.• If unharvested grapefruit and Valencia oranges are present from the previous crop, an application may be made to the new crop. However, a preharvest interval of 30 days must be observed for the unharvested crop. <p>Note: If soft brown scale is present in the grove, a suitable scalicide should be applied after the CARZOL SP application.</p> | |