
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 953A-S9 0716

GROUP	9	12	FUNGICIDES
-------	---	----	------------

Switch® 62.5WG

Fungicide

This supplemental label expires on July 12, 2019 and must not be used or distributed after this date.

Active Ingredients:

Cyprodinil*	37.5%
Fludioxonil**	25.0%
<hr/>	
<i>Other Ingredients:</i>	37.5%
Total:	100.0%

*CAS No. 121552-61-2

**CAS No. 131341-86-1

Switch 62.5WG is a water-dispersible granule containing 37.5% cyprodinil and 25% fludioxonil.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-953

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Switch 62.5WG as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Rotational Crop Restrictions

Do not plant any crop which is not registered for use with cyprodinil or fludioxonil for a period of 30 days, unless a shorter interval is specified on the following list.

Rotational Crop	Planting Time From Last Switch 62.5WG Application
Beans (dried and succulent except cowpeas)* Berries (bushberries 13-07B, caneberries 13-07A)* Brassica (Cole) Leafy Vegetables* Cucurbits* Herbs (fresh and dried)* Leafy Vegetables* Leaves of Root and Tuber Vegetables* Onions (dry bulb, garlic, and green) Peppers Tuberous and Corm Vegetables (crop subgroup 1C)* Root and Tuber Vegetables except Sugar beet* Strawberries Tomatoes Watercress Crops Not Intended for Food or Feed	0 days
All Other Crops Intended for Food or Feed	30 days

*See crop lists in **CROP USE DIRECTIONS** section.

In annual crops where multiple crops can be grown per year (double/triple cropping), do not apply more than 1.3 lb ai cyprodinil and 0.9 lb ai fludioxonil per acre per year to an individual plot of land.

For the crops to which aerial applications are allowed, refer to the specific crop directions for use.

Nassau and Suffolk counties of New York: use limited to strawberries and onions.

Crop	Disease	Product Rate oz/Acre	Remarks
Strawberry and Berry, Low Growing Subgroup 13-07G (except Cranberry)* Strawberry And cultivars and/or hybrids of these.	Gray Mold (<i>Botrytis cinerea</i>) Powdery mildew (<i>Sphaerotheca macularis</i>) Anthracnose (<i>Colletotrichum</i> spp.)	11-14	Begin application at or before bloom and continue on a 7-10 day interval. Resistance Management: After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications.
	Root and crown anthracnose at planting (<i>Colletotrichum</i> spp.)	5-8 oz per 100 gal water	Apply as a preplant dip to strawberry roots and crowns at the rate of 5 to 8 oz per 100 gallons of water for suppression of root and crown rot caused by anthracnose. Wash transplants to remove excess soil prior to dipping. Completely immerse planting stock in dip solution. Dip or expose plants for a minimum of 2 minutes or a maximum of 5 minutes. Completely drain the transplants after dip. DO NOT reuse solution. Dispose of dip solution according to local regulations. Plant treated plants as quickly as possible. For continued anthracnose control, follow with foliar applications of beginning 2-3 weeks after transplant.
*Additional Low Growing Berries: Bearberry; bilberry; cloudberry; muntries; partridgeberry; and cultivars and/or hybrids of these			
Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions 1) Make no more than two applications by air. 2) Make only one pre-plant dip application per crop. 3) Do not apply more than 56 oz/A of Switch 62.5WG per year. 4) Do not apply more than 1.3 lb ai/A of cyprodinil-containing products and 0.9 lb ai/A of fludioxonil-containing products per year. 5) May be applied on the day of harvest (0-day PHI).			

Switch® is a trademark of a Syngenta Group Company
©2016 Syngenta