SUPPLEMENTAL LABELING

Syngenta Crop Protection LLC

P. O. Box 18300 Greensboro, North Carolina 27419-8300 SCP 993B-S4 0316

GROUP 14 HERBICIDE

Ringside® Herbicide

This supplemental label expires on 05/15/2019 and must not be used or distributed after this date.

Active Ingredient/Guarantee:

Sodium salt of fomesafen

5-[2-chloro-4-(trifluoromethyl)phenoxy]-N-(methylsulfonyl)-2-nitrobenzamide......22.8%*

Other Ingredients: 77.2%

Total: 100.0%

Ringside Herbicide is formulated as a soluble liquid.

Ringside Herbicide contains 1,2-benzisothiazolin-3-one at 0.02% as a preservative.

*Ringside Herbicide is equivalent to 21.7% or 2 pounds per U.S. gallon or 240 grams per liter of fomesafen active ingredient.

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. 100-993

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Ringside Herbicide as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

GROUND APPLICATION

Preplant Surface and Preemergence Application - Use a minimum of 10 gallons per acre. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for preplant surface or preemergence applications.

Postemergence Application - Use sufficient spray volume and pressure to ensure complete coverage of the target weed. A spray volume of 10-20 gallons per acre and 30-60 psi at the nozzle tip is recommended. On large weeds and/or dense foliage, use 60 psi and a minimum of 20 gallons per acre to ensure coverage of weed foliage.

The use of flat fan nozzles will result in the most effective postemergence application of Ringside Herbicide. Use nozzles that are set up to deliver medium quality spray (ASAE Standard S-572).

RESTRICTIONS

- A maximum of 1.5 pt of Ringside Herbicide (or a maximum of 0.375 lb ai/A of fomesafen from any product containing fomesafen) may be applied per acre per year in Region 1 (see Regional Use Map).
- A maximum of 1.5 pt of Ringside Herbicide (or a maximum of 0.375 lb ai/A of fomesafen from any product containing fomesafen) may be applied per acre in ALTERNATE years in Region 2 (see Regional Use Map).
- A maximum of 1.25 pt of Ringside Herbicide (or a maximum of 0.313 lb ai/A of fomesafen from any product containing fomesafen) may be applied per acre in ALTERNATE years in Region 3 (see Regional Use Map).
- A maximum of 1 pt of Ringside Herbicide (or a maximum of 0.25 lb ai/A of fomesafen from any product containing fomesafen) may be applied per acre in ALTERNATE years in Region 4 (see Regional Use Map).
- A maximum of 1 pt of Ringside Herbicide (or a maximum of 0.25 lb ai/A of fomesafen from any product containing fomesafen) may be applied per acre in ALTERNATE years in Region 4a (see Regional Use Map). Apply only to soybeans in Region 4a. Do not make a Ringside Herbicide application later than June 20th. Cumulative rainfall plus overhead irrigation must total 15 inches from the period of Ringside Herbicide application to soybean crop maturity to allow planting of rotational crops listed in this label (refer to Rotational Crop Restrictions section). If the soybean crop is lost or the required cumulative rainfall plus irrigation is not received as outlined above, plant only soybeans the following growing season.
- A maximum of 0.75 pt of Ringside Herbicide (or a maximum of 0.1875 lb ai/A of fomesafen from any product containing fomesafen) may be applied per acre in ALTERNATE years in Region 5 (see Regional Use Map).
- Do not make ground or aerial application during temperature inversions.
- Do not apply when wind velocity exceeds 15 mph.
- Do not use on potatoes in Nassau and Suffolk Counties, New York.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying Ringside Herbicide at recommended rates:

Rotational Crops	Planting Time From Last Ringside Herbicide Application
Bean, Dry	J 11
Bean, Snap	
Cotton	0 months
Potato	O months
Soybean	
Soybean, Succulent (edamame)	
Bean, Lima	
Pea, Succulent	4 months
Peanut	4 1110111115
Small Grains such as Wheat, Barley, Rye	
Corn, Field	
Corn, Seed	
Corn, Sweet⁵	
Pepper (transplanted) ¹	
Popcorn ⁴	10 months
Pumpkin ²	
Rice	
Tomato (transplanted) ¹	
Watermelon ²	
Bean, Succulent (other than edamame, snap bean and	
lima bean)	
Cantaloupe ²	
Cucumber ²	
Edible-podded beans and peas not otherwise specified	
in this table	12 months
Eggplant	12 months
Pea, Dry	
Pepper, (direct-seeded)	
Squash ²	
Sweet Potato	
Tomato (direct-seeded)	
Sorghum ³	18 months
All other crops not listed above	18 months

¹ 4 months in Region 1

Restriction: Do not graze rotated small grain crops for harvest forage or straw for livestock.

² 8 months in Region 1

³ 10 months in Region 1


 ^{4 12} months in the states of Ohio, Kentucky, Illinois, Indiana, Iowa, and Regions 4 and
 4 4 when applied at rates of 1 pint per acre or more

^{5 18} months in the states of Connecticut, Maine, Massachusetts, New Hampshire, New York, Rhode Island, Vermont and Region 5


RINGSIDE HERBICIDE - USE RATES AND WEEDS CONTROLLED

REFER TO MAP FOR DEFINITION OF SPECIFIED GEOGRAPHIC REGIONS

RINGSIDE HERBICIDE REGIONAL USE MAP



REGION 1 (Maximum Rate 1.5 pt/A per year)



Inc	ludes the following	states or portion of states where Ringside Herbicide may be applied:
	Alabama	All areas.
	Arkansas	All areas.
	Florida	All areas except for Miami-Dade County.
	Georgia	All areas.
	Louisiana	All areas.
Region 1	Mississippi	All areas.
	Missouri	Counties of Bollinger, Butler, Cape Giradeau, Dunklin, Madison, Mississippi, New Madrid, Pemiscot, Perry, Ripley, Scott, Stoddard and Wayne.
	North Carolina	All areas.
	Oklahoma	All areas east of U.S. Highway 75 and East of Indian Nation Parkway.
	South Carolina	All areas.
	Tennessee	All areas.
	Texas	All areas east of U.S. Highway 77 to State Road 239, including all of Calhoun County.

REGION 2 (Maximum Rate 1.5 pt/A, alternate years)



Includes the following states or portion of states where Ringside Herbicide may be applied:		
	Delaware	All areas.
	Illinois	All areas south of Interstate 70.
	Indiana	All areas south of Interstate 70.
Region 2	Kentucky	All areas.
Region 2	Maryland	All areas.
	Ohio	All areas south of Interstate 70.
	Pennsylvania	All areas south of Interstate 80 to the intersection of U.S. Highway 15 and east of U.S. Highway 15 and U.S. Highway 522.
	Virginia	All areas.
	West Virginia	All areas.

REGION 3 (Maximum Rate 1.25 pt/A, alternate years)



Inc	ludes the following	states or portion of states where Ringside Herbicide may be applied:
	Connecticut	All areas.
	Illinois	All areas north of Interstate 70.
	Indiana	All areas north of Interstate 70.
	Iowa	All areas.
	Maine	All areas.
	Massachusetts	All areas.
Region 3	Missouri	All counties except for those listed in Region 1.
	Ohio	All areas north of Interstate 70.
	New	All areas.
	Hampshire	
	New Jersey	All areas.
	New York	All areas. Do not use on potatoes in Nassau and Suffolk counties, New York.
	Pennsylvania	All areas except those listed in Region 2.
	Rhode Island	All areas.
	Vermont	All areas.
	Wisconsin	All areas south of U.S. Highway 18 between Prairie Du Chien and Madison, and south of Interstate 94 between Madison and Milwaukee.

REGION 4 (Maximum Rate 1 pt/A, alternate years)



Includes the following states or portion of states where Ringside Herbicide may be applied:		
	Kansas	All counties east of or intersected by U.S. Highway 281.
	Michigan	Southern Peninsula.
	Minnesota	All areas south of Interstate 94.
	Nebraska	All counties east of or intersected by U.S. Highway 281.
	North Dakota	All areas east of Interstate 29 from Fargo south to the South Dakota state line.
Region 4	South Dakota	All areas east of Interstate 29 from the North Dakota state line to Watertown, all areas east of Highway 81 from Watertown to Madison and all areas east and south of State Road 34 and U.S. Highway 281 to the Nebraska state line.
	Wisconsin	All areas south of Interstate 94 (except those in Region 3) from Minnesota state line to Eau Claire and south of U.S. Highway 29 from Eau Claire to Green Bay plus Barron, Burnett, Chippewa, Clark, Door, Dunn, Eau Claire, Langlade, Lincoln, Kewaunee, Marathon, Marinette, Menominee, Oconto, Polk, Price, Rusk, Sawyer, Shawano, St. Croix, Taylor, and Washburn counties. The following counties are excluded: Adams, Marquette, Portage, Waupaca, Waushara and Wood.

REGION 4a (Maximum Rate 1 pt/A, Alternate Years*)



	Includes the following portions of states where Ringside Herbicide may be applied:		
Region	Kansas	All areas west of U.S. Highway 281 to the Colorado state line.	
4a	Nebraska	All areas that intersect west of U.S. Highway 281 and east of U.S. Highway 83.	

*Note: Refer to the Use Precautions section for additional requirements that must be followed to use Ringside Herbicide in Region 4a.

REGION 5 (Maximum Rate 0.75 pt/A, alternate years)



Includes the following states or portion of states where Ringside Herbicide may be applied:		
	Minnesota	All areas south of U.S. Highway 2 (except those areas in Region 4), plus
		Betrami, Clearwater, Lake of the Woods, Kittson, Marshall, Pennington,
Region 5		Polk, Red Lake, and Roseau.
	North Dakota	All areas east of U.S. Highway 281, except those areas in Region 4.
	South Dakota	All areas east of U.S. Highway 281, except those areas in Region 4.

Ringside® is a trademark of a Syngenta Group Company.

©2016 Syngenta