

FIFRA Section 24(c) Special Local Need Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Stinger®

EPA Reg. No. 62719-73

EPA Sec 24(c) SLN Registration OR-190008

For Distribution and Use Only in the State of Oregon

This label is valid until December 31, 2024 or until otherwise amended, withdrawn, canceled, or suspended.

For Control of Post Emergent Broadleaf Weeds in Pear Orchards

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Read this SLN and the label affixed to the container for Stinger® herbicide before applying.
- Use of Stinger according to this SLN label is subject to all use directions, restrictions, and precautions imposed by the label affixed to the container for Stinger.

Active Ingredient:

cloprialid: 3,6-dichloro-2- pyridinecarboxylic acid, monoethanolamine salt	40.9%
Other Ingredients	59.1%
Total	100.0%

Acid Equivalent: cloprialid: 3,6-dichloro-2-pyridinecarboxylic acid - 31% (3 lb/gal)

Keep Out of Reach of Children

CAUTION

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

Cloprialid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply cloprialid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Ground Water Advisory

Cloprialid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply cloprialid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching surface water via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of clopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Directions for Use

Use Stinger for postemergence control of broadleaf weeds that are infesting pear orchards. Refer to the product label for Use Precautions, Restrictions, Mixing and Application Instructions, and a complete list of weeds controlled or suppressed.

Target Broadleaf Weeds	Application Rate	Specific Use Restrictions
aster burdock clover, red clover, white curly dock dandelion goldenrod horseweed (marestalk) nightshade, black nightshade, hairy pineappleweed sowthistle, annual thistle, Canada thistle, musk vetch volunteer alfalfa	1/3 to 2/3 pint/acre (0.125 to 0.25 lb ae/acre)	<ul style="list-style-type: none">• Preharvest interval: Do not apply within 30 days of harvest.• Make one to two broadcast applications per crop per year. Do not exceed a total of 2/3 pint per acre.• Apply Stinger to non-bearing trees (well established trees, 1 year or older) and bearing trees.• Avoid direct contact with tree foliage, fruit, or tree trunks.

Application Timing

Apply Stinger to clover and vetch from weed emergence up to the 5-leaf stage of growth. Apply Stinger to nightshade (black and hairy) at the 2-leaf to 4-leaf stage of growth. For control of Canada thistle and annual sowthistle, apply Stinger from rosette up to bud stage.

Application Rate

Apply 1/3 to 2/3 pint of Stinger per acre with ground equipment in 10 gallons or more total spray volume per acre. Use a higher rate in the rate range for heavy infestations or when greater residual control is desired.

Tank Mixtures

Stinger may be tank mixed with other herbicides labeled for use in pears. Follow all applicable use directions, precautions, restrictions, and limitations on the labels for each product used in the tank mix.

Use Restrictions

- Do not apply within 6 to 8 hours of expected rainfall or irrigation.
- Do not apply by aircraft.
- **Chemigation:** Do not apply this product through any type of irrigation system.

24(c) Registrant: Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268

® ™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R043-089
Accepted 04/24/19
Initial Printing