



GENERAL INFORMATION

MAX SET® 99 is designed primarily for foliar applications to prevent or correct nutrient deficiencies in a wide range of agronomic and ornamental plants. Its use is suggested as a supplement to a regular, balanced fertilizer program to enhance yields and improve quality.

Application of MAX SET 99 is a means of obtaining a quick response to needed elements. Foliar applications of MAX SET 99 may be particularly beneficial during periods of peak nutrient demand, for crops grown on soils having poor nutrient availability or to crops suffering from a weakened root system. MAX SET 99 should be used as part of a comprehensive Total Nutrition System® for optimizing plant growth, development, yield and quality.

DIRECTIONS FOR USE

MAX SET 99 may be applied as a foliar spray by air, ground concentrate or dilute. Maintain agitation to provide uniform distribution within spray tank. Use of a high quality CPDA approved spreader, such as R-11®, at recommended rates to improve spray cover is usually advised. Do not apply when plants are under excessive heat or moisture stress. Applications of foliar nutrients to leafy vegetables or soft-skinned fruit during periods of excessive temperatures may cause cosmetic injury. Applications of certain micronutrients to fleshy fruit within 30 days of harvest may result in staining of rind or skin.

FOLIAR APPLICATIONS: MAX SET 99 may be applied by air or with all types of ground spraying equipment. For aerial applications, apply 1-4 pints per acre in a minimum of 6% to 25 gallons of water per acre. For ground applications, apply 1-4 pints per acre. Do not exceed a product concentration of 2% volume/volume maximum. For the following product rates, do not use less than the corresponding gallons of water per acre: 1 pint/acre = 6% gallons of water/acre; 2 pints/acre = 12½ gallons of water/acre; 3 pints/acre = 18% gallons of water/acre.

SOIL APPLICATIONS: MAX SET 99 may be applied to the soil by all types of soil spraying/injection equipment and through certain types of irrigation systems. Refer to the IRRIGATION SYSTEM APPLICATIONS section. Apply 1-4 pints per acre.

VEGETABLE CROPS

TOMATOES, PEPPERS, LETTUCE, SPINACH AND OTHER LEAFY VEGETABLES, BROCCOLI, CAULIFLOWER AND OTHER BRASSICA VARIETIES: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply the first application after transplanting, thinning or at 2nd true leaf stage. Apply 1-2 subsequent applications at 10 to 15-day intervals or as needed to supplement nutritional requirements.

MELONS, CUCUMBERS AND OTHER CUCURBITS: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply the first application after transplanting or at 2nd true leaf stage. Apply 1-2 subsequent applications at 10 to 15-day intervals or as needed to supplement nutritional requirements.

BEANS AND PEAS: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply the first application after thinning or at 2nd true leaf stage. Apply 1-2 subsequent applications at 10 to 15-day intervals or as needed to supplement nutritional requirements.

STRAWBERRIES: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply the first application after transplanting. Apply at 30-day intervals or as needed to supplement nutritional requirements.

POTATOES, ONIONS AND OTHER VEGETABLE ROOT, BULB OR TUBER CROPS: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply the first application after transplanting or at 2^{nd} true leaf stage. Apply 1-2 subsequent applications at 10 to 15-day intervals or as needed to supplement nutritional requirements.

PERMANENT CROPS

ALMONDS, WALNUTS, AND OTHER NUT CROPS: Apply 1-4 pints per acre (+/- 1-4 liters/ hectare) per application. Apply first application at bud break. Apply subsequent applications at 30-day intervals up to nut fill or as needed to supplement nutritional requirements.

PLUMS, NECTARINES AND OTHER STONE FRUIT: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply first application at bud break or pink bud. Apply subsequent applications at fruit set or jacket fall, fruit thinning and 2-3 weeks prior to harvest or as needed to supplement nutritional requirements.

APPLES, PEARS AND OTHER POME FRUIT: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply first application at green tip. Apply subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

99

GUARANTEED ANALYSIS

Soluble Potash (K ₂ O)	9.00%
Calcium (Ca)	

DERIVED FROM: Potassium Phosphite and Calcium Carbonate.

KEEP OUT OF REACH OF CHILDREN



PRECAUTIONARY STATEMENTS

Wash thoroughly after handling. Do not eat, drink or smoke when using this product.

FIRST AID

In all cases, call a poison control center or doctor for further treatment advice. IF SWALLOWED: Call a poison control center or doctor if you feel unwell. Rinse mouth.

For chemical spills, leaks, fire or exposure, call CHEMTREC: (800) 424-9300.

STORAGE AND DISPOSAL

Store in a safe manner. Store in original container only and keep tightly sealed when not in use. Store product and dispose of unused product and empty containers in accordance with federal, state and local regulations.

KIWI: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply first application pre-bloom. Apply subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

CITRUS AND AVOCADOS: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply first application pre-bloom. Apply subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

OLIVES: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply first application 2-3 weeks pre-bloom. Apply subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

GRAPES: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. **Table Grapes:** Apply a single application 2 weeks prior to bloom. Do not apply after bloom. **Wine Grapes:** Apply first application 2 weeks prior to bloom. Apply subsequent applications at bunch preclosing, veraison and 2-3 weeks prior to harvest or as needed to supplement nutritional requirements.

RASPBERRIES, BLACKBERRIES, AND OTHER CANEBERRIES: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply first application pre-bloom. Apply

WEIGHT PER GALLON: 12.7 POUNDS at 68°F

subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

BLUEBERRIES, CRANBERRIES, AND OTHER BUSH BERRIES: Apply 1-4 pints per acre (+/- 1-4 liters/hectare) per application. Apply first application pre-bloom. Apply subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

HOPS: Apply 1-4 pints per acre (+/- 1 to 4 liters/hectare) per application. Apply first application pre-bloom. Apply subsequent applications at 30-day intervals up to harvest or as needed to supplement nutritional requirements.

IRRIGATION SYSTEM APPLICATIONS

Applications of MAX SET 99 to the soil can be made through certain irrigation systems including, but not limited to, low volume systems (e.g., micro sprinklers, drippers, dripp

COMPATIBILITY

MAX SET 99 is compatible with most commonly used foliar pesticides, chemically neutral fertilizers and micronutrients. It is not recommended to use this product with surfactants or adjuvants that lower pH. If the tank mix containing MAX SET 99 is below a pH of 7, DO NOT tank mix with fungicidal copper products OR apply the spray solution to plants that have been or will be treated with a fungicidal copper spray within 10 days of the planned MAX SET 99 application. Always jar test new combinations for compatibility prior to field mixing.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of the product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Buyer and User accept all risks arising from any use of this product, including without limitation uses contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

To the extent permitted by law, neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS logo, Ideas to Grow With, MAX SET, R-11, and Total Nutrition System are registered trademarks of Wilbur-Ellis Company.

K-082015

Manufactured for: **WILBUR-ELLIS COMPANY**7 E. Washington Ave., Yakima, WA 98903
(509) 248-6171

NET CONTENTS: 2.5 GALLONS (9.46 Liters)