



Fungicide For Control and / or Suppression of Listed Diseases in Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Watermelon, Zucchini, (Cucurbit Vegetables Crop Group 9); Eggplant, Ground cherry, Bell and non-bell Pepper, Pepino, Tomato, and Tomatillo (Fruiting Vegetables Crop Group 8-10).

SUPPELMENTAL LABELING

This supplemental label expires on December 16, 2019, and must not be used or distributed after this date.

Read the entire label for **METTLE 125 ME** before proceeding with the use direction contained in this supplemental labeling.

"Label" as used in this supplemental labeling refers to the label booklet for **METTLE 125 ME** and this supplemental.

Active Ingredient:

Tetraconazole* 11.6%

Other Ingredients 88.4%

Total 100.0%

*1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2-tetrafluoroethoxy)propyl]1H-1,2,4-triazole

Contains 1 lb active ingredient (Tetraconazole) per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

If you do not understand this label, find someone to explain it to you in detail.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This supplemental labeling must be in possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- These directions can be found on the currently registered EPA Stamped label.

PRODUCT INFORMATION

METTLE 125 ME is formulated as a one pound active ingredient per gallon micro emulsion (ME). The active ingredient in **METTLE 125 ME** is Tetraconazole, a triazole fungicide (Group 3) that works by inhibiting demethylation and other processes in sterol biosynthesis. Tetraconazole is a systemic, protectant and curative fungicide and is absorbed quickly into the plant tissue. Optimal disease control is achieved when **METTLE 125 ME** is applied in a regularly scheduled spray program.



EPA Registration No. 80289-8
 Manufactured by Isagro S.p.A. for:
 Isagro USA, Inc.
 430 Davis Drive, Suite 240
 Morrisville, NC 27560

Crop	Target Diseases	Product Use Rate per Application fl oz/A (lb ai/A)	Use Directions
Cucurbit Vegetable (Crop Group 9): Balsam apple & pear Casaba Cantaloupe Chayote (fruit) Cucumber Chinese cucumber Chinese okra Chinese waxgourd Cucuzza Edible gourd Gherkin Hechima Honey balls Honeydew Hyotan Melon (Bitter, Chinese preserving, Citron, Crenshaw, Golden Pershaw, Mango, Persian, Pineapple, Snake and Santa Claus) <i>Momordica</i> spp. Muskmelon Pumpkin Squash (Acorn, Butternut, Calabaza, Crookneck, Hubbard, Scallop and Spaghetti) True cantaloupe Vegetable marrow Watermelon Zucchini (Cultivars, varieties, and/or hybrids of these)	Powdery mildew (<i>Sphaerotheca</i> spp. and <i>Erysiphe</i> spp.)	8 (0.0625 lb. ai/A)	Begin applications prior to onset of disease when conditions are favorable for disease development. Make applications on a 7- to 10-day protectant schedule. Make no more than 2 sequential applications of METTLE 125 ME before alternating to another fungicide with a different mode of action. To control other foliar cucurbit diseases, tank mix application of registered fungicides should be made according to label use directions.
	Alternaria leaf blight and leaf spot (<i>Alternaria</i> spp.) Cercospora leaf spot (<i>C. citrullina</i>) Septoria leaf blight (<i>S. cucurbitacearum</i>) phoma blight (<i>P. exigua</i>) Suppression: gummy stem blight (<i>Didymella bryoniae</i>)	8 (0.0625 lb ai/A)	Consult your local university, extension agent, crop consultant or other expert for current recommendations regarding application timing and recommendations for managing gummy stem blight.
Application Instructions: <ul style="list-style-type: none"> Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results. Application may be made by ground, air, or chemigation. Apply in 0.1 to 0.25 inches/A of water for chemigation applications. Chemigation application using excessive water could lead to reduced efficacy. 			
Restrictions and Limitations <ul style="list-style-type: none"> Do not apply more than 24 fluid ounces per acre per year of METTLE 125 ME. Do not apply more than 0.188 lb ai per acre per year of a Tetraconazole containing product. Do not apply more than 5 applications of METTLE 125 ME per acre per year. There must be a retreatment interval of at least 7 days between applications of METTLE 125 ME. Applications may be made up to the day of harvest (PHI = 0 days). 			



EPA Registration No. 80289-8
Manufactured by Isagro S.p.A. for:
Isagro USA, Inc.
430 Davis Drive, Suite 240
Morrisville, NC 27560

Crop	Target Diseases	Product Use Rate per Application fl oz/A (lb ai/A)	Use Directions
Fruiting Vegetable (Crop Group 8-10): African eggplant Bush tomato Cocona Currant tomato Eggplant Garden huckleberry Goji berry Ground cherry Martynia Naranjilla Okra Pea eggplant Pepino Pepper, bell Pepper, non-bell Roselle Scarlet eggplant Sunberry Tomato Tomatillo Tree tomato (Cultivars, varieties, and/or hybrids of these)	Powdery mildew <i>Leveillula</i> spp. <i>Oidium</i> spp.	6 to 8 (0.047 to 0.0625 lb ai/A)	Begin applications prior to onset of disease when conditions are favorable for disease development. Reapply on a 7- to 14-day interval when conditions remain favorable for disease development. Make no more than 2 sequential applications of METTLE 125 ME before alternating to another fungicide with a different mode of action. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate.
	Anthracnose (<i>Colletotrichum</i> spp.) Septoria leaf spot (<i>S. lycopersici</i>) Early blight (<i>Alternaria solani</i>) Gray leaf spot (<i>Stemphylium solani</i> , <i>S. lycopersici</i>) Target spot (<i>Corynespora</i> spp.)	6 to 8 (0.047 to 0.0625 lb ai/A)	

Application Instructions:

- Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results.
- Application may be made by ground, air, or chemigation. Apply in 0.1 to 0.25 inches/A of water for chemigation applications. Chemigation application using excessive water could lead to reduced efficacy.

Restrictions and Limitations

- Do not apply more than 16 fl oz per acre per year of **METTLE 125 ME**.
- Do not apply more than 0.125 lb ai per acre per year of a Tetraconazole containing product.
- Do not apply more than 5 applications of **METTLE 125 ME** per acre per year.
- Do not exceed 21 days between applications.
- There must be a retreatment interval of at least 7 days between applications of **METTLE 125 ME**.
- Applications may be made up to the day of harvest (PHI = 0 days).

ROTATIONAL CROP RESTRICTIONS

Use the time intervals listed below to determine the minimum required time interval between last **METTLE 125 ME** application and new crop planting.

Crop	Replant Interval
Bearberry, bilberry, blueberry (lowbush), cloudberry, corn, gooseberry, grape, kiwifruit (hardy), lingonberry, maypop, muntries, partridgeberry, peanut, pecan, schisandra berry, soybean, strawberry and sugar beet	0 day
All other crops after application to Crop Groups 8-10, 9, and subgroups 13-07F or 13-07G	15 days
Small Grains: (after sugar beet application) Barley, buckwheat, millet, oats, rice, rye, triticale, and wheat	40 days
Sugarcane	45 days
All Other Crops- after application to sugar beet, peanut and pecan	120 days

Read the entire label before using this product, including the "**LIMITATION OF WARRANTY AND LIABILITY**" in the label booklet for **METTLE** before using this product.

METTLE is a registered trademark of Isagro USA, Inc.

EPA stamp (approved 12-15-16)



EPA Registration No. 80289-8
Manufactured by Isagro S.p.A. for:
Isagro USA, Inc.
430 Davis Drive, Suite 240
Morrisville, NC 27560