# Supplemental Label



Walnut Creek, CA 94596-8025

## SELECT® 2 EC HERBICIDE

EPA Reg. No. 59639-3

### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND PAMPHLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

#### FOR POSTEMERGENCE GRASS CONTROL IN:

Crop <sup>(1)</sup> Beet, Garden		Minimum Time From Application to Harvest (PHI) 30 days	Use Rate Per Acre <sup>(2)</sup> 6-8 fl. oz.	Crop Oil Concentrate* Rate Per Acre <sup>(3)</sup> 1% v/v in the finished spray volume.
Canola <sup>(4)</sup>		75 days	4-5 fl. oz.	1% v/v in the finished spray volume.
Celery, Rhubarb		30 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Clover (for use on clover grown in the states of ID, OR and WA only)		15 days	6-16 fl. oz.	1% v/v in the finished spray volume.
Cranberry		30 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Cucurbits, including: Cantaloupes (all) Cucumber Gherkin Honeydew Melon	Muskmelons (all) Pumpkin Squash (all) Watermelon	14 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Flax <sup>(5)</sup>		75 days	4-5 fl. oz.	1% v/v in the finished spray volume.
Fruiting Vegetable, (except Tomato) including: Eggplant	Groundcherry Pepino Peppers (all) Tomatillo	20 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Head and Stem Brassica Vegetables, including:	Broccoli Cabbage Cauliflower Brussels sprouts	30 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Leafy Brassica Greens, (6) including: Broccoli Raab Cabbage, Chinese (Bok Choy)	Collards Kale Mustard Greens Mustard Spinach Rape Greens Turnip Greens	14 days	6-8 fl. oz.	1 % v/v in the finished spray volume.
Leaf Lettuce		14 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Mint <sup>(6)</sup>		21 days	6-16 fl. oz.	1 % v/v in the finished spray volume.
Mustard Seed <sup>(4)</sup>		75 days	4-5 fl. oz.	1% v/v in the finished spray volume

Crop <sup>(1)</sup>	Minimum Time From Application to Harvest (PHI)	Use Rate Per Acre <sup>(2)</sup>	Crop Oil Concentrate* Rate Per Acre <sup>(3)</sup>
Potato, Sweet Potato, Yam and other tuberous and corm vegetables including: Artichoke Cassava Chinese Bitter Jerusalem Sweet Ginger	30 days	6-16 fl. oz.	1% v/v in the finished spray volume for all crops except potatoes.  For potatoes: add crop oil concentrate at the rate of 1 qt./A for ground applications. AMS at the rate of 2.5 - 4.0 lbs./A may be added in potatoes.
Radish	15 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Root Vegetables Ginseng (except Radish), Horseradish including: Chicory Turnip	30 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Spinach <sup>(6)</sup>	14 days	6-8 fl. oz.	1 % v/v in the finished spray volume.
Strawberry	4 days	6-8 fl. oz.	1% v/v in the finished spray volume.
Sugar Beet	40 days	6-16 fl. oz.	1 qt./A for ground application. AMS at the rate of 2.5 - 4.0 lbs./A may be added.
Sunflower	70 days	6-16 fl. oz.	1 qt./A for ground application. AMS at the rate of 2.5 - 4.0 lbs./A may be added.

<sup>\*</sup> Acceptable crop oil concentrates would be those that contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

- (1) SELECT 2 EC Herbicide is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided
- (2) Do not apply more than 8 fl. oz./A in a single application, except for crops noted above that allow for rates up to 16 fl. oz./A. For repeat applications make on a minimum of a 14-day interval. For clover and radish do not apply more than 16 fl. oz. (0.25 lb. ai) per acre in a season. (3) Do not add nitrogen fertilizer source spray adjuvants to the spray mixture, unless specifically advised otherwise.
- (4) Do not apply SELECT 2 EC Herbicide to canola or mustard seed after the crop has begun bolting. Crop injury may occur when applied during the bloom period. Do not exceed 5 fl. oz. (0.08 lb. ai) per acre in a season.
- Do not apply SELECT 2 EC Herbicide to flax during the bloom period. Crop injury may occur when applied during the bloom period. Do not exceed 5 fl. oz. (0.08 lb. ai) per acre in a season.
- Registration pending in the state of New York 8/02.

#### **RESTRICTIONS AND LIMITATIONS**

Do not apply more than 32 fl. oz. of SELECT 2 EC Herbicide (0.5 lb. ai) per acre per season.

Application on Long Island, New York is restricted to no more than 16 fl. oz. of SELECT 2 EC Herbicide (0.25 lb. ai) per acre per

#### RECOMMENDATIONS FOR ANNUAL OR PERENNIAL GRASSES

- Apply only to actively growing grasses at recommended weed heights. Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
  Use the high rate under heavy grass weed pressure and/or when grasses are at maximum height.
- For the control of annual grasses in crops other than canola, flax and mustard seed use 6 to 8 fl. oz./A, for canola, flax and mustard seed use 4 to 5 fl. oz./A. Do not exceed 5 fl. oz./A in canola, flax or mustard seed per acre per application. For the control of perennial grasses use 8 fl. oz./A, except for those crops listed above that allow for rates
- up to 16 fl. oz./A..
- Please refer to container label for specific use rates, weeds and weed sizes.

#### THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Copyright © 2002 by Valent U.S.A. Corporation

Manufactured for Valent U.S.A. Corporation 1333 N. California Blvd., Suite 600 Walnut Creek, CA 94596-8025 www.valent.com Form 2002-SEL-0011 Made in U.S.A.