

TECHNICAL INFORMATION BULLETIN

Restricted Use Pesticide. Due to toxicity to aquatic invertebrate animals. For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

GROUP 15 INSECTICIDE



EPA Reg. No. 400-461

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency

FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY.

The addition of Redneck Peanut Worm control in PEANUTS.

This FIFRA Section 2(ee) Recommendation expires December 31, 2018 and must not be used or distributed after this date.

Insect Growth Regulator

Not for Homeowner/Residential Use

DIRECTIONS FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions for use, restrictions, worker protection standard requirements, and precautions on the EPA registered main label attached to the container of **DIMILIN® 2L Insecticide (EPA Reg. No. 400-461)**.

103116V071TX2ee

Crops	Pests	Application Rate (fl oz/acre)	Application Timing
PEANUTS	PEANUT RESTRICTIONS: Do not make more than 3 applications per season. Do not exceed 24 fl oz (6 oz ai) per acre per year. Pre-harvest Interval: Do not harvest within 28 days of application.		
	Velvet bean caterpillar Mexican bean beetle Green cloverworm	2 – 4	Make applications when larvae are small (< 0.5 inches) to give greater control and minimum insect damage to leaves. Repeat application if damaging numbers reappear. The minimum reapplication interval is 14 days. Use the higher listed rate in the range if the crop has a history of heavy infestations, dense foliage is present, or greater residual control is desired.
	Redneck peanut worm Armyworms, such as: <ul style="list-style-type: none"> - Beet armyworm - Fall armyworm - Southern armyworm - Yellow-striped armyworm Lesser cornstalk borer Soybean looper (suppression)	4 – 8	
	Grasshopper	2	For best results, apply when the majority of infesting grasshoppers have reached the 2nd to 3 rd nymphal stage of development. DIMILIN 2L is not effective in controlling grasshoppers once they reach the adult stage. If a large influx from neighboring fields should occur, the time to reduce that population may not be short enough to minimize extensive foliage feeding. Use a tank mix with a knockdown insecticide under these conditions.
	Aerial Application: Apply in sufficient water (3 to 5 gallons per acre) to achieve uniform coverage of foliage. Ground Application: Apply in 9 to 35 gallons of water per acre to give uniform coverage. Adjuvant Usage: See Cotton section. Since DIMILIN 2L is an insect growth regulator, larvae/nymphs must ingest treated plant material and then molt before populations are reduced. Thus initial signs of control may not be seen until 5 to 7 days after treatment.		

Manufactured For:
MacDermid Agricultural Solutions, Inc.
245 Freight Street
Waterbury, CT 06702-1818

®DIMILIN is a Registered Trademark of MacDermid Agricultural Solutions, Inc.

©Copyright 2016, MacDermid Agricultural Solutions, Inc.