

Supplemental Labeling

Tenkoz Inc.

1725 Windward Concourse

Alpharetta, GA 30005 USA

Staunch™ Herbicide

EPA Reg. No. 62719-570-55467

Increased Use Rates For Longer Residual Control

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Staunch™ herbicide before applying. Follow all applicable use directions and precautions on the EPA-registered label.
- Use of Staunch according to this supplemental labeling is subject to all restrictions and limitations imposed by the label affixed to the product container.

Directions for Use

Staunch™ herbicide may be used in preplant, preemergence, or early postemergence applications in corn. It is designed to provide early season control of grass and broadleaf weeds to allow for optimal timing of the in-crop postemergence application of glyphosate or glufosinate.

Staunch may be used at rates from 1.5 to 3 pints per acre. Use rates in the higher end of the rate range for longer residual activity.

See the product label for Directions for Use, General Use Precautions, Mixing and Application instructions.

Use Rates* for Staunch by Soil Texture and Organic Matter Content

Soil Texture	Soil Organic Matter Content	
	Less Than 3%	3% or Greater
	Pints/Acre	Pints/Acre
Coarse	1.5 – 2.0	1.5 – 2.0
Medium	1.5 – 2.5	1.75 – 3.0
Fine	2.0 – 3.0	2.0 – 3.0

* Use rates in the higher end of the rate range for longer residual activity.

The following table provides a reference to determine the amount of Staunch to be mixed per ton of dry bulk fertilizer for a range of herbicide rates.

Fertilizer Rate (lb/acre)	Acres Covered (per ton)	Staunch (pints/acre)					
		1.5	1.75	2	2.5	2.75	3
		Pints Herbicide/Ton Fertilizer					
200	10	15	17.5	20	25	27.5	30
300	6.7	10	11.7	13.4	16.8	18.4	20.1
400	5	7.5	8.8	10	12.5	13.8	15
500	4	6	7	8	10	11	12
600	3.3	5	5.8	6.6	8.3	9.1	9.9
700	2.9	4.4	5.1	5.8	7.3	8	8.7

Application Precautions

- Uneven application or uneven incorporation of Staunch can result in erratic weed control or crop injury. Over application may result in crop injury or rotational crop damage from soil residue.
- **Maximum Application:** The total cumulative maximum application amount of Staunch on corn is 3.5 pints per acre per crop season.
- Do not exceed 3 pints per acre in a single application.

TMTrademark of Tenkoz Inc.
EPA Accepted: 01/28/09
Initial printing.