

Supplemental Label



Acephate 90 WSP Insecticide EPA Reg. No. 34704-862 (Except in CA)

FOLIAR APPLICATION OF ACEPHATE 90 WSP INSECTICIDE FOR INSECT CONTROL IN SOYBEANS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

GENERAL USE PRECAUTIONS

Do not graze or cut vines for hay or forage. Do not apply more than 1.5 lbs. ai/A of Acephate 90 WSP Insecticide per season. Do not apply Acephate 90 WSP Insecticide within 14 days of harvest. Always read and follow all label directions, restrictions and precautions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

APPLICATION METHOD	PESTS CONTROLLED	RATES OF PROD/ ACRE	REMARKS	DAYS TO HARVEST
FOLIAR: By Air: 5 to 10 gal/A of spray By Ground: 10 to 50 gals/A of spray	Grasshopper Thrips	0.28 to 0.56 lb.	Application Intervals for all agricultural crops: 3 days for applications rates up to 0.5 lb ai/A 7 days for application rates greater than 0.5 lb ai/A.	14
	Potato Leafhopper Stinkbugs	0.56 to 1.1 lbs.		
	Armyworms (except Beet) Bean Leaf Beetle Cabbage Looper Green Clovenworm Mexican Bean Beetle Soybean Aphid Threecornered Alfalfa Hopper Velvetbean Caterpillar	0.83 to 1.1 lbs.		

Loveland Products Inc.
P.O. Box 1286
Greeley, CO 80632-1286