

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Duramax™

EPA Reg. No. 62719-556

### For Control of Annual and Perennial Weeds in Sunflower and Safflower

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

**Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.**

Refer to the label booklet for this product for general information on product use, mixing, application equipment and techniques, labeled uses, and weed rate tables.

**Types of Applications:** Chemical fallow, preplant fallow beds, preplant, preemergence, at-planting, hooded sprayers in row-middles, shielded sprayers in row-middles, wiper applications in row-middles, pre-harvest, and post-harvest treatments.

**Note:** Unless otherwise specified, weed control applications may be made according to the rates listed in the "Annual Weeds" and "Perennial Weeds Rate Table" section in the Duramax™ herbicide label.

#### **Chemical fallow, preplant fallow beds, preplant, preemergence, at-planting**

**Specific Use Instructions:** This product may be applied before, during or after planting sunflower or safflower crops. Broadcast applications must be made prior to emergence.

**Precautions and Restrictions:** In safflower, the total combined application rate of all preemergence and hooded/shielded sprayer treatments must not exceed 2.25 quarts of this product per acre per year. In sunflower, the total combined application rate of all preemergence and hooded/shielded sprayer treatments must not exceed 24 fluid ounces of this product per acre per year.

#### **Hooded or Shielded Sprayers in Row Middles, Wiper Applications in Row Middles**

**Specific Use Instructions:** Wiper applicators or hooded/shielded sprayers may be used between the rows once the crop is established. See the "Selective Equipment" part of the "Application Equipment and Techniques" section of the Duramax label for information on proper use and calibration of this equipment.

**Precautions and Restrictions:** In safflower, the total combined application rate of all preemergence and hooded/shielded sprayer treatments must not exceed 2.25 quarts of this product per acre per year. In sunflower, the total combined application rate of all preemergence and hooded/shielded sprayer treatments must not exceed 24 fluid ounces of this product per acre per year.

### **Preharvest**

**Specific Use Instructions:** This product provides weed control when applied as a harvest aid to a physiologically mature crop prior to harvest of safflower or sunflower. For safflower, apply when the seed has lost its opaque character, approximately 20 to 30 days after the end of flowering of the secondary branches. For sunflower, apply when the backsides of sunflower heads are yellow and bracts are turning brown and seed moisture content is less than 35%.

**Precautions and Restrictions:** Allow a minimum of 7 days between treatment and harvest or livestock feeding. Apply no more than 2.25 quarts of this product at a preharvest timing to safflower. Apply no more than 24 fluid ounces of this product at a preharvest timing to sunflower. Applications must be made at least 30 days prior to planting any crop not listed on the Duramax label.

### **Post-harvest**

**Specific Use Instructions:** This product may be applied after harvest of safflower or sunflower. Higher rates may be required for control of larger weeds, which were growing in the crops at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

**Precautions and Restrictions:** Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation. Applications must be made at least 30 days prior to planting any crop not listed on the Duramax label.