

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Treflan* TR-10*

EPA Reg. No. 62719-131

24(c) Special Local Need Registration EPA SLN No. UT-870002
(For Distribution and Use Only in the State of Utah)

Weed Control in Established Alfalfa

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan TR-10 before applying. Carefully follow precautionary statements and applicable use directions.
- Use of Treflan TR-10 according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan TR-10.

Directions for Use

Apply Treflan* TR-10* herbicide to established alfalfa prior to weed emergence with properly calibrated ground or aerial equipment. A single rainfall or overhead sprinkler irrigation of 0.5 inches or more, flood irrigation or furrow irrigation after application is required to activate Treflan TR-10. If activated using furrow irrigation, the surface of beds between furrows should be thoroughly wetted. If activation does not occur within 10 days after application, Treflan TR-10 should be activated using incorporation equipment that will insure thorough soil mixing with minimum damage to the established alfalfa.

Weeds Controlled in Established Alfalfa

Barnyardgrass	Cheat [†]	Junglerice
Bromegrass [†]	(chess)	Sandbur
(cheatgrass)	Crabgrass	Wild barley
(downy brome)	Woolly cupgrass	Dodder [†]
Canarygrass	Foxtail	(suppression only)

[†] See specific instructions below for control of these weeds.

General Weed Control

Apply Treflan TR-10 at a rate of 20 pounds per acre for all soil textures. Treflan TR-10 may be applied during dormancy or throughout the growing season immediately after a cutting. Because Treflan does not control established weeds, application must be made prior to the expected time of weed germination.

Extended Weed Control and Suppression of Dodder: For extended control of the weeds listed above and suppression of dodder, a second treatment of 20 pounds per acre may be applied (immediately after a cutting) after at least two cutting cycles or 60 days of active alfalfa growth.

Control of Bromegrass and Cheat (Fall Application Only): Apply immediately after a cutting between August 1 and October 1. When fall applied, Treflan controls bromegrass and cheat in addition to other weeds listed above that germinate after application. Bromegrass and cheat begin to germinate in the fall with the onset of cooler weather.

Precautions: Do not make more than two 20 pound per acre applications of Treflan TR-10 to alfalfa within a 12-month period. In the season following application of 20 or more pounds per acre of Treflan

TR-10, where established alfalfa is to be rotated to another crop, plant only those crops for which Treflan can be applied as a preplant treatment or crop injury will result.

Preharvest Interval: Do not graze or harvest hay from treated areas within 21 days after application.

*Trademark of Dow AgroSciences LLC

D06-048-009

Accepted: 01/28/2002

Replaces 122-33-068N.