



SUPPLEMENTAL LABEL

Nufarm Diquat 2L Herbicide

EPA Reg. No. 228-675

LABEL REVISIONS: CANOLA CROP- PREHARVEST DESICCATION BROADCAST

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR THIS PRODUCT BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. "Label" as used in this supplemental labeling refers to the label booklet for this product and this supplemental.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.

This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS IN THE PRODUCT LABEL BOOKLET MUST BE FOLLOWED, INCLUDING STATEMENTS PERTAINING TO THE WORKER PROTECTION STANDARDS, ON THE EPA REGISTERED LABEL.

AGRICULTURAL USES

Apply this product to desiccate canola as a preharvest aid to facilitate harvesting. This product can also be applied as a general herbicide to control weeds in nonbearing crops and noncrop areas.

This product acts on contact with actively growing green plant tissue and for effective control, complete coverage of all green plant tissue is required. Signs of herbicidal activity are usually apparent within a few days of application. Improper application technique and/or application to large, stressed, or mowed weeds will generally result in unacceptable control. Weeds that emerge after application of this product will not be controlled or suppressed. Weeds that are established or larger than 6" may require retreatment.

NOTE: Cool (below 55°F) or cloudy, overcast weather will slow this product activity but will not affect performance.

USE PRECAUTIONS

- Do NOT allow spray to contact or drift to desirable vegetation or severe plant injury or death will result.
- Do NOT use dirty or muddy water when diluting this product.
- Do NOT apply this product through any type of irrigation system.
- Rain or irrigation occurring within 30 minutes of application negatively impacts herbicidal activity.
- Be sure to rinse all spray equipment thoroughly with water after use.

As with any desiccant consider the following before use:

- Premature application will result in the harvest of green seed
- Pod drop will increase during heavy winds following application
- Effects may be more pronounced with some varieties

The user of this product accepts the responsibility for:

- Knowing the sensitivity of the variety treated,
- Properly timing the application, and
- Accepting the risk of unfavorable weather conditions after application.

APPLICATION INSTRUCTIONS

Apply this product to newly emerged weeds while actively growing and before they become too large (weeds 1 - 6 inches in height are easiest to control). Be sure to follow the specified rates listed in the specific crop instructions below, using the higher rates when weeds are large or dense or when applying for harvest aid and the crop vegetation is dense.

If weeds have been mowed or grazed (removing much of the green foliage) be sure to allow weeds to regrow to a height of 2 - 4 inches before application. For harvest aid applications, refer to the specific crop instructions below for application timing specifications.

NOTE: Because dust can coat target surfaces and reduce product activity, avoid applying in extremely dusty conditions (e.g., dust caused by high winds or the passage of equipment tires).

Spray Equipment

Be sure to follow the specific instructions for minimum spray volumes listed in the crop specific instructions below. Note that the specifications are minimum volumes only, and increase spray volumes as necessary to obtain complete coverage of the target weed or plant (without causing foliage runoff). For best results use flat fan nozzles. Other nozzles provide less than complete coverage resulting in reduced performance.

NOTE: When spraying less than 20 gals. of spray carrier per acre, target weeds must not exceed 6 inches in height.

Adjuvants

When applying this product, be sure to always add one of the following adjuvants:

Nonionic Surfactant (NIS) - Add a NIS containing 75% or greater surface active agent at 0.06-0.5% v/v (1/2 – 4 pts. per 100 gals.) of the finished spray volume.

Other Adjuvants - Use adjuvants other than NIS but that meet the following criteria:

- Contains only EPA exempt ingredients.
- Is compatible when mixed with this product (compatibility can be determined using a jar test).
- Is supported locally for use with this product through proven field trials and through university and extension recommendations.

SPECIFIC USE INSTRUCTIONS

| Crop | Use Pattern | Product Rate per Acre | Minimum Total Spray Volume per Acre | Preharvest Interval (Days) | Precautions, Restrictions, and Comments |
|--------|----------------------------------|-----------------------|--------------------------------------|----------------------------|--|
| Canola | Preharvest desiccation broadcast | 1-1/2 to 2 pints | Ground: 15 gallons Air: 5 gallons | 7 | A maximum of one application per crop is allowed. For best results, harvest no later than 10 days after application. |

NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

**Manufactured For Nufarm Americas Inc.
150 Harvester Drive | Burr Ridge, IL 60527**