

PULL HERE TO OPEN ►

RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

GROUP 3 INSECTICIDE



Insecticide

ACTIVE INGREDIENT:

Lambda-cyhalothrin^{1,2} 22.8%

OTHER INGREDIENTS: 77.2%

TOTAL: 100.0%

Lambda T-2 contains 2.08 lb of active ingredient per gal and is a capsule suspension.

¹CAS No. 91465-08-6

²Synthetic pyrethroid

Contains petroleum distillate.

KEEP OUT OF REACH OF CHILDREN. WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1295-5905

EPA Est. 39578-TX-1

SCPSR-HEL-1295D-L1 1114

4062274

AD 111314

NET CONTENTS: 1 gallon

MANUFACTURED FOR

HELENA CHEMICAL COMPANY

225 SCHILLING BOULEVARD, SUITE 300 • COLLIERVILLE, TN 38017

7849-12628 01-15-16 SCP 01-02 (02-09-16)
Lambda T-2 (Helena) Sub – 1 gal bkl/base – Mech
SCP 7849...

AGI: 107327
Print Size – Bkl: 4.75" wide X 5.5" high
Base: 5.5" wide X 5.5" high
Pantone Colors: Pantone Black, Pantone 7687,
Pantone Cool Gray 6, Pantone Cool Gray 10
No. of pages: 36

NOTES

| FIRST AID | |
|--|--|
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| NOTE TO PHYSICIAN | |
| Contains petroleum distillate – vomiting may cause aspiration pneumonia | |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. | |
| HOT LINE NUMBER | |
| For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call CHEMTREC at 1-800-424-9300 | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hrs. after exposure and may last 2 – 30 hrs., without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, Category G, such as barrier laminate or Viton® ≥ 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

continued...

PRECAUTIONARY STATEMENTS (continued)

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife.

For terrestrial uses: do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

CONDITIONS OF SALE – LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale – Limited Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man, or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use, subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used.

To the extent allowed by law, the Company shall not be liable, and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product, and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability, and remedies.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, Category G, such as barrier laminate or Viton® ≥ 14 mils
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or aerial equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gal per acre by air or 10 gal per acre by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, Lambda T-2 may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control.

RESISTANCE MANAGEMENT

Lamba T-2 is a Group 3 Insecticide (contains the active ingredient lambda-cyhalothrin). Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

SPRAY DRIFT PRECAUTIONS

BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing Lambda T-2 onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: *Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21 pp.* www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf

In the State of New York, a 25 ft vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial or ground application. For aerial applications, the 25 ft vegetated non-cropped buffer strip for runoff protection would be part of the larger 150 ft buffer strip (or 450 ft buffer strip for ULV application) required for spray drift.

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airstblast)

Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

Buffer Zone for ULV Aerial Application

Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

Buffer Zone for Non-ULV Aerial Application

Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

SPRAY DRIFT REQUIREMENTS

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition.

Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make aerial or ground applications into temperature inversions.

Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airstblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downward. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

TANK MIX APPLICATION

When tank-mixing with any other agricultural products, **always add Lamba T-2 last**. Fill the tank with $1/2 - 2/3$ volume of the mixing diluent. Make sure all other products are fully dispersed in the mixing diluent before adding the recommended rate of Lamba T-2 to the tank. Add the remainder of the mixing diluent volume. It is recommended that mixing and spray equipment have continuous agitation for best results. Follow the precautions and limitations of the most restricted product in the tank mixture.

While Lamba T-2 has good flexibility for tank mixing with other agricultural products, a jar test for physical compatibility is recommended for untried mixtures, using proper ratios and mixing sequences of all ingredients to be included in the mixture.

Lamba T-2 is an aqueous-based formulation. It is recommended that no type of non-emulsifiable oils be used in combination with Lamba T-2. If adjuvants are used, use only:

- Nonionic Surfactant (NIS) containing at least 75% surface agent, or
- Nonphytotoxic Crop Oil Concentrate (COC), including once-refined Vegetable Oil Concentrate (VOC), or,
- Methylated Sunflower Oils (MSO) containing a minimum of 17% emulsifier.

Adjuvants other than NIS or COC may be used providing the product meets the following criteria:

1. Contains only EPA exempt ingredients.
2. Is nonphytotoxic to the target crop.
3. Is compatible in mixture. (May be established through a jar test.)
4. Is supported locally for use with Lamba T-2 on the target crop through proven field trials and through university and extension recommendations.

In addition, the following may be used as diluents:

- Crop Oil Concentrate
- Methylated Sunflower Oils
- Urea-Ammonium Nitrate

It is recommended that the following not be used in combination with Lamba T-2 as diluents or adjuvants:

- Nonemulsifiable oils
- Diesel Fuel
- Straight Mineral Oil

CHEMIGATION

Sprinkler Irrigation Application

Apply Lamba T-2 at rates and timing described elsewhere in this label. As local recommendations differ, consult your local State Extension Service or other local experts for recommendations on adjuvant or diluent types, (see **TANK MIX APPLICATION**) rates and mixing instructions. These recommendations should be proven, through university and extension field trials, to be effective with Lamba T-2 applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period.

Apply by injecting the recommended rate of Lamba T-2 into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1-0.2 acre-inch of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of Lamba T-2 for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

It is not recommended that Lamba T-2 be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Use Precautions - Sprinkler Irrigation Applications

- A. Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- C. If you have any questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and are capable of being fitted with a system interlock.
- L. Any alternatives to the above required safety devices must conform to the list of EPA-approved alternative devices.
- M. Do not apply when wind speed favors drift beyond the area intended for treatment or non-uniform distribution of treated water.
- N. Do not apply through chemigation systems connected to public water systems.

SPECIFIC USE DIRECTIONS

AGRICULTURAL USES

| Crop | Target Pests | Rate | |
|---|---|---------------|-------------|
| | | lb ai/A | fl oz/A |
| ALFALFA AND ALFALFA GROWN FOR SEED | | | |
| | Alfalfa Caterpillar Army Cutworm Cutworm species Green Cloverworm Leafhopper species Looper species Threecornered Alfalfa Hopper Velvetbean Caterpillar Webworm species | 0.015 – 0.025 | 0.96 – 1.60 |
| | Alfalfa Seed Chalcid (Adult) Alfalfa Weevil Armyworm | 0.02 – 0.03 | 1.28 – 1.92 |

| Crop | Target Pests | Rate | |
|---|---|-------------|-------------|
| | | lb ai/A | fl oz/A |
| ALFALFA AND ALFALFA GROWN FOR SEED (continued) | | | |
| | Bean Leaf Beetle (Adult) Blister Beetle species Blue Alfalfa Aphid Clover Leaf Weevil species Clover Root Borer (Adult) Clover Root Curculio species (Adult) Clover Stem Borer (Adult) Corn Earworm Cowpea Aphid Cowpea Curculio (Adult) Cowpea Weevil (Adult) Cucumber Beetle species (Adult) Egyptian Alfalfa Weevil Fall Armyworm ¹ Grape Colaspis (Adult) Grasshopper species Green June Beetle (Adult) Green Peach Aphid ³ Japanese Beetle (Adult) Meadow Spittlebug Mexican Bean Beetle Pea Aphid Pea Weevil (Adult) Plant Bug species including Lygus species ³ Spotted Alfalfa Aphid Stink Bug species Sweet Clover Weevil (Adult) Thrips species ⁴ Western Yellowstriped Armyworm Whitefringed Beetle species (Adult) Yellowstriped Armyworm | 0.02 – 0.03 | 1.28 – 1.92 |
| | Beet Armyworm ^{1,3} Blotch Leafminer ³ Spider Mites ² | 0.03 | 1.92 |

Remarks

- Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gal per acre by air or 10 gal per acre by ground. When foliage is dense and/or pest populations are high 5–10 gal per acre by air or 20 gal per acre by ground and higher use rates are recommended. Use higher rates for increased residual control.
- Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2–3 days following application. Avoid direct application to bee shelters.
- **Do not** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per cutting.
- **Do not** apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per season.
- **Do not** apply within 1 day of harvest for forage or within 7 days of harvest for hay.

continued...

- 1 Use higher rates for large larvae.
 2 Suppression only
 3 See **Resistance** statement under **General Directions for Use**.
 4 Does not include Western Flower Thrips.

| Crop | Target Pests | Rate | |
|---------------|---|--------------|-------------|
| | | lb ai/A | fl oz/A |
| CANOLA | | | |
| | Armyworm species Cabbage Seedpod Weevil Cutworm species Diamondback Moth Flea Beetle Grasshoppers Looper species Lygus Bug | 0.015 – 0.03 | 0.96 – 1.92 |
| | Cabbage Aphid | 0.03 | 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply a minimum of 2 gal of water per acre.
- **Do not** apply within 7 days of harvest.
- **Do not** apply more than 0.09 lb ai (5.76 fl oz or 0.36 pt of product) per acre per year.

| Crop | Target Pests | Rate | |
|--|--|--|---|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS | | | |
| Corn (at Plant): Field Corn Popcorn Seed Corn Sweet Corn | Corn Rootworm Larvae: Mexican Northern Southern Western Cutworm species Lesser Cornstalk Borer Red Imported Fire Ant ¹ Seedcorn Beetle Seedcorn Maggot White Grub species Wireworm species | 0.005 lb ai per 1000 ft of row ² | 0.33 fl oz per 1000 ft of row ² |

Remarks

- **Banded Applications** – Apply at planting as a 5–7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel.
- **In-Furrow Applications** – Apply into the seed furrow through spray nozzles or microtubes, behind the planter furrow openers and in front of the press wheel.
- Apply a minimum of 3 gal finished spray per acre.

- **Do not** harvest or graze livestock or cut treated crops for feed within 21 days of at plant application.
- **Do not** apply more than 0.09 lb ai (5.76 fl oz or 0.36 pt of product) per acre per crop at plant.
- For field corn, popcorn, and seed corn **do not** apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per crop from at plant and foliar applications. For sweet corn **do not** apply more than 0.48 lb ai (30.72 fl oz or 1.92 pt of product) per acre per crop from at plant and foliar applications.

¹ Suppression only

²Lb ai and fl oz/A of Lamba T-2 Applied at 0.33 fl oz/1000 ft of Row for
Various Row Spacings

| Row Spacing | 40" | 38" | 36" | 34" | 32" | 30" |
|-------------|--------|--------|--------|--------|--------|--------|
| Linear Ft/A | 13,068 | 13,756 | 14,520 | 15,374 | 16,335 | 17,424 |
| Lb ai/A | 0.067 | 0.07 | 0.075 | 0.079 | 0.084 | 0.09 |
| Fl oz/A | 4.3 | 4.55 | 4.8 | 5.05 | 5.4 | 5.75 |

| Crop | Target Pests | Rate | |
|---|--|---------------|-------------|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS | | | |
| Corn (Foliar) Field Corn Popcorn Seed Corn | Corn Earworm ¹ Cutworm species Green Cloverworm Meadow Spittlebug Western Bean Cutworm ¹ | 0.015 – 0.025 | 0.96 – 1.60 |
| | Armyworm ² Bean Leaf Beetle Bird Cherry-Oat Aphid ³ Cereal Leaf Beetle Corn Leaf Aphid ³ Corn Rootworm Beetle (Adult): Mexican Northern Southern Western English Grain Aphid ³ European Corn Borer ¹ Fall Armyworm ² Flea Beetle species Grasshopper species Hop Vine Borer ¹ Japanese Beetle (Adult) Lesser Cornstalk Borer Sap Beetle (Adult) Seedcorn Beetle Southwestern Corn Borer ¹ Stalk Borer ¹ Stink Bug species Tobacco Budworm ^{1,4} Webworm species Yellowstriped Armyworm ² | 0.02 – 0.03 | 1.28 – 1.92 |

| Crop | Target Pests | Rate | |
|---|---|---------|---------|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS (continued) | | | |
| Corn (Foliar) Field Corn Popcorn Seed Corn | Beet Armyworm ⁴ Chinch Bug Greenbug ^{3,4} Mexican Rice Borer ¹ Rice Stalk Borer ¹ Southern Corn Leaf Beetle ³ Sugarcane Borer ¹ | 0.03 | 1.92 |

Remarks

- Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gal of water per acre.
- For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3- to 5-day intervals if needed. Lambda T-2 may only suppress heavy infestations and/or subsequent migrations.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 0.03 lb ai (1.92 fl oz of product) per acre.
- **Do not** apply within 21 days of harvest.
- **Do not** allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. **Do not** feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.
- **Do not** apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per crop from at plant and foliar applications.
- **Do not** apply more than 0.06 lb ai (3.84 fl oz or 0.24 pt of product) per acre after silk initiation.
- **Do not** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre after corn has reached the milk stage (yellow kernels with milky fluid).

¹ For control before the larva bores into the plant stalk or ear.

² Use higher rates for large larvae.

³ Suppression only

⁴ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|----------------------|---|-------------|-------------|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS | | | |
| Sweet Corn (Foliar) | Aphid species ^{2,3} Armyworm ¹ Aster Leafhopper Beet Armyworm ^{1,3} Chinch Bug Common Cornstalk Borer Corn Earworm Corn Rootworm Beetle (Adult): Mexican Northern Southern Western Cutworm species | 0.02 – 0.03 | 1.28 – 1.92 |

| Crop | Target Pests | Rate | |
|----------------------------------|---|-------------|-------------|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS (continued) | | | |
| Sweet Corn (Foliar) | European Corn Borer Fall Armyworm ¹ Flea Beetle species Grasshopper species Japanese Beetle (Adult) Sap Beetle (Adult) Southern Armyworm ¹ Southwestern Corn Borer Spider Mite species ² Stink Bug species Tarnished Plant Bug Webworm species Western Bean Cutworm Yellowstriped Armyworm ¹ | 0.02 – 0.03 | 1.28 – 1.92 |
| | Corn Silkworm (Adult) ² | 0.03 | 1.92 |

Remarks

- Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 4 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods and should be targeted for control before insects enter the stalk or ear.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage and ears (if present). When applying by air, apply in a minimum of 2 gal of water per acre.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 0.025 lb ai (1.60 fl oz of product) per acre.
- **Do not** apply within 1 day of harvest.
- **Do not** allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. **Do not** feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.
- **Do not** apply more than 0.48 lb ai (30.72 fl oz or 1.92 pt of product) per acre per crop from at plant and foliar applications.

¹ Use higher rates for large larvae.

² Suppression only

³ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|----------------------|---|--------------|------------|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS | | | |
| Rice Wild Rice | Bird Cherry-Oat Aphid Chinch Bug Fall Armyworm Grasshopper species Greenbug Leafhopper species Rice Stink Bug Rice Water Weevil (adult) Riceworm Sharpshooter species True Armyworm Yellow Sugarcane Aphid Yellowstriped Armyworm | 0.025 - 0.04 | 1.6 - 2.56 |

continued...

| Crop | Target Pests | Rate | |
|----------------------------------|--|-------------|-------------|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS (continued) | | | |
| Rice Wild Rice | European Corn Borer ¹ Mexican Rice Borer ¹ Rice Seed Midge ¹ Rice Stalk Borer ¹ Sugarcane Borer ¹ | 0.03 – 0.04 | 1.92 – 2.56 |

Remarks

- Apply as required by scouting. Timing and frequency of application should be based upon insect populations reaching locally determined economic thresholds. Determine the need for repeat applications, usually at intervals of 5 - 7 days, by scouting.
- Lamba T-2 can be safely used when propanil products are being used for weed control.
- Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water (or total carrier volume) per acre, but ensure sufficient volume is used to provide adequate coverage. In addition, adding an emulsified crop oil (e.g., 1 pt per acre) when lower aerial application volumes are used is recommended to help improve coverage, reduce evaporation and improve efficacy.
- For control of rice water weevil in dry-seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations.
- For control of rice water weevil in water-seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars, usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations.
- California: In addition to above directions for control of rice water weevil in water seeded rice, Lamba T-2 may be applied at the 1-3 leaf growth stage, with the majority at the 2-leaf growth stage. Adults are vulnerable on levees and in the water. Larvae are vulnerable while feeding on the leaf prior to entering the soil. Monitor for adults, based upon field history and density of population. Monitor field edges and levee areas for adults. Treat in the following manner: a) spray the inside perimeter of the field, or b) spray the entire field.
- Greenbug is known to have many biotypes. Lamba T-2 may only provide suppression. If satisfactory control is not achieved with the first application of Lamba T-2, a resistant biotype may be present. Use alternate chemistry for control.
- For control of stem borers, scout fields, when rice growth is near panicle differentiation, for early symptoms of damaging populations exhibited as discoloration (orange-tan) around the junction of the leaf sheath and leaf blade which is caused by feeding of young larvae within the sheath. Applications must be made before larvae bore into rice stems. Make the first application at panicle differentiation to 2 inch panicle for partial control. Make the second application at boot to heading for maximum control. All rice varieties are susceptible to stem borer damage, but Cocodrie and Priscilla are particularly susceptible.
- Mixers/loaders supporting aerial applications to wild rice at a rate of 0.04 lb ai per acre, and treating 1200 acres (or more) per day must wear dust-mist respirator.
- **Do not** release flood water within 7 days of an application.
- **Do not** apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per season.
- **Do not** apply more than 0.04 lb ai (2.56 fl oz or 0.16 pt of product) per acre within 21 to 27 days of harvest.
- **Do not** apply within 21 days of harvest.
- **Do not** use treated rice fields for the aquaculture of edible fish and crustacea.
- **Do not** apply as an ultra-low volume (ULV) spray.

¹ For control before the larvae bores into the plant stalk.

| Crop | Target Pests | Rate | |
|----------------------|--|--------------|-------------|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS | | | |
| Sorghum (Grain) | Cutworm species Sorghum Midge | 0.015 – 0.02 | 0.96 – 1.28 |
| | Armyworm Beet Armyworm ³ Corn Earworm European Corn Borer ² Fall Armyworm ¹ Flea Beetle species Grasshopper species Lesser Cornstalk Borer ² Southwestern Corn Borer ² Stink Bug species Webworm species Yellowstriped Armyworm ¹ | 0.02 – 0.03 | 1.28 – 1.92 |
| | Chinch Bug Mexican Rice Borer ² Rice Stalk Borer ² Sugarcane Borer ² | 0.03 | 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or aerial equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gal of water per acre.
- For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5-day intervals if needed.
- For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3- to 5-day intervals if needed. Lambda T-2 may only suppress heavy infestations and/or subsequent migrations.
- **Do not** apply more than 0.08 lb ai (5.12 fl oz or 0.32 pt of product) per acre per season.
- **Do not** apply more than 0.06 lb ai (3.84 fl oz or 0.24 pt of product) per acre per season after crop emergence.
- **Do not** apply more than 0.02 lb ai (1.28 fl oz or 0.08 pt of product) per acre per season once crop is in soft-dough stage.
- **Do not** apply within 30 days of harvest.

¹ Use higher rates for large larvae.

² For control before the larva bores into the plant stalk.

³ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|----------------------|------------------------------------|---------------|-------------|
| | | lb ai/A | fl oz/A |
| CEREAL GRAINS | | | |
| Barley | Army Cutworm | 0.015 – 0.025 | 0.96 – 1.60 |
| Buckwheat | Cutworm species | | |
| Oats | Armyworm | 0.02 – 0.03 | 1.28 – 1.92 |
| Rye | Bird Cherry-Oat Aphid ¹ | | |
| Triticale | Cereal Leaf Beetle | | |
| Wheat | English Grain Aphid ¹ | | |
| Wheat Hay | Fall Armyworm | | |
| | Flea Beetle species | | |
| | Grasshopper species | | |
| | Hessian Fly ⁴ | | |
| | Orange Blossom Wheat Midge | | |
| | Russian Wheat Aphid ¹ | | |
| | Stink Bug species | 0.025 – 0.03 | 1.60 – 1.92 |
| | Yellowstriped Armyworm | | |
| | Grass Sawfly | 0.03 | 1.92 |
| | Chinch Bug | | |
| | Corn Leaf Aphid ² | | |
| | Greenbug ^{1,3} | | |
| | Mite species ² | | |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- For chinch bug control, repeat applications at 3- to 5-day intervals if needed. Lambda T-2 may only suppress heavy infestations and/or migrations.
- Greenbug is known to have many biotypes. Lambda T-2 may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.
- **Do not** apply within 30 days of harvest.
- **Do not** allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals within 7 days after treatment. **Do not** feed treated straw to meat or dairy animals within 30 days after the last treatment.
- **Do not** apply more than 0.06 lb ai (3.84 fl oz or 0.24 pt of product) per acre per season.

¹ Best control is obtained before insects begin to roll leaves. Once crop has started to boot, Lambda T-2 may provide suppression only. Higher rates and increased coverage will be necessary.

² Suppression only

³ See Resistance statement under **General Directions for Use**.

⁴ Make applications when adults emerge.

| Crop | Target Pests | Rate | |
|--|--|---------------|-------------|
| | | lb ai/A | fl oz/A |
| COLE CROPS (HEAD AND STEM BRASSICA) | | | |
| Broccoli Brussels Sprouts Cabbage Cavalo Broccolo Cauliflower Chinese Broccoli (gai lon) Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy) Kohlrabi | Alfalfa Looper Cabbage Looper Cabbage Webworm Cutworm species Imported Cabbageworm Southern Cabbageworm | 0.015 – 0.025 | 0.96 – 1.60 |
| | Aphid species ^{2,3} Armyworm Beet Armyworm ^{1,3} Corn Earworm Diamondback Moth ³ Fall Armyworm ¹ Flea Beetle species Grasshopper species Japanese Beetle (Adult) Leafhopper species Meadow Spittlebug Plant Bug species including Lygus species ³ Spider Mite species ² Stink Bug species Thrips species ² Vegetable Weevil (Adult) Whitefly species ^{2,3} Yellowstriped Armyworm | 0.02 – 0.03 | 1.28 – 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- **Do not** apply within 1 day of harvest.
- **Do not** apply more than 0.24 lb ai (15.36 fl oz or 0.96 pt of product) per acre per season.

¹ For control of first and second instar only.

² Suppression only

³ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|---------------|--|--------------|-------------|
| | | lb ai/A | fl oz/A |
| COTTON | | | |
| | Cutworm species Soybean Thrips Tobacco Thrips | 0.015 – 0.02 | 0.96 – 1.28 |
| | Cabbage Looper Cotton Fleahopper Cotton Leafperforator Cotton Leafworm Lygus Bug species ³ Pink Bollworm Saltmarsh Caterpillar | 0.02 – 0.03 | 1.28 – 1.92 |
| | Bandedwing Whitefly ^{2,3} Beet Armyworm ^{1,3} Boll Weevil Brown Stink Bug Cotton Aphid ^{2,3} Cotton Bollworm European Corn Borer Fall Armyworm Green Stink Bug Southern Green Stink Bug Sweet Potato Whitefly ^{2,3} Tobacco Budworm ³ Twospotted Spider Mite ² | 0.025 – 0.04 | 1.60 – 2.56 |

Remarks

- Apply as required by scouting, usually at intervals of 5 - 7 days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or aerial equipment using sufficient water to obtain full coverage of foliage.
- Applications may also be made with equipment adapted and calibrated for ULV sprays. Lamba T-2 may be mixed with once-refined vegetable oil and applied in a minimum of at least one qt of finished spray per acre.
- Under light bollworm/budworm infestation levels, 0.02 lb ai (1.28 fl oz of product) per acre may be applied in conjunction with intense field monitoring.
- For boll weevil control, spray on a 3- to 5-day schedule.
- When applied according to label directions for control of cotton bollworm and tobacco budworm, Lamba T-2 also provides ovicidal control of unhatched *Heliothis* species eggs.
- **Do not** apply within 21 days of harvest.
- **Do not** graze livestock in treated areas.
- **Do not** apply more than 0.2 lb ai (12.8 fl oz or 0.8 pt of product) per acre per season.
- **Do not** make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season.

¹ For control of the first and second instar only.

² Suppression only

³ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|---|---|-----------|-----------|
| | | lb ai/A | fl oz/A |
| CUCURBIT VEGETABLES | | | |
| Chayote (fruit) Chinese Waxgourd (Chinese preserving melon) Citron Melon Cucumber Gherkin Gourd (edible) <i>Lagenaria</i> species – includes: hyotan, cucuzza <i>Luffa acutangula</i> , <i>L. cylindrical</i> – includes: hechima, Chinese okra <i>Momordica</i> species – includes: balsam apple, balsam pear, bitter melon, Chinese cucumber Muskmelon (hybrids and/or cultivars of <i>Cucumis melo</i>) – includes: true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon Pumpkin Squash, summer (<i>Cucurbita pepo</i> var. <i>melopepo</i>) – includes: crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini Squash, winter (<i>Cucurbita maxima</i> ; <i>C. moschata</i>) – includes butternut squash, calabaza, hubbard squash (<i>C. mixta</i> ; <i>C. pepo</i>) – includes: acorn squash, spaghetti squash Watermelon – includes: hybrids and/or varieties of <i>Citrullus lanatus</i> | Armyworm species ¹ Blister Beetle species Cabbage Looper Corn Earworm Cricket species Cucumber Beetle species (adults) Cutworm species Flea Beetle species Grasshopper species June Beetle species Leaffooted Bug Leafhopper species Lygus Bug species ¹ Melonworm Pickleworm Plant Bug species Rindworm species complex Saltmarsh Caterpillar Squash Beetle Squash Bug species Squash Vine Borer species Stink Bug species Thrips species ^{1, 2} Tobacco Budworm ¹ Webworm species | 0.02-0.03 | 1.28-1.92 |
| | Aphid species ¹ Leafminer species ^{1, 3} Whitefly species ^{1, 3} Spider Mite species ³ | 0.03 | 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of all plant parts. When applying by air, apply in a minimum of 2 gal total solution per acre. When applying by ground, a minimum of 10 gal total solution per acre is recommended.
- Use higher application volumes and/or rates when foliage is dense, pest populations are high, larvae are large, weather conditions are adverse and/or as plant size increases. Use higher rates for longer residual.
- Insects that bore or tunnel into leaves, vines, stems or fruit must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of Lamba T-2.
- **Do not** apply more than 0.18 lb ai (11.5 fl oz or 0.72 pt of product) per acre per season. **Do not** apply within 1 day of harvest.

continued...

¹ See Resistance statement under General Directions for Use.

² Does not include Western Flower Thrips

³ Suppression only

| Crop | Target Pests | Rate | |
|----------------------------|-------------------------------------|---------------|-------------|
| | | lb ai/A | fl oz/A |
| FRUITING VEGETABLES | | | |
| Eggplant | Cabbage Looper | 0.015 – 0.025 | 0.96 – 1.60 |
| Ground cherry | Cutworm species | | |
| Pepino | Hornworm species | 0.02 – 0.03 | 1.28 – 1.92 |
| Peppers (bell and nonbell) | Aphid species ^{2,3} | | |
| Tomatillo | Beet Armyworm ^{1,3} | | |
| Tomato | Blister Beetle species | | |
| | Colorado Potato Beetle ³ | | |
| | Cucumber Beetle species (Adult) | | |
| | European Corn Borer ⁴ | | |
| | Fall Armyworm ¹ | | |
| | Flea Beetle species | | |
| | Grasshopper species | | |
| | Japanese Beetle (Adult) | | |
| | Leafhopper species | | |
| | Leafminer species ² | | |
| | Meadow Spittlebug | | |
| | Pepper Weevil (Adult) ² | | |
| | Plant Bug species | | |
| | Southern Armyworm ¹ | | |
| | Spider Mite species ² | | |
| | Stalk Borer ⁴ | | |
| | Stink Bug species | | |
| | Thrips ⁵ | | |
| | Tobacco Budworm ³ | | |
| | Tomato Fruitworm | | |
| | Tomato Pinworm | | |
| | Tomato Psyllid ^{2,3} | | |
| | Vegetable Weevil (Adult) | | |
| | Whitefly species ^{2,3} | | |
| | Yellowstriped Armyworm ¹ | | |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- **Do not** apply within 5 days of harvest.
- **Do not** apply more than 0.36 lb ai (23.04 fl oz or 1.44 pt of product) per acre per season.

¹ For control of first and second instar only

² Suppression only

³ See Resistance statement under General Directions for Use.

⁴ For control before the lava bores into the plant stalk or fruit

⁵ Does not include Western Flower Thrips.

| Crop | Target Pests | Rate | |
|---|---|-------------|-----------|
| | | lb ai/A | fl oz/A |
| GRASS FORAGE, FODDER AND HAY | | | |
| Pasture and Rangeland Grass, Grass Grown for Hay or Silage and Grass Grown for Seed | Army Cutworm Cutworm species Essex Skipper Range Caterpillar Striped Grass Looper | 0.015-0.025 | 0.96-1.6 |
| | Beet Armyworm Billbug species ³ Bird Cherry-Oat Aphid ¹ Black Grass Bug Black Turfgrass Beetle (adult) Blue Stem Midge Cereal Leaf Beetle Chinch Bug Crane Fly species Cricket species English Grain Aphid ¹ Fall Armyworm Flea Beetle species Grass Mealybug Grass Sawfly (adult) Grasshopper species Green June Beetle (adult) Greenbug ^{1, 2} Japanese Beetle (adult) Katydid species Leafhopper species Mite species ³ Russian Wheat Aphid ¹ Southern Armyworm Spittlebug species Stink Bug species Sugarcane Aphid Thrips species Tick species True Armyworm Webworm species Yellowstriped Armyworm | 0.02-0.03 | 1.28-1.92 |

Remarks

- Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal total solution per acre. When applying by ground, a minimum of 7 gal total solution per acre is recommended.
- Use higher application volumes and rates when foliage is dense, pest populations are high, larvae are large and/or weather conditions are adverse. Use higher rates for longer residual.
- For chinch bug control, Lamba T-2 may only suppress heavy infestations and/or migrations. In this situation, a second application using an alternative chemistry may be needed.
- Greenbug is known to have many biotypes. Lamba T-2 may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.

- Pasture and rangeland grass may be used for grazing or cut for forage 0 days after application. **Do not** cut grass to be dried and harvested for hay until 7 days after the last application.
Grass grown for seed:
 - Straw, hay and mature seed (seed screenings) may be used as feed 7 days after the last application. Regrowth of grass grown for seed may be used for grazing, cut for forage or cut to be dried and harvested for hay.
 - **Do not** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per cutting for pastures, rangeland and grasses grown for seed. A minimum re-treatment interval (RTI) of 30 days is required for pastures and rangeland receiving 0.03 lb ai per acre which have not been cut between applications.
 - **Do not** apply more than 0.09 lb ai (5.76 fl oz or 0.36 pt of product) per acre per season.
- 1 Best control is obtained before insects begin to roll leaves.
 2 See **Resistance** statement under **General Directions for Use**.
 3 Suppression only

| Crop | Target Pests | Rate | |
|---|---|---------------|-------------|
| | | lb ai/A | fl oz/A |
| LEGUME VEGETABLES (BEANS AND PEAS) | | | |
| Edible Podded (Only) | | | |
| <i>Canavalia ensiformis</i> – jackbean | Cutworm species Green Cloverworm Imported Cabbageworm | 0.015 – 0.025 | 0.96 – 1.60 |
| <i>Canavalia gladiata</i> – sword bean | Mexican Bean Beetle Saltmarsh Caterpillar Velvetleaf Caterpillar | | |
| <i>Glycine max</i> – soybean (immature seed) | Alfalfa Caterpillar Aphid species ⁴ Armyworm ² | 0.02 – 0.03 | 1.28 – 1.92 |
| Edible Podded, Succulent Shelled or Dried Shelled | | | |
| <i>Cajanus cajan</i> – Pigeon pea | Bean Leaf Beetle Bean Leafskeltonizer Blister Beetle species Corn Earworm | | |
| <i>Phaseolus</i> species – includes: field, kidney, lima, navy, pinto, runner, snap, tepary and wax beans | Corn Rootworm Beetle species (Adult) Cucumber Beetle species (Adult) Curculio and Weevil species ¹ (foliage and pod feeding adults and larvae) European Corn Borer Fall Armyworm ² | | |
| <i>Pisum</i> species – includes: dwarf, edible-pod, English, field, garden, green, snow and sugar snap peas | Flea Beetle species (Adult) Flea Hopper species Grasshopper species Japanese Beetle (Adult) | | |
| <i>Vigna</i> species – includes: adzuki, asparagus, moth, mung, rice, urd and yardlong beans, black-eye pea, catjang, Chinese longbean, cowpea, Crowder pea, and Southern pea | Leafhopper species Leaf-tier species Looper Species Meadow Spittlebug Painted Lady Butterfly (Larva) Plant Bug species including Lygus species ⁴ Stalk Borer ¹ Stink Bug species Threecornered Alfalfa Hopper Thrips species ^{4,5} Tobacco Budworm ⁴ Webworm species Western Bean Cutworm Western Yellowstriped Armyworm ² Yellowstriped Armyworm ² | | |

| Crop | Target Pests | Rate | |
|--|---|---------|---------|
| | | lb ai/A | fl oz/A |
| LEGUME VEGETABLES (BEANS AND PEAS) | | | |
| (continued) | Beet Armyworm ^{3,4} Leafminer species ^{3,4} Lesser Cornstalk Borer ³ Soybean Looper ^{3,4} Spider Mite species ³ Whitefly species ^{3,4} | 0.03 | 1.92 |
| Succulent Shelled or Dried Shelled | | | |
| <i>Vicia faba</i> – broadbean (favabean) | | | |
| Dried Shelled (Only) | | | |
| <i>Cicer arietinum</i> – chickpea (garbonzo bean) | | | |
| <i>Cyamopsis tetragonoloba</i> – guar | | | |
| <i>Lablab purpureus</i> – Lablab bean (hyacinth bean) | | | |
| <i>Lupinus</i> species – includes: grain, sweet, white and sweet white lupines | | | |
| <i>Lens esculata</i> – Lentils | | | |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- For edible podded and succulent shelled legume vegetables, do not apply within 7 days of harvest.
- For dried shelled legume vegetables, do not apply within 21 days of harvest.
- **Do not** apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per season.
- For succulent and dried shelled peas and beans, **do not** graze livestock in treated areas or harvest vines for forage or hay.

¹ For control before the larva bores into the plant stalk or pods.

² Use higher rates for large larvae.

³ For suppression only

⁴ See **Resistance** statement under **General Directions for Use**.

⁵ Does not include Western Flower Thrips.

| Crop | Target Pests | Rate | |
|-------------------------------------|---|---------------|-------------|
| | | lb ai/A | fl oz/A |
| LEGUME VEGETABLES (SOYBEANS) | | | |
| Soybeans | Bean Leaf Beetle Cabbage Looper Corn Earworm Corn Rootworm Beetle (Adult): Mexican Northern Southern Western | 0.015 – 0.025 | 0.96 – 1.60 |

continued...

| Crop | Target Pests | Rate | |
|---|---|---------------|-------------|
| | | lb ai/A | fl oz/A |
| LEGUME VEGETABLES (SOYBEANS) <i>(continued)</i> | | | |
| Soybeans | Cutworm species Green Cloverworm Mexican Bean Beetle Painted Lady (Thistle) Caterpillar Potato Leafhopper Saltmarsh Caterpillar Soybean Aphids ⁴ Threecornered Alfalfa Hopper Thrips species ⁵ Velvetbean Caterpillar Woollybear Caterpillar | 0.015 – 0.025 | 0.96 – 1.60 |
| | Armyworm ¹ Blister Beetle species European Corn Borer Fall Armyworm ¹ Grasshopper species Japanese Beetle (Adult) Plant Bug species Silverspotted Skipper Stink Bug species Tobacco Budworm ³ Webworm species Yellowstriped Armyworm ¹ | 0.025 – 0.03 | 1.60 – 1.92 |
| | Beet Armyworm ^{2,3} Lesser Cornstalk Borer ² Soybean Looper ^{2,3} Spider Mite species ² | 0.03 | 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- **Do not** graze or harvest treated soybean forage, straw, or hay for livestock feed.
- Apply with ground or aerial equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial-applied corn rootworm control program use a minimum of 0.02 lb ai (1.28 fl oz of product) per acre.
- **Do not** apply within 30 days of harvest.
- **Do not** apply more than 0.06 lb ai (3.84 fl oz or 0.24 pt of product) per acre per season.

¹ Use higher rates for large larvae.

² Suppression only

³ See **Resistance** statement under **General Directions for Use**.

⁴ Use lower rates for early season applications and/or lighter populations.

⁵ Does not include Western Flower Thrips.

| Crop | Target Pests | Rate | |
|--------------------------------|---|---------------|-------------|
| | | lb ai/A | fl oz/A |
| LETTUCE (HEAD AND LEAF) | | | |
| | Alfalfa Looper Cabbage Looper Cutworm species Green Cloverworm Imported Cabbageworm Saltmarsh Caterpillar | 0.015 – 0.025 | 0.96 – 1.60 |
| | Aphid species ^{2,3} Armyworm Beet Armyworm ^{1,3} Corn Earworm Diamondback Moth ³ European Corn Borer Fall Armyworm ¹ Flea Beetle species Grasshopper species Japanese Beetle (Adult) Leafhopper species Meadow Spittlebug Plant Bug species including Lygus species ³ Southern Armyworm Spider Mite species ² Stink Bug species Tobacco Budworm ³ Vegetable Weevil (Adult) Whitefly species ^{2,3} | 0.02 – 0.03 | 1.28 – 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- **Do not** apply within 1 day of harvest.
- **Do not** apply more than 0.3 lb ai (19.2 fl oz or 1.2 pt of product) per acre per season.

¹ For control of first and second instar only

² Suppression only

³ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|--------------------------------|---|---------------|-------------|
| | | lb ai/A | fl oz/A |
| ONION (BULB) AND GARLIC | | | |
| | Cutworm species Leafminer species (Adult) Onion Maggot (Adult) Seedcorn Maggot (Adult) | 0.015 – 0.025 | 0.96 – 1.60 |
| | Aphid species ² Armyworm species ¹ Flower Thrips ^{4,3} Onion Thrips ³ Plant Bug species Stink Bug species Tobacco Thrips ³ Western Flower Thrips ^{2,3} | 0.02 – 0.03 | 1.28 – 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Use the higher label rates as thrips population increases and avoid rescue situations.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- For thrips control by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturers use directions) may enhance the deposition of the spray and increase plant coverage.
- **Do not** apply within 14 days of harvest.
- **Do not** apply more than 0.24 lb ai (15.36 fl oz or 0.96 pt of product) per acre per season.

¹ For control of the first and second instar only

² Suppression only

³ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|----------------|---|---------------|-------------|
| | | lb ai/A | fl oz/A |
| PEANUTS | | | |
| | Cutworm species Green Cloverworm Potato Leafhopper Rednecked Peanut Worm Threecornered Alfalfa Hopper Velvetbean Caterpillar | 0.015 - 0.025 | 0.96 - 1.60 |
| | Bean Leaf Beetle Corn Earworm Fall Armyworm ¹ Grasshopper species Southern Corn Rootworm (Adult) Stink Bug species Tobacco Thrips Vegetable Weevil Whitefringed Beetle (Adult) | 0.02 - 0.03 | 1.28 - 1.92 |
| | Aphid species ² Beet Armyworm ^{2,3} Lesser Cornstalk Borer ² Soybean Looper ^{2,3} Spider Mite species ² | 0.03 | 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or aerial equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- **Do not** apply within 14 days of harvest.
- **Do not** apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per season.

¹ Use higher rates for large larvae.

² Suppression only

³ See **Resistance** statement under **General Directions for Use**

| Crop | Target Pests | Rate | |
|---|--|-------------|-------------|
| | | lb ai/A | fl oz/A |
| POME FRUITS | | | |
| Apple Crabapple Loquat Mayhaw Oriental Pear Pear Quince | Apple Aphid Apple Maggot (Adult) Cherry Fruit Fly species (Adult) Codling Moth Green Fruitworm Japanese Beetle Leafhopper species Leafroller species Lesser Appleworm Omnivorous Leafroller Orange Tortrix Oriental Fruit Moth Pear Psylla ¹ Pear Sawfly Periodical Cicada Plant Bug species Plum Curculio Rosy Apple Aphid San Jose Scale (fruit infestations only) Spirea Aphid ¹ Stink Bug species Tent Caterpillar species Tentiform Leaf Miner species Tree Borer species Tufted Apple Budworm Webworm species | 0.02 – 0.04 | 1.28 – 2.56 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds and IPM recommendations.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gal of water per acre, but use higher volumes as appropriate for thorough coverage.
- **Do not** apply within 21 days of harvest.
- **Do not** apply more than 0.2 lb ai (12.8 fl oz or 0.80 pt of product) per acre per year.
- **Do not** apply more than 0.16 lb ai (10.24 fl oz or 0.64 pt of product) per acre per year post bloom.

¹Suppression only

| Crop | Target Pests | Rate | |
|--|--|-------------|-------------|
| | | lb ai/A | fl oz/A |
| STONE FRUITS | | | |
| Apricot Chickasaw Plum Damson Plum Japanese Plum Nectarine Peach Plum Plumcot Prune Sweet and Tart Cherry | American Plum Borer Apple Maggot (Adult) Black Cherry Aphid Cherry Fruit Fly species (Adult) Codling Moth Green Fruitworm Japanese Beetle June Beetle Leafhopper species Leafroller species Oriental Fruit Moth Peach Twig Borer Peachtree Borer species Pear Sawfly Periodical Cicada Plant Bug species Plum Curculio Rose Chafer Stink Bug species Tent Caterpillar species Thrips species | 0.02 – 0.04 | 1.28 – 2.56 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold and IPM recommendations.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 5 gal of water/per acre, but use higher volumes as appropriate for thorough coverage.
- **Do not** apply within 14 days of harvest.
- **Do not** apply more than 0.2 lb ai (12.8 fl oz or 0.80 pt of product) per acre per year.
- **Do not** apply more than 0.16 lb ai (10.24 fl oz or 0.64 pt of product) per acre per year post bloom.

| Crop | Target Pests | Rate | |
|------------------|---|--------------|-------------|
| | | lb ai/A | fl oz/A |
| SUGARCANE | | | |
| | Mexican Rice Borer ¹ Pygmy Mole Cricket Rice Stalk Borer ¹ Sugarcane Aphid ³ Sugarcane Beetle (Adult) ² Sugarcane Borer ¹ West Indian Crane-fly Yellow Sugarcane Aphid ³ | 0.025 – 0.04 | 1.60 – 2.56 |

Remarks

- Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 2 gal of water per acre.
- **Do not** apply within 21 days of harvest.
- **Do not** apply more than 0.16 lb ai (10.24 fl oz or 0.64 pt of product) per acre per season.

¹ For control before the larva bores into the plant stalk

² Suppression only of beetles active above ground

³ See Resistance statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|------------------|--|---------------|-------------|
| | | lb ai/A | fl oz/A |
| SUNFLOWER | | | |
| | Cutworm species Sunflower Beetle | 0.015 – 0.025 | 0.96 – 1.60 |
| | Banded Sunflower Moth Fall Armyworm ¹ Grasshopper species Head-Clipper Weevil (Adult) Japanese Beetle (Adult) Leafhopper species Meadow Spittlebug Painted Lady (Thistle) Caterpillar Seed Weevil (Adult) Spotted Cabbage Looper Stem Weevil (Adult) Stink Bug species Sunflower Maggot (Adult) Sunflower Moth Woollybear Caterpillar | 0.02 – 0.03 | 1.28 – 1.92 |
| | Beet Armyworm ^{2,3} Spider Mite species ² | 0.03 | 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

- Apply with ground or air equipment using sufficient water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- **Do not** apply within 45 days of harvest.
- **Do not** apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per season.
- **Do not** apply more than 0.09 lb ai (5.76 fl oz or 0.36 pt of product) per acre per season after bloom initiation.
- **Do not** apply as an ultra-low volume (ULV) spray.

¹ Use higher rates for large larvae.

² Suppression only

³ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|----------------|--|--------------|-------------|
| | | lb ai/A | fl oz/A |
| TOBACCO | | | |
| | Armyworm species ¹ Blister Beetle species Cabbage Looper Corn Earworm Cucumber Beetle species (Adult) Cutworm species Grasshopper species Japanese Beetle (Adult) Katydid species Plant Bug species ³ Potato Tuberworm Salt Marsh Caterpillar Stinkbug species Tobacco Aphid species ^{2,3} Tobacco Budworm ³ Tobacco Flea Beetle (Adult) Tobacco Hornworm Tobacco Thrips species ² Tomato Hornworm Tree Cricket species Vegetable Weevil (Adult) Webworm species | 0.015 – 0.03 | 0.96 – 1.92 |

Remarks

- Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage. When applying by air, apply in a minimum of 2 gal of water per acre.
- **Do not** apply within 40 days of harvest.
- **Do not** apply more than 0.09 lb ai (5.76 fl oz or 0.36 pt of product) per acre per year.

¹ For control of first and second instars only

² Suppression only

³ See **Resistance** statement under **General Directions for Use**.

| Crop | Target Pests | Rate | |
|--|--|-------------|-------------|
| | | lb ai/A | fl oz/A |
| TREE NUTS | | | |
| Almond Beech Nut Brazil Nut Butternut Cashew Chestnut Chinquapin Filbert (Hazlenut) Hickory Nut Macadamia Nut (Bush Nut) Pistachio Walnut, Black Walnut, English (Persian) | Ants Chinch Bug Codling Moth Filbertworm Leaffooted Bug Leafroller species Navel Orangeworm Peach Twig Borer Plant Bug species Stink Bug species Walnut Aphid Walnut Husk Fly species (Adult) | 0.02 – 0.04 | 1.28 – 2.56 |
| Pecan | Hickory Shuckworm Pecan Aphid species Pecan Casebearer species Pecan Phylloxera species Pecan Spittlebug Pecan Weevil Stink Bug species | 0.02 – 0.04 | 1.28 – 2.56 |

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gal of water/per acre, but use higher rates as appropriate for thorough coverage.
- **Do not** apply within 14 days of harvest.
- **Do not** apply more than 0.16 lb ai (10.24 fl oz or 0.64 pt of product) per acre per year.
- **Do not** apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per year post bloom.

| Crop | Target Pests | Rate | |
|--|-------------------------------------|-------------|-----------|
| | | lb ai/A | fl oz/A |
| TUBEROUS AND CORM VEGETABLES (Potato, Sweet Potato, Yams and Related) | | | |
| Arracacha | Cutworm species | 0.015-0.025 | 0.96-1.60 |
| Arrowroot | Leafhopper species | | |
| Artichoke (Chinese and Jerusalem only) | Saltmarsh Caterpillar | 0.02-0.03 | 1.28-1.92 |
| Canna (edible) | Sweet Potato Hornworm | | |
| Cassava (bitter and sweet) | Woollybear Caterpillar species | | |
| Chayote (root) | Aphid species ¹ | | |
| Chufa | Armyworm species ¹ | | |
| Dasheen | Blister Beetle species | | |
| Ginger | Colorado Potato Beetle ¹ | | |
| Leren | Corn Earworm | | |
| Potato | Cricket species | | |
| Sweet Potato | Cucumber Beetle species (adults) | | |
| Tanier | European Corn Borer | | |
| Turmeric | Flea Beetle species (adults) | | |
| Yam (bean and true) | Grasshopper species | | |
| | Looper species ¹ | 0.03 | 1.92 |
| | Lygus Bug species ¹ | | |
| | Plant Bug species | | |
| | Potato Psyllid | | |
| | Potato Tuberworm | | |
| | Stink Bug species | | |
| | Sweet Potato Leaf Beetle (adults) | | |
| | Sweet Potato Vine Borer | | |
| | Thrips species ^{1, 2} | | |
| | Tortoise Beetle species | | |
| | Webworm species | | |
| | Weevil species (adults) | | |
| | Leafminer species ^{1, 3} | | |
| | Spider Mite species ³ | | |
| | Whitefly species ^{1, 3} | | |

Remarks

- Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of all above ground plant parts. When applying by air, apply in a minimum of 2 gal total solution per acre. When applying by ground, a minimum of 10 gal total solution per acre is recommended.
- Use higher application volumes and/or rates when foliage is dense, pest populations are high, larvae are large, weather conditions are adverse and/or as plant size increases. Use higher rates for longer residual.
- Insects that bore or tunnel into leaves, vines, stems, tubers or corms must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of Lamba T-2.
- Do not apply more than 0.12 lb ai (7.68 fl oz or 0.48 pt of product) per acre per season.
- Do not apply within 7 days of harvest.

¹ See **Resistance** statement under **General Directions for Use**.

² Does not include Western Flower Thrips.

³ Suppression only

| NON-AGRICULTURAL USES | | | |
|------------------------------------|--|-------------|-------------|
| Crop | Target Pests | Rate | |
| | | lb ai/A | fl oz/A |
| CONIFER AND DECIDUOUS TREES | | | |
| Plantations and Nurseries | Bagworm Balsam Twig Aphid Balsam Woolly Aphid Birch Leafminer Black Pine Weevil Elm Leaf Beetle European Elm Bark Beetle Gypsy Moth Japanese Beetle June Beetle species Leaf Beetle species Leafroller species May Beetle species Mealybug species ¹ Pales Weevil Pine Chafer Pine Colaspis Beetle Pine Conelet Bug Pine Leaf Chermid Pine Needle Scale Pine Sawfly species Pine Tip Moth species Pine Tortoise Scale Pine Weevil species Poplar Aphid species Sawfly species Spittlebug species Spruce Budworm Tent Caterpillar species Tussock Moth species Webworm species | 0.02 – 0.04 | 1.28 – 2.56 |

Remarks

- To control exposed foliage, flower, cone, seed and bark feeding insects, apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground equipment using sufficient water to obtain full coverage of target site. When applying by air, apply a minimum of 2 gal of water per acre.
- **Do not** apply more than 0.24 lb ai (15.36 fl oz or 0.96 pt of product) per acre per year.

¹ Suppression only

| Crop | Target Pests | Rate | |
|------------------------------------|--|-------------|-------------|
| | | lb ai/A | fl oz/A |
| CONIFER AND DECIDUOUS TREES | | | |
| Seed Orchards | Coneworm species Seed Bug species Thrips species | See Remarks | See Remarks |

Remarks

- For high volume sprayers, dilute 2.56 fl oz per 100 gal of water and apply 5-10 gal of finished spray per tree.
- For low volume sprayers, dilute 10 fl oz per 100 gal of water and apply 100 gal of finished spray per acre.
- For aerial applications, apply 7.5 fl oz per acre in a minimum of 10 gal finish spray per acre.
- **Do not** apply more than 0.5 lb ai (32 fl oz or 2 pt of product) per acre per year.

| Crop | Target Pests | Rate | |
|--------------------------------------|--|------------------|------------------|
| | | lb ai/A | fl oz/A |
| Non-Cropland (Excluding Public Land) | See Crop Outlets on this Lamba T-2 label for target pests and rates. | See Crop Outlets | See Crop Outlets |

Remarks

- Spray non-cropland adjacent to agricultural areas to control migratory insects, which may threaten crops.
- Follow **General Use Directions**, rates and spray recommendations found elsewhere in this label for the adjacent crop outlet and target pests.
- Use highest labeled rates for dense/large foliage, high insect populations and larger larval stages.
- Repeat as necessary to maintain control.
- **Do not** exceed 0.2 lb ai (12.8 fl oz or 0.8 pt of product) per acre per year.
- **Do not** graze livestock in treated areas.

Rate Conversion Chart

| Pp | Fl oz Per Acre | Pints Per Acre | Treated Acres Per Gal |
|-------|----------------|----------------|-----------------------|
| 0.015 | 0.96 | 0.06 | 133 |
| 0.02 | 1.28 | 0.08 | 100 |
| 0.025 | 1.60 | 0.10 | 80 |
| 0.03 | 1.92 | 0.12 | 67 |
| 0.035 | 2.24 | 0.14 | 57 |
| 0.04 | 2.56 | 0.16 | 50 |

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area. **DO NOT ALLOW PRODUCT TO FREEZE.**

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons – bulk]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Viton® is a registered trademark of E.I. DuPont de Nemours and Company

©2016 Helena Holding Company

Manufactured for:
Helena Chemical Company
225 Schilling Boulevard, Suite 300
Collierville, Tennessee 38017

SCPSR-HEL-1295D-L1 1114
4062274
AD 111314

RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

GROUP 3 INSECTICIDE



Insecticide

ACTIVE INGREDIENT:
Lambda-cyhalothrin^{1,2} 22.8%
OTHER INGREDIENTS: 77.2%
TOTAL: 100.0%

Lambda T-2 contains 2.08 lb of active ingredient per gal and is a capsule suspension.

¹CAS No. 91465-08-6

²Synthetic pyrethroid

Contains petroleum distillate.

KEEP OUT OF REACH OF CHILDREN. WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

©2016 Helena Holding Company

NET CONTENTS: 1 gallon

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 - 30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Contains petroleum distillate - vomiting may cause aspiration pneumonia

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call CHEMTREC at 1-800-424-9300.

EPA Reg. No. 100-1295-5905 EPA Est. 39578-TX-1

SCPSR-HEL-1295D-L1 1114 4062274 AD 111314

MANUFACTURED FOR

HELENA CHEMICAL COMPANY

225 SCHILLING BOULEVARD, SUITE 300 • COLLIERVILLE, TENNESSEE 38017