FIFRA SECTION 24(c) SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OREGON

FORFEIT[™] 280

EPA Reg. No. 34704-1080 EPA SLN No.:OR-140011

For Weed Control in Grass Grown for Seed Production Only

This label is valid until December 31, 2019 or until otherwise amended, withdrawn, cancelled, or suspended.

ACTIVE INGREDIENT:

Glufosinate-ammonium*	. 24.5%**
OTHER INGREDIENTS:	75.5%
TOTAL	100.0%

^{*}CAS Number 77182-82-2

WARNING — AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This pesticide is toxic to vascular plants and should be used strictly in accordance with the drift and run-off precautions on this SLN label and the EPA-registered label in order to minimize off-site exposures. Under some conditions, FORFEIT 280 may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur to minimize water run-off is recommended.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this Special Local Need (SLN) label and on the EPA-registered label for FORFEITTM 280, EPA Reg. No. 34704-1080.
- This SLN label and the EPA registered product label must be in the possession of the user at the time of application.

^{**}Equivalent to 2.34 pounds of active ingredient per U.S. gallon.

Do not use this product until you have read the entire label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

GENERAL INFORMATION FOR WEED CONTROL IN GRASS GROWN FOR SEED PRODUCTION:

For use on grasses grown for seed production, such as, perennial ryegrass, tall fescue or bentgrass.

Tolerance of grasses to FORFEIT 280 is species dependent. Species tolerance in descending order is: 1) perennial ryegrass; 2) tall fescue; 3) bentgrass; 4) fine fescue; and 5) *Poa* species. For example, perennial ryegrass tends to be more tolerant to FORFEIT 280 than the *Poa* species.

Directed shielded sprays are suggested in fine fescue species, orchardgrass and *Poa* species grown for seed production. Broadcast applications made after first node development can increase the potential for crop injury and yield loss.

- 1. **DO NOT** apply FORFEIT 280 when grass grown for seed production is stressed due to drought, heat, frost, flooding, poor fertility, diseases, insects, or other reason.
- 2. Stunting of grass grown for seed production may occur following application and, in some instance, seed yields may be adversely affected.

DO NOT broadcast apply FORFEIT 280 on fine fescue, orchardgrass, or Poa species grown for seed.

WEEDS CONTROLLED OR SUPPRESSED:

Poa annua Manna Grass

Poa trivialis Bromus species (suppression only)

TIMING OF APPLICATION:

For best weed control and crop safety, apply when the grass seed crop is actively growing in the fall and/or spring to minimize potential for crop injury.

FALL APPLICATION

Broadcast Only:

For severe weed pressure in newly established grass seedling fields, applications may occur after the 1st tiller of the crop is established, but do not exceed the 10.0 fluid ounces rate per acre. For established fields with a minimum of 4 tillers, do not exceed a rate of 16.5 fluid ounces per acre. Do not make applications after December 1 in either seedling or established grass seed stands. Use a minimum of 20.0 gallons per acre of water at 30- 40 psi.

If additional control is required in the spring, one spring broadcast application may also be made.

SPRING APPLICATION

Broadcast:

Apply FORFEIT 280 to actively growing grass seed fields in the 4-6 tiller growth state at a rate of 16.5-20.0 fluid ounces per acre. Do not make applications after April 1 except when severe weed pressure necessitates control.

Hooded or Shielded Directed Spray

To avoid excessive crop phytotoxicity, use a hooded or shielded sprayer. Apply by directing spray between rows and using shielded directed sprayers to prevent spray contact with crop plants. Adjust shields to prevent spray from contacting the crop.

Apply 55.0 fluid ounces per acre with shielded directed spray when the weeds are less than 8 inches tall. Apply 82.0 fluid ounces per acre with shielded directed spray when the weeds are 8 inches or taller.

RESTRICTIONS TO THE DIRECTIONS FOR USE:

- 1. **DO NOT** graze in treated areas for one year (365 days) following application.
- 2. If you make a hooded or shielded direct spray application, do not also make a broadcast application.
- 3. **DO NOT** make more than one hooded or shielded direct spray application per crop year.
- 4. **DO NOT** make more than two broadcast applications per crop year (one in the fall and one in the spring).
- 5. **DO NOT** apply this product aerially.
- 6. **DO NOT** apply this product through any type of irrigation system.
- 7. **DO NOT** allow product to drift off the intended field of grass grown for seed production. Drift control is the responsibility of the applicator. Refer to local state laws, regulations, and guidelines for proper application to avoid off-target movement.

Forfeit is a registered trademark of Loveland Products, Inc.

24[c] Registrant: Loveland Products Inc. PO Box 1286 Greeley, CO 80632-1286