



## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR SPUR HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL

When using SPUR herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling.

# SPUR

EPA Reg. No. 42750-89

## FOR WEED CONTROL IN APPLE ORCHARDS

KEEP OUT OF REACH OF CHILDREN.

## CAUTION

In case of an emergency involving this product, Call CHEMTREC toll free at 1-800-424-9300

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

This supplemental labeling must be in the possession of the user at the time of pesticide application.

**APPLE**  
**Postemergence Broadleaf Weed Control in Apple Orchards**

(Not for Use or Distribution in the State of Florida or in Nassau and Suffolk Counties in New York State)

Target Broadleaf Weeds	Application Rate (pint/acre)	Use Restrictions
Aster Burdock Clover, red Clover, white Curly-dock Cocklebur Dandelion Goldenrod Horseweed Nightshade, black nightshade, hairy pineappleweed sowthistle, annual Thistle, canada Thistle, musk Vetch Volunteer alfalfa	1/4 - 2/3  (0.094 - 0.25 lb ae/acre)	<ul style="list-style-type: none"> <li>• Pre-Harvest Interval: Do not apply within 30 days of harvest.</li> <li>• Make one to two broadcast applications per crop per year. Do not apply more than a total of 2/3 pint per acre (0.25 lb ae/acre) per year.</li> <li>• Maximum single application rate is 2/3 pint per acre per growing season.</li> <li>• Apply SPUR to non-bearing (well established, 1 year or older) and bearing trees.</li> <li>• East of the Rocky Mountains, do not apply SPUR during bloom.</li> <li>• Avoid direct contact of foliage, fruit, or tree trunks.</li> </ul>

**Broadcast Application:**

Apply to the orchard floor on each side of the apple tree row with a minimum swath width of 3 feet on each side of the tree row. Apply uniformly with ground equipment in a minimum of 10 gallons of water per acre. For Canada thistle, apply after the majority of basal leaves have emerged but prior to bud stage and at least 30 days prior to harvest.

**Tank Mixtures:**

SPUR may be tank mixed with other herbicides labeled for use on apple. Follow all applicable use directions, precautions, restrictions and limitations on the labels for each product used in the tank mix.

MANUFACTURED BY:  
Albaugh, Inc.  
Ankeny, IA 50021

AD021804