

SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1254A-S7 0519

IMPORTANT NOTICE

U.S. LABEL – It is a violation of the United States law to use this product in the United States in a manner inconsistent with its United States labeling.

MANDIPROPAMID	GROUP	40	FUNGICIDE
----------------------	--------------	-----------	------------------

Revus®

Fungicide

This supplemental label expires on 05/01/2022 and must not be used or distributed after this date.

Active Ingredient:

Mandipropamid*	23.3%
----------------------	-------

<i>Other Ingredients:</i>	76.7%
---------------------------------	-------

Total:	100.0%
--------	--------

* CAS No. 374726-62-2

Revus is formulated as a suspension concentrate (SC), containing 23.3% Mandipropamid equivalent to 2.08 pounds per gallon or 250 grams per liter of active ingredient.

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative.

KEEP OUT OF REACH OF CHILDREN.

EPA Reg. No. 100-1254

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Revus as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

SPECIFIC DIRECTIONS FOR USE

Crop	Disease	Rate fl oz/Acre (lb ai/A)	Remarks
Citrus, Crop Group 10-10	<i>Phytophthora</i> brown rot (<i>Phytophthora</i> spp.)	5.5 - 8.0 (0.09 -0.13)	<p>Make first foliar application prior to the onset of disease, and a second application after a minimum interval of 30 days. Do not make more than 2 applications per year.</p> <p>Apply by ground application in a spray volume sufficient to ensure good coverage of the foliage and fruit.</p> <p>Make no more than 2 consecutive applications of Revus before switching to another effective non-Group 40 fungicide.</p> <p>Revus may be tank mixed with a non-Group 40 fungicide labeled for <i>Phytophthora</i> brown rot.</p>

Citrus Crop Group 10-10: Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these

Specific Use Restrictions:

- 1) Do not apply more than 8 fl oz product/A per application.
- 2) Do not apply more than 16 fl oz product/A/year (0.26 lb ai/A/year).
- 3) Revus may be applied the day of harvest (0-day PHI).

Revus and the Syngenta logo are trademarks of a Syngenta Group Company

©2019 Syngenta