

CIDETRAK® CM-OFM COMBO

A MATING DISRUPTANT FORMULATION FOR:

CODLING MOTH (*Cydia pomonella*), **HICKORY SHUCKWORM** (*Cydia caryana*),
ORIENTAL FRUIT MOTH (*Grapholita molesta*), **MACADAMIA NUT BORER**
(*Cryptophlebia ambrodelta*), **KOA SEEDWORM** (*Cryptophlebia illepidia*)

ACTIVE INGREDIENTS:	Percent by Wt.
CM: (8E,10E)-8,10-dodecadien-1-ol.....	4.60 %
OFM: Z-8-Dodecen-1-yl Acetate*.....	1.88%
E-8-Dodecen-1-yl Acetate*.....	0.12%
Z-8-Dodecen-1-ol*.....	0.02%
OTHER INGREDIENTS:	93.38%
TOTAL:	100.00%

NET CONTENTS:

Dispensers per package: 400 units
One dispenser contains: 331 mg, active ingredient

EPA Registration No. 51934-11

EPA Establishment No. 51934-OK-002

MANUFACTURED BY: TRÉCÉ INCORPORATED, 7569 Hwy 28 W · Adair, Oklahoma 74330

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wear chemical resistant gloves and protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID STATEMENT

IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice
IF IN EYES	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER:	Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-866-785-1313 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of product, wash water, or when cleaning equipment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING

CIDETRAK® CM-OFM COMBO disrupts the mating communication between adult male and female moths, reduces fertile egg laying, suppresses subsequent larval infestation, and may reduce the need for applying insecticides.

For maximum effectiveness, growers should remove and destroy infested fruit from trees and the ground. Growers should maintain accurate records from monitoring of CM and OFM adult populations and larval damage within their orchards. Consult your local extension specialist, certified crop advisor, or Trece representative for assistance in designing the best CIDETRAK® CM-OFM COMBO program for your orchard.

Crop	Apple, Pear, Walnut, Pecan, Quince, and other pome fruits; Peaches, Nectarines, Apricots, Plums, Cherries, and other stone fruits; Almonds, Macadamia, and other tree fruit crops.
Targeted Pest	Codling Moth (<i>Cydia pomonella</i>), Hickory Shuckworm (<i>Cydia caryana</i>), Oriental Fruit Moth (<i>Grapholita molesta</i>), Macadamia nut borer (<i>Cryptophlebia ambrodelta</i>), Koa seedworm (<i>Cryptophlebia illepidia</i>).
Rate	Minimum of 200-220 dispensers per acre (500-550 dispensers per hectare) or 46-51 gm CM a.i. and 20-22 gm OFM a.i. per acre per application (115-127 gm CM a.i. and 51-56 gm OFM a.i. per hectare). Maximum of 400-440 dispensers per acre (1000-1100 dispensers per hectare) or 92-101 gm CM a.i. and 40-44 gm OFM a.i. per acre per application (230-253 gm CM a.i. and 101-111 gm OFM a.i. per hectare). Apply the maximum dispenser rate to all orchard borders, to orchards that are moderately to heavily infested with CM and OFM, or to borders of adjacent untreated orchards. Do not exceed 150 grams of each active ingredient per acre per year.
Application	Attach dispensers securely to lateral branches in upper 14" or 35 cm of tree canopy. Attach the dispensers from the ground using a pole applicator or from a moving trailer.
Timing	Apply prior to moth emergence in early spring. Monitor moth activity using PHEROCON® IIB or PHEROCON VI traps and lures. Begin monitoring in early spring and continue throughout the season to assess treatment effectiveness. A second application may be necessary in areas with a long field season (more than 120 days) based on monitoring results, prior to subsequent CM and OFM flights. Consult your fieldman or local extension office for advice on proper timing and retreatment. Note - CIDETRAK® CM-OFM COMBO will not prevent crop damage from immigration of mated females moths into treated orchards. Nearby (500 yards) untreated Codling Moth and Oriental Fruit Moth hosts, such as apple, walnut, pear, quince, crabapple, plum, peaches, nectarines, almonds or other stone fruit and nut crops, can be a source of these mated females. Treatment with CIDETRAK CM-OFM COMBO is recommended only when all host crops within or near treated blocks have been treated with CIDETRAK® CM-OFM COMBO and recommended insecticides.
Treatment Tips	CIDETRAK® CM-OFM COMBO suppresses mating of Codling Moth, Hickory Shuckworm, Oriental Fruit Moth, Macadamia Nut Borer, and Koa Seedworm. Immigration of mated female moths from adjacent, infested orchards can reduce the level of control. Area-wide treatment of the entire host plant block is the most effective strategy. To supplement CIDETRAK® CM-OFM COMBO, use insecticides to control high populations. Monitor all pest populations to determine timely use of insecticides. To manage immigration and high population pressures, consider: 1. Treatment of external sources of infestation with CIDETRAK® CM-OFM COMBO 2. Treatment of external sources of infestation with an insecticide 3. Treatment of pheromone treated orchard with insecticide Consult your local extension specialist, certified crop advisor, or Trece representative for local field condition management strategies.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

PESTICIDE STORAGE: Store unopened original package in a dry location at a temperature below 40° F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Dispose of empty containers and foil envelopes in trash.

WARRANTY AND LIMITATION OF DAMAGES

To the extent consistent with applicable law, all statements concerning the use of this product apply only when used as directed as stated on this product label. Trece Inc. warrants that the product conforms to this label and under normal conditions of use according to this label, is reasonably fit for the stated label purpose. The manufacturer makes no express or implied warranties of merchantability or fitness for a purpose concerning this product, or its use, extending beyond the label. Under no circumstance will Trece Inc. be liable for damage (special, consequential or incidental) that result from the handling, storage, or use of this product which is not in compliance with the label. Read all label directions carefully.

Buyer's exclusive remedy, should this product be defective, is replacement of the product, or if this is impractical, a purchase price refund.



Mailing Address: P.O. Box 129 · Adair, Oklahoma 74330
Shipping Address: 7569 Hwy 28 W. · Adair, Oklahoma 74330
Phone: (918) 785-3061 • Fax: (918) 785-3063

PHEROCON® and CIDETRAK® are registered Trademarks of TRECE Incorporated, Adair, OK USA
©2016 TRECE, Adair, OK USA • TRE0901