THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA



EPA Reg. No. 279-3594

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN ARIZONA & CALIFORNIA

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

Application Rates and Timing: Cotton

When applied as a preplant injection, early post emergence application or early post emergence injection, Topguard® Terra fungicide will provide control of cotton root rot (*Phymatotrichopsis omnivora*) in cotton.

Methods of Application

- Topguard[®] Terra fungicide may be injected into the row up to 30 days prior to planting. Fungicide should be placed 2-3 inches directly below seed placement zone.
- Topguard® Terra fungicide may be applied postemergence (over the top of cotyledon to 2 leaf cotton) as a 4-inch band.
- Topguard® Terra fungicide may be applied to emerged cotton as a directed spray at the base of cotton not larger than 4 true leaves. Directed spray should be concentrated in band no wider than 4 inches at base of plants.
- Topguard® Terra fungicide may be injected into the row after cotton has emerged up to 6 leaf cotton. (or prior to cotton reaching 7-leaf growth stage).

Crop	Pest	Rate (fl. oz/A)	Application Timing
Cotton	Cotton Root Rot (Phymatotrichopsis omnivora)	4 – 8	 Injected into the row up to 30 days prior to planting Postemergence (over the top of cotyledon to 2 leaf cotton) Directed postemergence 2-4 leaf cotton, directed at the base of the plants Injected into the row after cotton has emerged up to 6 leaf cotton

- Use higher rates on irrigated fields or where high infestations of cotton root rot are common.
- Rainfall or irrigation is required to move Topguard® Terra fungicide into the infection zone

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED TOPGUARD TERRA FUNGICIDE LABEL (EPA REG. NO. 279-3594) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION

