Brandt Sequestar 6% Fe EDDHA

F76

Water Soluble EDDHA Chelated Micronutrient

Guaranteed Analysis

Iron (Fe)	6.0%
6.0% Chelated iron	

Derived from ferric EDDHA.

General Information

BRANDT SEQUESTAR 6% Fe EDDHA is a water soluble form of iron EDDHA that is readily available to plants. It can be applied to prevent or correct iron deficiency (chlorosis). This product may be used at any time, but spring applications before the flush of growth or around planting time are recommended to provide iron when the plant's need is greatest.

Directions For Use

Foliar Application: BRANDT SEQUESTAR 6% Fe EDDHA may be applied in water or in combination with most pesticides. Thorough coverage and the use of wetting agents often enhance nutrient uptake from foliar sprays. Before general application, apply the spray solution to a small test area of the crop or foliage to assess any undesirable effects or phytotoxicity. Avoid applying this product during periods of high temperature and low humidity, or when plants are suffering from moisture stress.

Vegetable Crops (including beans, carrots, celery, coel crops, cucumbers, lettuce, melons, onions, peas, peppers, potatoes, spinach, sweet corn, tomatoes):	For ground equipment, apply 1 to 1-1/2 lb per acre in sufficient spray volume for thorough coverage. For aerial applications, apply 1/2 to 1 lb. per acre in at least 10 gal water. Begin	
Field Crops (including alfalfa, corn, dry beans, peas, forage grasses, rice, safflower, sorghum, sugar beets, sugarcane):	4-6 weeks after planting or when deficiency symptoms appear and repeat as needed at 2-3 week intervals until bloom.	
Trees (citrus):	Apply 1/2 to 1 lb per 100 gal of cover spray any time after harvest and until bloom for the succeeding crop. DO NOT apply when fruit is on the tree. DO NOT use in combination with oils or miticides.	
Trees (excluding citrus):	Apply 1/2 to 1 lb per 100 gal of cover spray at 4-6 weeks after full bloom. If deficiency persists, apply 1-2 additional sprays at 2-3 week intervals. DO NOT use in combination with oils or miticides.	
Vine & Berry Crops (including cranberries, grapes, and strawberries):	Apply 1/2 to 1 lb per 100 gal of cover spray.	

Aerial Application: Apply 1/3 lb of BRANDT SEQUESTAR 6% Fe EDDHA per acre in at least 10 gal of water. For application timing and frequency, refer to appropriate foliar application recommendation.

Soil Application: BRANDT SEQUESTAR 6% Fe EDDHA is formulated with EDDHA chelate to improve iron availability to plants, especially those grown in calcareous, alkaline or acid soils. It may be dissolved in water or liquid fertilizers, then applied to the soil as a spray or via drip, sprinkler or furrow irrigation. It may also be dry blended with dry fertilizers or inert carriers (such as sand) then broadcast, banded or sidedressed. For best results, this product should either be incorporated into the root zone at planting, injected into the root zone, or surface applied and moved into the root zone by tillage, drenching, irrigation or rainfall.

	beans, carrots, celery, cole crops, cucumbers, lettuce, melons, onions, peas, peppers, potatoes, spinach, sweet corn, tomatoes):	Band or sidedress 2-1/2 to 5 lb pe acre at planting or when deficienc symptoms appear. Repeat as neede
	Field Crops (including alfalfa, corn, dry beans, peas, forage grasses, rice, safflower, sorghum, sugar beets, sugarcane):	at 2-3 week intervals. Incorporate or irrigate into soil
	Tree Crops (including citrus, fruit and nuts):	Broadcast 1/2 to 2 oz per tree for each inch of trunk diameter. Use lower rates on sandy soils and higher rates on soils with high clay content. DO NOT apply more than 1 lb per tree. Apply annually to the drip line before or at the time of spring growth. For nursery stock, apply 1/3 to 1/2 lb per 300 sq ft. Irrigate into soil.
	Vine And Berry Crops (including cranberries, grapes, and strawberries):	Band or sidedress 1/2 to 1 lb per 100 ft of row in early spring or when deficiency symptoms appear. Incorporate or irrigate into soil.
	Ornamental Shrubs And Trees (including azaleas, gardenias, junipers, pines, roses):	Broadcast 2 oz for small shrubs and up to 2 oz per inch of trunk diameter for large shrubs and trees. Apply annually in the spring before the flush of growth or when deficiency symptoms appear. Irrigate into soil.
	Ornamental Bedding Plants / Groundcovers (including asters, chrysanthemums, geraniums, iris, petunias and verbena):	Broadcast up to 2 oz per 100 sq ft or band 2-4 oz per 100 feet of row. Irrigate into soil.

Vegetable Crops (including

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.html

Information about the components of this lot of fertilizer may be obtained by writing to Brandt Consolidated, Inc. and giving the lot number which is found on the container or label.

Precautionary Statements: Based on currently available data, this product is not classified as a hazardous substance. However, observe good industrial hygiene practices. Wash hands after handling. **Storage and Disposal**: Do not contaminate water, food or feed by storage or disposal. Keep out of reach of children. Store away from incompatible materials. Dispose of contents/container in accordance with local authority requirements.

△ WARNING: Cancer and Reproductive Harm-www.P65Warnings.ca.gov

LIMITED WARRANTY: Brandt Consolidated, Inc. ("BCI") warrants that the Product conforms to the chemical description given on this label and is reasonably fit for the purpose stated when used in accordance with label directions under normal conditions of use. BCI and Seller make no other warranties, express or implied, including warranty of merchantability or fitness for a particular purpose. Buyer and User accept all risks arising from any use of this Product. To the extent allowed by law, BCI and the Seller shall not be liable to the Buyer or User of this Product for any consequential, special, incidental or indirect damages.

