### 5481-452\_Tre-Hold RTU Sprout Inhibitor\_20141015\_17\_5481\_.pdf

# **Tre-Hold® RTU Sprout Inhibitor**

## Ready-to-Use Formulation

Controls Sprouts and Sucker Growth on Apples, Olives, Pears, Non-Bearing Citrus and Nectarines, and Ornamental Woody Plants. For Commercial/Ag Use: Not Residential Use.

#### **ACTIVE INGREDIENT:**

Ethyl 1-Naphthaleneacetate	
INERT INGREDIENTS:	
TOTAL:	100.00%

# KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID				
If on skin or	Take off contaminated clothing.			
clothing:	• Rinse skin immediately with plenty of water for 15-20 minutes.			
	Call a poison control center or doctor for treatment advice.			
If swallowed:	Immediately call a poison control center or doctor.			
	<ul> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> </ul>			
	<ul> <li>Do not give and liquid to the person.</li> </ul>			
	<ul> <li>Do not give anything to an unconscious person.</li> </ul>			
If inhaled:	Move person to fresh air.			
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration,			
	preferably mouth-to-mouth if possible.			
	• Call a poison control center or doctor for further treatment advice.			
If in eyes:	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes			
	• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.			
	• Call a poison control center or doctor for treatment advice.			
	EMERGENCY INFORMATION			
Have the product of	container or label with you when calling a poison control center or doctor, or going for			
treatment.	FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:			
For Medical Emer	gencies phone:1-888-681-4261			
For Transportation	Emergencies, including spill, leak or fire, phone:			
CHEMTREC	1-800-424-9300			
For Product Use In				
AMVAC	1-888-462-6822			

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

EPA Reg. No. 5481-452
EPA Est. No. 5481-CA-01

Net Weight:
As Marked on Container



#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or inhaled. Causes eye and skin irritation. Avoid contact with eyes, skin or clothing.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been heavily drenched or heavily contaminated with this product. Do not reuse them.

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, PVC and viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

#### **User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

#### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply Tre-Hold® RTU Sprout Inhibitor through any type of irrigation system.

Proper pruning of young non-bearing trees promotes the growth and development of strong, well-spaced branches that can support more larger sized fruit, resist wind and ice damage, improve spray penetration for better insect and disease control and improve light penetration for better fruit set, size and color. Non-bearing citrus and nectarine trees sprout vigorously from the root stock and from pruning cuts made on hard wood branches. Tre-Hold® RTU Sprout Inhibitor controls sprouting from pruning cuts, trunks and root stocks. Tre-Hold® RTU Sprout Inhibitor is a ready to use product. For non-bearing nectarines, dilute 1 part to 3 parts water (see dilution chart) and thoroughly treat cut surfaces immediately after pruning to obtain maximum sprout inhibition. Do not allow Tre-Hold® RTU Sprout Inhibitor to contact buds or foliage as injury may result. Mix thoroughly before using.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. These requirements only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product NOT in the scope of the Worker Protection Standard, for agricultural pesticides 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep children, pets and other unprotected persons out of treated areas until sprays have dried.

#### SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

Wind speed restriction:

Do not apply when wind speeds are greater than 10 mph at the application site.

Specific language for orchard air blast:

Sprays must be directed into the crop canopy.

Outward pointing nozzles should be turned off at row ends and when spraying outer rows.

For ground based applications and aerial applications, use only medium or courser spray droplets (or nozzles) according to ASABE (S572) definition for standard nozzles.

#### FOR USE ON APPLES AND PEARS


Do not apply Tre-Hold<sup>®</sup> RTU Sprout Inhibitor to trees less than one year old as injury and stand loss will occur. Do not treat any tree that is not vigorous, healthy, and free from stress. The maximum single application for apples and pears is 1 gallon of product per acre. The maximum annual application for apples and pears is 3 gallons of product per acre. The minimum retreatment interval is 5 days.

#### **New Plantings**

Apple and pear trees, healthy, vigorous and free from stress, may be treated after planting in the grove when the trees show established growth. Remove sprouts and follow the application procedures described below. Do not apply to non-established or stressed trees as injury and stand loss will result.

#### **Established Plantings— Scaffold Limbs**

Prune existing sprouts and treat during the dormant season. Thoroughly cover area where existing sprouts were removed but restrict treatment to the cut surfaces and 2 to 3 inches of the surrounding area. Apply Tre-Hold® RTU Sprout Inhibitor with a brush or small hand held pump-up type sprayer. Use low pressure (not to exceed 25 psi) to avoid fine spray particles from drifting or splashing on surrounding fruiting wood or buds.



Do not treat this area

Do not treat scaffold limbs after bud activity starts in the Spring. There has been some fruit size reduction on Golden Delicious and Anjou pears from excessive treatments to scaffold limbs on older, weaker trees. To minimize the chance of injury, limit treated areas to 10 percent of the total bark area of the tree. Do not treat weak trees, and do not repeat applications on the same tree for at least one year.

#### Trunk and Basal Sprouts and Root Suckers

Prune existing sprouts and apply Tre-Hold<sup>®</sup> RTU Sprout Inhibitor during the dormant season prior to green tip stage or during the Summer pruning season when the new shoots are 6 to 12 inches in height. The later applications have been the most effective. On bearing trees do not treat suckers during bud swell, bloom or fruit set. This period is from the start of growth to 4 weeks after petal fall. Application during this time can result in excessive thinning.

Apply Tre-Hold® RTU Sprout Inhibitor with a brush, small hand held sprayer or back pack sprayer. For large projects, power operated low pressure spray equipment with attached hand gun or trunk directed nozzle can be used. Maintain low spray pressure (not to exceed 25 psi) to avoid spray drift of fine spray particles.

#### **Pruning Cuts on Large Lateral or Upright Limbs**

To control sprouts in the vicinity of pruning cuts, paint the Tre-Hold® RTU Sprout Inhibitor on the bark in the immediate vicinity of the cut. Treat several inches of bark around the cut. Apply with a brush, small hand held sprayer or back pack sprayer. Do not allow the mixture to splash or drip onto other parts of the tree. Treat only in the dormant season prior to bud activity. Spray application is not recommended, due to the possibility of spray contacting fruit buds.

#### OLIVES

The maximum single application for olives is 3 gallons of product per acre.

The maximum annual application for olives is 3 gallons of product per acre.

The minimum retreatment interval is 5 days.

#### **Trunk Basal Sprouts and Root Suckers**

(Do not treat scaffold branches or pruning cuts on olive trees.) Prune off existing sprouts or suckers and allow new sprouts to regrow. Treat with Tre-Hold® RTU Sprout Inhibitor when sprouts are not more than 10 inches tall. Apply with a brush, small hand held sprayer or back pack sprayer. Tre-Hold® RTU Sprout Inhibitor will control new sprout growth of olives, but will not control old woody sprouts. Do not treat during olive bloom or fruit set. Limit treated areas to 10% of the plant's total bark surface.

#### ORNAMENTAL WOODY PLANTS

The maximum single application for ornamental woody plants is 1 gallon of product per acre.

The maximum annual application for ornamental woody plants is 3 gallons of product per acre.

The minimum retreatment interval is 5 days.

#### **Controls Regrowth of Tree Sprouts Following Proper Pruning**

Tre-Hold<sup>®</sup> RTU Sprout Inhibitor prevents and inhibits vegetative growth in woody plants. After proper pruning, allow new sprouts to regrow and apply Tre-Hold<sup>®</sup> RTU Sprout Inhibitor with a brush, a small hand held sprayer or back pack sprayer when sprouts are no more than 10 inches tall. It is effective on many ornamental, flowering and shade trees such as:

American Elm	Crabapples	Red Maple	Sycamore
Carob	Crepe Myrtle	River Birch	Water Oak
Chinese Elm	Flowering Plum	Silver Maple	Wax Myrtle

Coast Redwood Pyracanthas Sugar Maple Willow spp. (Black; Marsh)

For most effective control of resprouts, prune to a lateral branch (drop crotch pruning), and apply Tre-Hold<sup>®</sup> RTU Sprout Inhibitor after resprout growth has begun. Do not leave stubs and avoid severe trimming. Treatment of all cuts is important.

#### **NON-BEARING CITRUS**

Use the product as supplied (undiluted) and follow the procedures described below for treating sprouts on scaffold limbs, trunks and rootstocks.

#### NON-BEARING NECTARINES

For non-bearing nectarines, first dilute the product to 1/4 the standard rate as shown in the following examples and follow the procedures described below for treating sprouts on scaffold limbs, trunks and rootstocks.

PRODUCT DILUTION FOR NON-BEARING NECTARINES				
Amount of Tre-Hold® RTU Sprout Inhibitor	Amount of Water	Total Volume		
1 Quart	3 Quarts	1 Gallon		
1 Gallon	3 Gallons	4 Gallons		

This product is not for use on non-bearing nectarines in California.

#### APPLICATION TO NON-BEARING CITRUS AND NECTARINES

#### **Sprouting from Scaffold Limbs**

Remove unwanted sprouts and thoroughly coat cut surfaces and surrounding bark.

#### **Sprouting from Trunk**

Remove trunk suckers and water sprouts and apply Tre-Hold® RTU Sprout Inhibitor to the entire trunk from ground line to scaffold limbs

#### Rootstock

Non-bearing citrus rootstock which is healthy, vigorous and free from stress may be treated. After planting, apply Tre-Hold<sup>®</sup> RTU Sprout Inhibitor from the ground line to the first scaffold branches. Do not allow the product to puddle at the base of the trees as injury and stand loss may occur. Do not apply to rootstock if the bud union has not callused. In temperate zones where freezing may injure scaffold branches, restrict treatment to the frost-free period to allow for a short period of growth between treatment and damaging freeze.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not store near fertilizers, seeds, insecticides or fungicides.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container.

Non refillable container equal or less than 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

Tre-Hold<sup>®</sup> is a registered trademark of Amvac Chemical Corporation.

Amvac Chemical Corporation 4100 E. Washington Blvd.

Los Angeles, CA 90023 U.S.A.