



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)

PROPULSE™

For control of Early Leaf Spot, Late Leaf Spot, Rust, and Nematodes in Peanuts
 EPA Reg. No. 264-1084

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ALABAMA, ARKANSAS, FLORIDA, GEORGIA, MISSISSIPPI, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TEXAS, VIRGINIA

This recommendation is valid until December 31, 2020

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Peanut		
Pest Control	Application Rate	Product Instructions
Early Leaf Spot (<i>Cercospora arachidicola</i>) Late Leaf Spot (<i>Cercosporidium personatum</i>) Rust (<i>Puccinia</i> spp.)	13.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.
Nematodes	13.6 fl oz/acre	For nematode suppression, either apply PROPULSE by chemigation using overhead irrigation equipment using 0.10 to 0.25 acre-inch of water or by foliar spray using a minimum of 15 gallons of water per acre. Foliar applications should be immediately followed by overhead irrigation using 0.10 to 0.25 acre-inch of water the move PROPULSE into the soil. For maximum nematode suppression, PROPULSE should be applied approximately 45 days after planting and following an at-plant nematicide application.

Restrictions:

- Do not apply more than 34.2 fl oz/acre per year.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs fluopyram or 0.712 lbs prothioconazole per acre per year, including seed treatment, soil and foliar uses.
- Apply by ground, aerial, or chemigation application equipment.
- Do not apply PROPULSE within 14 days of harvest.
- Do not feed hay or threshings or allow livestock to graze in treated areas.
- To limit the potential for development of disease resistance to this fungicide, do not make more than 2 sequential applications of PROPULSE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).