



ITERA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF LOUISIANA

Quilt Xcel® Fungicide

For the Control of Red Rot (*Glomerella tucumanensis*, *Colletotrichum falcatum*) and
Pineapple Disease (*Ceratosystis paradoxa*) in Sugarcane

EPA Reg. No. 100-1324

EPA SLN No. LA19-0004

This label expires and must not be distributed or used in accordance with this SLN
registration after July 29, 2024

KEEP OUT OF REACH OF CHILDREN
CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Restrictions:

- Quilt Xcel Fungicide may be applied **either** as an In-Furrow Application, **or** a Conveyor Belt Treatment.
- Regardless of which application method used, **DO NOT** apply more than 21 fl oz/A of Quilt Xcel Fungicide at planting.

In-Furrow Application

Apply Quilt Xcel Fungicide to sugarcane seed pieces as an in-furrow spray at planting at the rate of 1.4 – 3.0 fl oz per 1000 row feet using sufficient water to ensure coverage. Mount the spray nozzle so the spray is directed into the furrow just before the seed cane is covered. Use the higher rate when the weather conditions are expected to be conducive for disease development.

Conveyer Belt Treatment (Attached to Planting Equipment)

Treat seed pieces with Quilt Xcel Fungicide spray solution at the rate of 1.4 – 3.0 fl oz per 1000 row feet, using in-line directed spray sufficient to wet cut ends.

©2019 Syngenta

Quilt® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: LA1324037AA0719