



FIFRA Section 24(c) Special Local Need Label

Praiz™

Flowable Agricultural Fungicide

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NORTH DAKOTA FOR CONTROL OF DISEASES ON LONG-SEASON POTATOES GROWN UNDER IRRIGATION

This labeling must not be distributed or used after January 1, 2019.

ACTIVE INGREDIENT:

Chlorothalonil (tetrachloroisophthalonitrile)54.0%

OTHER INGREDIENTS:46.0%

TOTAL:100.00%

Contains 6 pounds chlorothalonil per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand this label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label and the Federal label for this product must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Praiz and this Section 24(c) label before applying. Do not use in a manner inconsistent with applicable use directions, restrictions, precautions and limitations provided on the label affixed to the container for Praiz and this Section 24(c) label.

Crop	Diseases	Causal Agent	Application Rate Pints/Acre (lbs. a.i./acre)
Potato (Long-season varieties grown under irrigation)	Late blight	<i>Phytophthora infestans</i>	Low rate: ¾ pint (0.6)
	Early blight	<i>Alternaria solani</i>	-then-
	Botrytis vine rot	<i>B. cinerea</i>	Higher rates: 1 to 1 ½ pints (0.75 to 1.125)

Application Directions: Begin applications of Praiz at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals. The minimum retreatment interval is 5 days.

Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur:

- Vines close between rows;
- Late blight forecasting measures 18 disease severity values (DSV); or
- The crop reaches 300 P-days.

Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.



This product may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). Do not exceed a 10 day interval between applications when using this technique. See *Application and Calibration Techniques for Sprinkler Irrigation* on the Praiz label.

Specific Use Restrictions: Do not apply more than 21 ½ pints (16 lb a.i.) of Praiz per acre on long-season varieties of potato during each growing season. Do not apply within 7 days of harvest.

EPA Reg. No. 9779-320

EPA SLN No. ND-160001

Distributed By:

Winfield Solutions, LLC

P.O. Box 64589, St. Paul, MN 55164-0589

1/0510/6