

## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL AFFIXED TO THE CONTAINER FOR WARRANT® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

Use of Warrant Herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container and in this supplemental labeling for Warrant Herbicide.

It is a violation of federal law to use this product in any manner inconsistent with its labeling. The labeling must be in the possession of the user at the time of herbicide application. This supplemental label expires on December 31, 2018 and must not be used or distributed after this date.

ACETOCHLOR	GROUP	15	HERBICIDE
------------	-------	----	-----------



EPA Reg. No. 524-591

WARRANT® and Rowel® are registered trademarks of Monsanto Technology LLC.

---

### Preplant, At-Planting, Preemergence and Postemergence Applications in Peanut

---

Keep Out of Reach of Children.

### CAUTION!

In case of an emergency involving this product, call collect, day or night, 314-694-4000.

---

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

See the "Product Information" and "Mixing, Spraying, and Handling Instructions" sections of the label booklet for Warrant Herbicide for essential product performance information.

**Read the "Limit of Warranty and Liability" in the label booklet for Warrant Herbicide before using. These terms apply to this supplemental label and if these terms are not acceptable, return the product unopened at once.**

This product, when applied preplant, at-planting, preemergence, or postemergence in peanut will provide preemergence control or reduced competition of the annual weeds listed in the "WEEDS CONTROLLED" section of this label. If weeds are emerged at the time of application, apply a labeled foliar herbicide with this product to control the emerged weeds. Observe the directions for use, precautions, and restrictions on the label of the foliar herbicide.

Application of Warrant Herbicide followed by conditions that do not foster adequate crop growth or cause stress (cold, wet soils), waterlogged conditions, excessive irrigation or rainfall, may result in crop injury.

Application made before weeds emerge, or after clean cultivation, is necessary, as this product will not control emerged weeds.

DO NOT exceed a total of 4 quarts (3 lbs acetochlor) per acre per season when making multiple applications.

Do not exceed a total of 3 applications per season.

Allow a minimum of 90 days between last application and grazing or feeding of peanut hay to livestock.

#### Approved Application Systems

**Ground:** Broadcast boom

**Aerial:** Fixed-wing and helicopter. **Allowed in selected states only** - see Aerial Application Equipment section of the label booklet for additional information.

## Approved Application Methods

### Preplant; At-planting; Preemergence

Warrant Herbicide may be applied preplant, at-planting, or preemergence to peanut at 1.25 to 2 quarts (0.94 to 1.5 lbs a.i.) per acre. Apply broadcast to the soil surface according to the rate table listed below.

**Tank Mixtures:** It is the applicator's responsibility to ensure that all products used in a tank mixture with this product are registered for the specific preplant or preemergence application timing in peanut. Read and follow all applicable restrictions and limitations and directions for use involving tank mixing on all product labels included in the tank mixture, including any applicable crop injury precautions. The end-user must follow the most restrictive directions for use and precautionary statements on the labeling of each product in the tank mixture.

This product may be applied early preplant or preemergence to peanut together with the following products in order to expand the weed control spectrum:

pendimethalin, ethalfuralin, diclosulam, trifluralin Prowl®, Sonalan®, Strongarm®, Treflan®, Rowel, Valor

Preplant **soil incorporated** applications together with Prowl, Sonalan, Strongarm or Treflan are not recommended due to risk of crop injury and reduced weed control.

Application of Warrant Herbicide in tank mixture with other products or to soils where other applications of soil applied herbicides have been made may increase the potential for injury with this product.

#### Application Rates (minimum and maximum range):

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE*					
	Less than 1.5% Organic Matter (quarts)			1.5% or More Organic Matter (quarts)		
Coarse	1.25	to	1.6	1.25	to	1.7
Medium	1.25	to	1.7	1.25	to	1.9
Fine	1.25	to	1.9	1.25	to	2.0

\* Use the higher rate in the range for areas of heavy weed infestation.

### Postemergence Surface

Warrant Herbicide may be applied postemergence to peanut at 1.25 to 2 quarts per acre after crop emergence up through the R1 growth stage (beginning bloom). R1 ends as 50% of the plants in an area have a visible peg (R2). Apply broadcast over top of the crop or directed to the soil surface according to the rate table listed below. Do not exceed 2 quarts (1.5 lbs a.i.) per acre as a single application. Allow at least 7 days between sequential applications.

**Tanks Mixtures:** It is the applicator's responsibility to ensure that all products used in a tank mixture with this product are registered for the specific postemergence application timing in peanut. Read and follow all applicable restrictions and limitations and directions for use involving tank mixing on all product labels included in the tank mixture, including any applicable crop injury precautions. The end-user must follow the most restrictive directions for use and precautionary statements on the labeling of each product in the tank mixture.

This product may be applied postemergence to peanut together with the following products to expand weed control spectrum or for control of emerged weeds at the time of application:

2,4-DB, flumioxazin, paraquat, imazapic, lactofen, bentazon, acifluofen

Gramoxone®, Firestorm®, Parazone®, Cadre®, Cobra®, Storm®, Ultra Blazer®

Application of Warrant Herbicide in tank mixture with other products or to soils where other applications of soil applied herbicides have been made may increase the potential for injury with this product.

#### Application Rates (minimum and maximum range):

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE*					
	Less than 1.5% Organic Matter (quarts)			1.5% or More Organic Matter (quarts)		
Coarse	1.25	to	1.6	1.25	to	1.7
Medium	1.25	to	1.7	1.25	to	1.9
Fine	1.25	to	1.9	1.25	to	2.0

\* Use the higher rate in the range for areas of heavy weed infestation.

All other trademarks are the property of their respective owners.

©2018 MONSANTO COMPANY  
MONSANTO COMPANY  
800 NORTH LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167 USA

36067S1-46