



MAKHTESHIM
A G of A N
NORTH AMERICA

FIFRA §2(ee) Recommendation

Bumper® 41.8 EC Fungicide

EPA Reg. No. 66222-42

For control of Rust and Scab on
 Almonds in California only

DIRECTIONS FOR USE

- The user must refer to the federally approved Bumper 41.8 EC Fungicide label and read and follow all directions for use, restrictions, and precautions.
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Users should have this recommendation in their possession at the time of use.
- This recommendation is permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the EPA.

CROP	PESTS CONTROLLED	RATE OF APPLICATION PER ACRE	APPLICATION INSTRUCTIONS
ALMONDS	Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>)	6.4 to 8 fl oz	Apply 6.4 to 8 fl oz (0.18 -0.225 lbs. active) per acre. Begin application approximately 2-5 weeks after petal fall when environmental conditions are favorable for disease development. For optimum disease control, apply on a 7- to 14-day application schedule. Make no more than two consecutive applications before rotating to a fungicide with a different mode of action (not a Group 3).
	<ul style="list-style-type: none"> • Do not apply more than 32 fl. oz. per acre per season of Bumper 41.8EC. • Do not apply more than 0.90 lb. a.i. propiconazole per acre per season. • Do not apply within 60 days of harvest. • Do not graze livestock in treated areas or cut treated cover crop for feed. 		

Bumper® is a registered trademark of Makhteshim Agan of North America, Inc.

Effective Date: 3-3-2010

Makhteshim Agan of North America Inc.
3120 Highwoods Blvd., Suite 100 •Raleigh, NC 27604