



TREATED SEED

This seed has been treated with INTEGO™ SUITE Soybeans which contains clothianidin, ethaboxam, ipconazole and metalaxyl at the rates recommended by the manufacturer.

DO NOT USE TREATED SEED FOR FEED, FOOD OR OIL PROCESSING

STORE AWAY FROM FEEDS AND OTHER FOODSTUFFS

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice.
Have a person sip a glass of water if able to swallow.
Do not induce vomiting unless told to by a poison control center or doctor.
Do not give anything to an unconscious person.

NOTE TO PHYSICIAN

Treatment is supportive and symptomatic.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

When handling treated seed wear long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves: barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride or Viton.

- After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. *Exception:* Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction.
- Regardless of application method, do not apply more than 0.2 lb (90.7 g) of clothianidin active ingredient per acre per year. (Based on the seed size of 2800 seeds/lb, at the label rate, the mg of clothianidin will be equal to 0.081 mg/seed).



TREATED SEED

This seed has been treated with INTEGO™ SUITE Soybeans which contains clothianidin, ethaboxam, ipconazole and metalaxyl at the rates recommended by the manufacturer.

DO NOT USE TREATED SEED FOR FEED, FOOD OR OIL PROCESSING

STORE AWAY FROM FEEDS AND OTHER FOODSTUFFS

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice.
Have a person sip a glass of water if able to swallow.
Do not induce vomiting unless told to by a poison control center or doctor.
Do not give anything to an unconscious person.

NOTE TO PHYSICIAN

Treatment is supportive and symptomatic.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

When handling treated seed wear long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves: barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride or Viton.

- After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. *Exception:* Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction.
- Regardless of application method, do not apply more than 0.2 lb (90.7 g) of clothianidin active ingredient per acre per year. (Based on the seed size of 2800 seeds/lb, at the label rate, the mg of clothianidin will be equal to 0.081 mg/seed).

- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Clothianidin is toxic to birds and mammals. Seeds treated with clothianidin exposed on the soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate water when disposing of equipment washwater or rinseate.
- Treated seeds must be planted at a minimum depth of 1 inch. This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- This product contains metolaxyl which is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Do not graze or feed soybean forage and hay to livestock.
- Areas planted with treated soybean seed may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- Do not re-use bags from treated seed to handle food or feed products.
- Do not use augers used for handling treated seed to move seed for feed, food or oil processing.
- Foliar spray of clothianidin can be applied 45 days after planting *NTEGO* SUITE Soybeans treated soybean seed for mid- to late-season foliar insects.
- Not for use on soybeans in California.
- Iproconazole may persist in soil, and long term use in the same field may result in accumulation of active ingredient.

EPA Reg. 59639-205

059639-20150717

NTEGO is a trademark of Valent U.S.A. Corporation

For **PRODUCT USE** information: Call 800-6-VALENT (800-682-5368)

IN CASE OF EMERGENCY ONLY: Call 24 Hours A Day 800-892-0099

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek CA 94596-8025

Form 2034-B

- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Clothianidin is toxic to birds and mammals. Seeds treated with clothianidin exposed on the soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate water when disposing of equipment washwater or rinseate.
- Treated seeds must be planted at a minimum depth of 1 inch. This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- This product contains metolaxyl which is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Do not graze or feed soybean forage and hay to livestock.
- Areas planted with treated soybean seed may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- Do not re-use bags from treated seed to handle food or feed products.
- Do not use augers used for handling treated seed to move seed for feed, food or oil processing.
- Foliar spray of clothianidin can be applied 45 days after planting *NTEGO* SUITE Soybeans treated soybean seed for mid- to late-season foliar insects.
- Not for use on soybeans in California.
- Iproconazole may persist in soil, and long term use in the same field may result in accumulation of active ingredient.

EPA Reg. 59639-205

059639-20150717

NTEGO is a trademark of Valent U.S.A. Corporation

For **PRODUCT USE** information: Call 800-6-VALENT (800-682-5368)

IN CASE OF EMERGENCY ONLY: Call 24 Hours A Day 800-892-0099

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek CA 94596-8025

Form 2034-B