

KPHITE® 7LP

SYSTEMIC FUNGICIDE BACTERICIDE

EPA REG. NO. 73806-1

FIFRA 2(ee) RECOMMENDATION FOR USE IN CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency.

CITRUS (Lemons)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered labels.

GENERAL APPLICATION INSTRUCTIONS

CROP	DISEASE	APPLICATION METHOD	RATE OF APPLICATION	APPLICATION PROGRAM
CITRUS (Lemons))	<i>Hyphoderma sambuci</i> ; symptom (<i>Hyphoderma gummosis</i>)	Foliar	2-3 quarts in a minimum of 100 gallons of water per acre.	<p>Apply K-PHITE 7LP at 4-6 week intervals.</p> <p>Do not apply K-PHITE 7LP at intervals of less than three days.</p> <p>Provide adequate coverage to reach the scaffolds, crotch, and trunk parts of the tree.</p> <p>Supplement and promote new growth and vigor through tank mixing RENEW 4-20-22 Urea, Di-Potassium Poly Phosphate / Phosphite Combination at rates of 2-4 gallons per acre in combination with K-PHITE 7LP.</p>
<p>*Make applications prior to disease development in conjunction with good cultural management practices.</p> <p>*Use the higher rate when disease pressure is severe.</p>				

All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. This FIFRA Section 2(ee) recommendation contains new or additional Diseases, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product label.

REGISTRANT : PLANT FOOD SYTEMS, INC

P.O. BOX 775, 2827 Union Street, Zellwood, FL. 32798

