



4.50"

**RESTRICTED USE PESTICIDE**  
 May injure (phytotoxic) susceptible non-target plants.  
 For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial and certified applicators must ensure that all persons involved in these activities are informed of the precautionary statements.






**CORVUS<sup>®</sup>**

Net Contents:

**2.5 Gallons**

Herbicide

**GROUP 2 | 27 HERBICIDE**



*For: weed control in field corn, seed corn and corn grown for silage in the states of: AR, AL, CO, DE, GA, IL, IN, IA, KS, KY, LA, MI, MN, MO, MS, MT, NE, NJ, NM, NC, ND, OH, OK, PA, SC, SD, TN, TX, VA, WI and WY.*

*In the states of CO, DE, KS, MD, MO, NJ, NM, SD, and WV use is only allowed under 24c registrations. A current 24c label must be in the possession of the user at the time of the pesticide application.*

*In the state of MN use is only allowed in accordance with the Minnesota Product Bulletin.*

*In the state of WI use is only allowed in accordance with the Wisconsin Product Bulletin.*

<b>ACTIVE INGREDIENTS:</b> Thiencarbazone-methyl: (Methyl 4-[[[(4,5-dihydro-3-methoxy-4-methyl-5-oxo-1H-1,2,4-triazol-1-yl)carbonyl]amino]sulfonyl]-5-methyl-3-thiophenecarboxylate) . . . . .	7.60%
Isoxaflutole [5-cyclopropyl-4-(2-methylsulfonyl-4-trifluoromethylbenzoyl) isoxazole] . . . . .	19.00%
<b>OTHER INGREDIENTS:</b> . . . . .	73.40%
<b>TOTAL</b>	<b>100.00%</b>

Contains 0.75 pounds Thiencarbazone-methyl per U.S. gallon  
 Contains 1.88 pounds Isoxaflutole per U.S. gallon

**EPA Reg. No. 264-1066**

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours a Day 1-800-334-7577  
 For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

**See additional precautionary statements and directions for use on label.**

Produced for:  
 Bayer CropScience LP  
 P.O. Box 12014, 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 CORVUS is a registered trademark of Bayer.  
 ©2016 Bayer CropScience

COPY AREA 7.50"

7.75"

4.25"

COPY AREA

4.50"

### FIRST AID

<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor for treatment advice.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a physician if irritation persists.</li> </ul>
<b>If on Skin:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p><b>In case of emergency, call the toll-free Bayer CropScience Emergency Response telephone number: 1-800-334-7577.</b></p> <p><b>Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</b></p>	
<p>Note to Physician: No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.</p>	

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

##### CAUTION

- Harmful if swallowed or absorbed through the skin.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.

##### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below.

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks
- Protective eye wear.

When mixing/loading or cleaning equipment, wear a chemical resistant apron in addition to the other required PPE. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

##### ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

##### USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

COPY AREA

7.375"

7.75"

4.0625"

COPY AREA

4.50"

**USER SAFETY RECOMMENDATIONS**

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Drift or runoff may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organism in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Do not apply when weather conditions favor drift from treated areas. Do not use the same spray equipment for other purposes unless thoroughly cleaned. Do not contaminate water used for irrigation or domestic purposes.

CORVUS Herbicide contains isoxaflutole which is known to leach through soil into shallow ground water under certain conditions as a result of agricultural use. Use of CORVUS Herbicide in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

This product can contaminate surface water through spray drift. Under some conditions, product residues may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips and areas over-laying tile drainage systems that drain to surface water.

A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of these chemicals from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

In fields having sands, loamy sands and sandy loam soils, special care should be taken not to over-irrigate since substantial over-irrigation promotes the leaching of chemicals.

This pesticide is toxic to some plants at very low concentrations. Non-target plants may be adversely affected if the pesticide is allowed to drift from areas of application. Exposure to isoxaflutole residues may injure or kill susceptible plants. Symptoms of phytotoxicity as a result of exposure to isoxaflutole include whitening or chlorosis of the foliage of affected plants. Cotton is particularly susceptible to isoxaflutole; therefore, exposure of cotton to isoxaflutole residues may affect cotton yield. To prevent damage to crops and other desirable plants, read and follow all directions and precautions on this label before using.

The chemicals in this product have properties and characteristics associated with chemicals detected in ground water. These chemicals may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff, according to the pesticide's mean soil partition coefficient (Kd) for several days after application.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

*(continued)*

4.0625"

COPY AREA

COPY AREA

7.375"

7.75"

4.50"

COPY AREA

7.375"

Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to exclude completely precipitation from contact shall be of sufficient capacity to contain at a minimum of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

**Endangered Species Advisory/Protection Requirements**

This product may have effects on federally listed threatened or endangered species or their critical habitat in some locations. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

(continued)

4.0625"

COPY AREA

7.75"

4.50"

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

**DIRECTIONS FOR USE**

**RESTRICTED USE PESTICIDE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**Read the entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**For Important crop safety information, refer to the Use Directions section under the specific crop.**

**In Minnesota, this product must only be used in accordance with the Minnesota Product Bulletin. The Minnesota Product Bulletin, which accompanies the sale and packaging of the product, must be in possession of the user at the time of pesticide application.**

**In Wisconsin, this product must only be used in accordance with the Wisconsin Product Bulletin. The Wisconsin Product Bulletin, which accompanies the sale and packaging of the product, must be in possession of the user at the time of pesticide application.**

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

**PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:**

- Coveralls over
- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Socks plus chemical resistant footwear
- Protective eye wear

**PRODUCT INFORMATION**

CORVUS Herbicide:

- is a selective herbicide for the control of important broadleaf and grass weeds in field corn, seed corn, corn grown for silage.

*(continued)*

COPY AREA

7.375"

7.75"

4.0625"

COPY AREA

4.50"

COPY AREA

7.375"

- is formulated as a suspension concentrate containing 2.63 pounds of active ingredients per gallon [0.75 lbs Thiencazuron-methyl a.i., 1.88 lbs Isoxaflutole a.i.].
- has multiple modes of actions: the first, inhibiting of enzymes that are essential to the protection of chlorophyll in plant leaves, and a second blocking the plant's synthesis of certain amino acids/protein synthesis.
- is effective in controlling glyphosate-, triazine-, PPO-, ALS- and auxin- herbicide resistant populations of weed species.

**APPLICATION INSTRUCTIONS**

CORVUS Herbicide:

- may be used in either conventional, conservation tillage, or no-till crop management systems.
- may be applied preplant [surface-applied or incorporated (less than 2" deep)], preemergence or early postemergence.
- will provide its most effective weed control when applied and subsequently moved into the soil by rainfall, sprinkler irrigation or mechanical tillage prior to weed emergence.
- may be tank mixed or applied in sequential applications with other herbicides to control additional weeds
- may be applied using either water or sprayable grade fluid fertilizer as a liquid carrier.
- may be applied by ground application only. Aerial application is not permitted.
- may be applied as either a broadcast spray or as a band application.

Refer to the 'Specific Use Directions' section of the label for application timing information specific from each registered use of CORVUS Herbicide.

**Ground Application (Banding)**

Banding application equipment must be carefully calibrated to prevent crop exposure to concentrations of CORVUS Herbicide that exceed the labeled rate for the soil type. It is critical to insure that the calibrated band width equates to actual band width realized in field applications. Bands actually delivered at a width narrower than targeted will concentrate the product and increase the risk for crop response.

Even flat spray tip nozzles and a band width of no less than 12" must be used.

Apply a broadcast equivalent rate and volume per acre. The following equations may be used to make the required calculations as follows:

$$\frac{\text{band width (inches)}}{\text{row width (inches)}} \times \text{broadcast rate per acre} = \text{banding rate per acre}$$

$$\frac{\text{band width (inches)}}{\text{row width (inches)}} \times \text{broadcast spray volume per acre} = \text{banding spray volume per acre}$$

**Ground Application (Broadcast)**

Apply CORVUS Herbicide either alone or in tank mixtures in a minimum of 10 gallons of spray mixture per acre. Uniform, thorough spray coverage is important to achieve consistent weed control. Keep the spray boom at the lowest possible spray height above the target surface. Refer to the nozzle manufacturer's recommendations for proper nozzle, pressure setting and sprayer speed for optimum product performance and minimal spray drift. Uneven application, sprayers not properly calibrated, or improper incorporation may decrease the level of weed control and/or increase the level of adverse crop response. Maintain a constant ground speed while applying this product to ensure proper distribution. Do not overlap spray patterns beyond equipment manufacturers recommendations as excessive rates may result in adverse crop responses and potential stand loss. Maintain adequate agitation at all times, including momentary stops.

4.0625"

COPY AREA

7.75"

4.50"

COPY AREA

7.375"

**USE RESTRICTIONS**

- Use on coarse textured soils with a shallow water table – All Registered Uses:
  - In the states of AL, AR, CO, DE, GA, KS, KY, LA, MD, MO, MS, NC, NM, OK, SC TN, TX, VA, and WV if the water table (i.e, level of saturation) is less than 25 feet below the ground surface, do not use on soils meeting all three of the following criteria. If the water table depth is unknown, do not use on any of the soils meeting all three of the following criteria. If less than three criteria are met or the water table is greater than 25 feet below the ground surface, there is no restriction against application:
    - The surface soil texture is loamy sand or sand
    - The subsoil texture is loamy sand or sand
    - The average organic matter (in the upper 12 inches) is less than 2% by weight
  - In the states of IA, IL, IN, MI, MT, ND, NE, NJ, OH, PA, SD, and WY, if the water table (i.e, level of saturation) is less than 25 feet below the ground surface, do not use on soils meeting all three of the following criteria. If the water table depth is unknown, do not use on any of the soils meeting all three of the following criteria. If less than three criteria are met or the water table is greater than 25 feet below the ground surface, there is no restriction against application:
    - The surface soil texture is sandy loam, loamy sand or sand
    - The subsoil texture is loamy sand or sand
    - The average organic matter (in the upper 12 inches) is less than 2% by weight
- Do not apply more than 5.6 fluid oz of CORVUS Herbicide per 365 day period or exceed the maximum labeled rate for any given soil type.
- Do not apply this product using aerial application equipment.
- Do not apply this product through any type of irrigation system.
- Do not use flood or furrow irrigation to apply, activate or incorporate this product.
- Do not allow cover crops in fields treated with CORVUS Herbicide to be grazed by livestock or harvested for food.
- To prevent off-site movement of soil containing this product to non-target areas, do not apply CORVUS Herbicide to areas receiving less than 15 inches of average annual precipitation unless supplemented to at least the equivalent of 15 inches of annual precipitation with irrigation water.
- In Minnesota, this product must only be used in accordance with the Minnesota Product Bulletin. The Minnesota Product Bulletin, which accompanies the sale and packaging of the product, must be in possession of the user at the time of pesticide application.
- In Wisconsin, this product must only be used in accordance with the Wisconsin Product Bulletin. The Wisconsin Product Bulletin, which accompanies the sale and packaging of the product, must be in possession of the user at the time of pesticide application.

Refer to the specific use directions and restrictions in each specific crop section.

**USE PRECAUTIONS**

- Application of CORVUS Herbicide at less than specified rates for the appropriate soil will only provide suppression of sensitive weeds.

**RESISTANCE MANAGEMENT**

Corvus Herbicide contains Group 27 & 2 Herbicides, i.e., an HPPD inhibitor (Group 27) and ALS/AHAS enzyme inhibitor (Group 2). A given weed population may contain or develop resistance to a herbicide after repeated use. Appropriate resistance-management strategies should be followed to mitigate or delay resistance. The following Integrated Weed Management Techniques are effective in reducing problems with herbicide resistant weed biotypes. It is best to use multiple practices to manage or delay resistance, as no single strategy is likely to be totally effective.

(continued)

4.0625"  
COPY AREA

7.75"

4.50"

COPY AREA

7.375"

- **Rotate crops.** Crop rotation diversifies weed management.
- **Rotate herbicide-tolerant traits.** Alternate herbicide-tolerant (HT) traits and/or use HT trait stacks for more efficient rotation.
- **Use multiple herbicide sites of action.** Use tankmix partners and multiple SOAs during both the growing season and from year to year to reduce the selection pressure of a single SOA.
- **Know your weeds, know your fields.** Closely monitor problematic areas with difficult-to-control weeds or dense weed populations.
- **Start with clean fields.** Effective tillage or the use of a burndown herbicide program can control emerged weeds prior to planting.
- **Stay clean – use residual herbicides.** Regardless of tillage system, pre-emergence or early post-emergence soil-applied residual herbicides should be used when possible.
- **Apply herbicides correctly.** Ensure proper application, including timing, full use-rates and appropriate spray volumes.
- **Control weed escapes.** Consider spot herbicide applications, row wicking, cultivation or hand removal of weeds or other techniques to stop weed seed production and improve weed management.
- **Zero tolerance – reduce the seed bank.** Do not allow surviving weeds to set seed, which will help decrease weed populations from year to year and prevent major weed shifts.
- **Clean equipment.** Prevent the spread of herbicide-resistant weeds and their seeds.

Contact your local extension specialist, certified crop advisory and /or Bayer CropScience representative for additional resistance management or IPM recommendation. Also for more information on Weed Resistance Management, visit the Herbicide Resistance Action Committee (HRAC) on the web at <http://www.hracglobal.com>.

## SPRAY DRIFT MANAGEMENT

### Droplet Size

To obtain uniform spray coverage and reduce spray drift, equip sprayers with nozzles that provide medium to coarse spray droplets as defined in ASABE Standard S572.1.

### Wind Speed

Only apply this product when the potential for drift to adjacent non-target areas is minimal (e.g., when the wind is 10 MPH or less and is blowing away from sensitive areas).

To avoid potential adverse effects to non-target areas, maintain a 25 foot buffer between the point of direct application and the **closest downwind edge** of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrub lands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

## COMPATIBILITY TESTING AND TANK MIX PARTNERS

### Compatibility

If CORVUS Herbicide is to be tank mixed with liquid fertilizers or other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5-15 minutes after mixing. Read and follow all parts of the label of each tank-mix product.

7

4.0625"

COPY AREA

7.75"



4.50"

COPY AREA

7.375"

**Order of Mixing**

CORVUS Herbicide may be used with other recommended pesticides, fertilizers, and micronutrients.

The proper mixing procedure for CORVUS Herbicide application with water or liquid fertilizer as a carrier:

1. Fill the spray tank 1/4 to 1/2 of the required volume of water or liquid fertilizer prior to the addition of CORVUS Herbicide.
2. Add the proper amount of CORVUS Herbicide, then add the rest of the water or liquid fertilizer to the desired level.
3. Maintain sufficient agitation to ensure a uniform spray mixture during application.
4. If CORVUS Herbicide is applied in a tank mixture with other pesticides, add CORVUS Herbicide to the spray tank first and ensure it is thoroughly dispersed before adding other pesticides.
5. Continue to fill the tank with carrier to the desired volume while agitating. Continue agitation during application to ensure a uniform spray mixture.

**RE-SUSPENDING SC PRODUCTS IN SPRAY SOLUTION**

Like other suspension concentrates (SC's), CORVUS Herbicide will settle if left standing without agitation. If the spray solution is allowed to settle for one hour or more, reagitrate the spray solution for a minimum of 10 minutes before application.

**Equipment Cleanup Procedures**

To avoid injury or exposure to non-target crops, thoroughly clean all mixing and spray equipment, including pumps, nozzles, lines and screens with a good quality tank cleaner, on approved rinse pad or on the field site where an approved crop is to be grown. Mix only as much cleaning solution as needed.

1. Flush tank, hoses, boom and nozzles with clean water.
2. Use a pressure washer with a high quality commercial spray tank cleaner in water to clean the inside of the spray tank. Take care to wash all parts of the tank, including the inside top surface. If a pressure washer is not available, completely fill the sprayer with the cleaning solution to ensure contact of the cleaning solution with all internal surfaces of the tank and plumbing. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
3. Flush hoses, spray lines, and nozzles for at least 1 minute with the cleaning solution.
4. Dispose of rinsate from steps 1-3 in an appropriate manner.
5. Repeat steps 2-4.
6. Remove nozzles, screens and strainers and clean separately in the cleaning solution after completing the above procedures.
7. Rinse the complete spraying system with clean water.
8. Cleanup should be conducted on an approved rinse pad or the field site where an approved crop is to be grown.

**ROTATIONAL CROPS**

Rotational crops vary in their response to low concentrations of CORVUS Herbicide remaining in the soil. The amount of CORVUS Herbicide that may be present in the soil depends on soil moisture, soil temperature, application rate, elapsed time since application and other environmental factors. When CORVUS Herbicide is used in combination with other products, always follow the most restrictive rotational crop requirements. The following rotational crops may be planted after applying CORVUS Herbicide.

(continued)

4.0625"  
COPY AREA

7.75"

4.50"

Crop	Rotational Interval <sup>2</sup>	Minimum Precipitation Requirement <sup>1</sup>
Field corn	0 Months	None
Wheat, Triticale, Cereal and rye	4 Months	None
Barley, Soybean, Sweet corn <sup>3</sup> , Popcorn <sup>3</sup>	9 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Rice <sup>3</sup> , Cotton <sup>3</sup>	10 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Peanuts <sup>3</sup>	11 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Tobacco <sup>3</sup>	12 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Alfalfa, Green and Dry Beans, Oats, Sorghum <sup>4</sup> , Sunflower, Canola, Potato, Sugar beet and all other crops <sup>5</sup>	17 Months <sup>3</sup>	30 inches of cumulative precipitation from application to planting of rotational crop

COPY AREA

7.375"

7.75"

<sup>1</sup> The amount of cumulative precipitation required before planting a rotational crop is in addition to the required rotational interval given in months. Furrow or flood irrigation should not be included in total. No more than 7 inches of overhead irrigation should be included in total.

<sup>2</sup> Crop varieties planted back at intervals of one year or less should not have known acute sensitivity to ALS-inhibiting and/or SU herbicides.

<sup>3</sup> When soil pH is 7.5 or above, crop plant back should be delayed to 17 months and to 24 months for crops listed in the 17 month interval above.

<sup>4</sup> For CORVUS Herbicide used at 2.25 - 3.3 fl oz. per acre or less and the total of Thiencarbazone-methyl from all sources is 0.014 pounds active ingredient per acre or less, sorghum can be planted at the 9 month or longer interval.

<sup>5</sup> All other crops may be seeded only after the completion of a successful bioassay after a CORVUS Herbicide application. Refer to the "Field/Small Scale Bioassay" section.

**In the event of crop failure:** If the corn crop treated with CORVUS Herbicide is lost, only field corn and corn grown for silage may be replanted immediately. Do not make an additional application of CORVUS Herbicide.

**Cover Crops**

Use of cover crops as a means of soil improvement, erosion control, weed and/or insect suppression, etc., following harvest of corn in the Fall is increasing. Planting of cover crops in fields treated with CORVUS Herbicide is allowed as long as these cover crops are not grazed by livestock nor harvested for food. Cover crops are to be tilled under or chemically controlled with burndown herbicides in the spring. Many cover crops can be planted within 90-120 days after application of CORVUS Herbicide. However, all potential cover crops have not been evaluated for tolerance to CORVUS Herbicide and significant injury may occur. Prior to seeding a cover crop, complete a successful field/small scale bioassay to provide an indication of the level of tolerance to the prior CORVUS Herbicide application. Refer to the "Field/Small Scale Bioassay" section. If used in tank mixtures with other herbicides, always follow the most restrictive label.

4.0625"

COPY AREA

4.50"

**Field/Small Scale Bioassay**

A field/small scale bioassay must be completed before rotating to a cover crop other than those specified in the "Rotational Crop Restrictions" section of this label. To conduct an effective **field bioassay**, grow strips of the crop(s) you intend to grow the following season in a field previously treated with CORVUS Herbicide. The test strip should be placed in a controlled area and should include low areas and knolls, and include variations in soil such as type and pH. Crop response to the bioassay will determine if the crop(s) grown in the test strips can be grown safely in the areas previously treated with CORVUS Herbicide.

For an effective **small scale bioassay**, collect uniform samples of all soil types from the CORVUS Herbicide-treated field (see example above for types of soil in the sample) and place the soil into a sturdy container. Plant the desired cover crop into the soil, apply water and place the container in a warm sunny area to allow germination and growth of the crop. Monitor growth of the cover crop over a three to four week period. If the crop emerges and grows normally, the risk to establish and grow the cover crop in the CORVUS Herbicide-treated field should be tolerable.

**WEEDS CONTROLLED**

CORVUS Herbicide applied as directed in this label will control or suppress the weeds listed below. Additional weeds may be controlled with tank mixtures or sequential applications (refer to the Tank Mix Instructions and Sequential Application Instructions sections of this label). Always refer to the tank mix partner labels for specific use rates and additional directions.

BROADLEAF WEEDS		
Amaranth, palmer	Mallow, Venice	Ragweed, common
Buffalobur	Marestail	Ragweed, giant 2,3,4
Burcucumber <sup>2</sup>	Medic, black 2,3	Russian thistle
Buttercup, small flower	Morningglory, annual 2,3,4	Sesbania, hemp
Carpetweed	Mustard, wild	Shepherd's-purse
Chamomile spp	Nightshade, black	Sicklepod 2,3,4
Chickweed, common	Nightshade, eastern black	Sida, prickly
Clover, purple 2,3,4	Nightshade, hairy	Smartweed, Penn.
Clover, white 2,3,4	Pennycress, field	Smartweed, ladysthumb
Cocklebur 2,3,4	Pepperweed, Virginia	Speedwell, corn 2,3
Copperleaf, Hophornbeam	Pigweed, prostrate	Spurge, toothed
Dandelion, (seedling)	Pigweed, redroot	Sunflower, wild 2,3,4
Deadnettle, purple	Pigweed, smooth	Velvetleaf
Galinsoga	Pigweed, tumble	Vetch, bird 2,3,4
Henbit	Plantain, broadleaf	Violet, field 2,3,4
Jimsonweed	Puncturevine, common	Waterhemp, tall
Kochia	Purslane, common	Waterhemp, common
Lambsquarters, common	Radish, wild	

GRASS/SEDGE WEEDS		
Barnyardgrass	Crabgrass, smooth	Foxtail, giant
Bluegrass, annual 2,3	Cupgrass, woolly 1	Foxtail, green
Crabgrass, large	Foxtail, bristly	Foxtail, robust white

(continued)

COPY AREA

7.375"

7.75"

4.0625"

COPY AREA

4.50"

Foxtail, robust purple	Millet, wild proso <sup>2</sup>	Panicum, Texas <sup>2</sup>
Foxtail, yellow	Nutsedge, yellow <sup>2,3</sup>	Sandbur, field <sup>2</sup>
Goosegrass	Oat, tame	Shattercane <sup>1</sup>
Johnsongrass, seedling	Oat, wild	Signalgrass, broadleaf
Millet, browntop	Panicum, fall	Witchgrass <sup>1</sup>

<sup>1</sup> These weeds may require an appropriate sequential postemergence herbicide treatment for control of late season escapes.

<sup>2</sup> These weeds will be partially controlled. Partially controlled weeds will be reduced competition by stunted growth and/or reduced populations as compared to non-treated areas. Commercially acceptable control may require the application of an appropriate preemergence tank mixture or sequential postemergence herbicide treatment.

<sup>3</sup> Control of these weeds can be gained with the addition of an approved label rate of atrazine.

<sup>4</sup> These weeds may require a postemergence application of DiFlexx<sup>®</sup> Herbicide or other appropriate postemergence herbicides.

### SPECIFIC USE DIRECTIONS

#### CORN (Field Corn, Seed Corn and Corn Grown for Silage)

CORVUS Herbicide may be used in either conventional, conservation tillage, or no-till crop management systems and may be applied either preplant, preplant incorporated (less than 2" deep), preemergence or early postemergence.

CORVUS Herbicide treatments are most effective in controlling weeds when adequate rainfall is received within 14 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain occurs, use shallow tillage such as rotary hoe to lightly incorporate CORVUS Herbicide. Make certain corn seeds are below the tilled area. If treated soil is moved during tillage practices in such a way that the herbicide barrier is no longer intact, weeds may emerge from areas where treated soil has been removed. Do not incorporate with a drag harrow after planting.

#### APPLICATION RATES

Application Timing	Fluid oz of CORVUS Herbicide per Acre <sup>1</sup>			
	Soil Texture			
	Coarse Soils 2.0% O.M. <sup>2</sup> or less Sand, Loamy sand, Sandy loam	Coarse Soils greater than 2.0% O.M. <sup>2</sup> Sand, Loamy sand, Sandy loam	Medium Soils Loam, Silt loam, Silt, Sandy clay loam	Fine Soils Silty clay loam, Clay loam, Sandy clay, Silty clay, Clay
Preplant <sup>3</sup> (Surface Applied or Incorporated) Preemergence Early postemergence	3.33	5.6 <sup>4</sup>	5.6 <sup>4</sup>	5.6 <sup>4</sup>

<sup>1</sup> If soils are 2.0% or less in O.M. and have a pH of 7.5 or greater, the rate selected from the table above can be reduced by 0.5 fluid oz.

<sup>2</sup> O.M. = Organic Matter by weight.

<sup>3</sup> CORVUS Herbicide may be applied alone or in recommended tank-mixes up to 21 days prior to planting. CORVUS Herbicide may be applied up to 30 days prior to planting when used in a planned sequential application program such as CORVUS followed by Liberty 280<sup>®</sup> Herbicide, DiFlexx<sup>®</sup> Herbicide, Laudis<sup>®</sup> Herbicide, or other postemergence applied herbicides appropriate for control of the target weeds.

<sup>4</sup> For coarse textured soils with greater than 2.0% O.M. or medium textured soils with 2.0% O.M. or less, and where densities of weeds controlled by CORVUS Herbicide are light to moderate, an appropriate rate down to 4.5 fluid oz per acre may be selected.

4.0625"  
COPY AREA

COPY AREA

7.375"

7.75"

4.50"

**APPLICATION TIMING**

**Preplant Surface-Applied**

CORVUS Herbicide may be applied up to 21 days before planting corn. CORVUS Herbicide may be applied up to 30 days prior to planting when used in a planned sequential application program such as CORVUS Herbicide followed by DiFlexx® Herbicide, Liberty 280® SL Herbicide, or other postemergence applied herbicides appropriate for control of the target weeds. Refer to all parts of the label of the respective sequential partner for specific use directions and restrictions. Split applications of CORVUS Herbicide can be made. It is recommended that 60% of the listed broadcast rate (refer to Application Rate Table) may be applied 15 – 30 days prior to planting and the remaining 40% applied at planting. Total CORVUS Herbicide applied may not exceed the listed rate for a preplant treatment on the predominant soil type in the field. Moving treated soil out of the row or moving untreated soil to the surface during planting may result in reduced weed control.

**Preplant Incorporated**

CORVUS Herbicide may be applied up to 21 days before planting corn. CORVUS Herbicide may be applied up to 30 days prior to planting when used in a planned sequential application program such as CORVUS Herbicide followed by DiFlexx® Herbicide, Liberty 280® SL Herbicide, or other postemergence applied herbicides appropriate for control of the target weeds. Refer to all parts of the label of the respective sequential partner for specific use directions and restrictions. Apply to the soil and uniformly incorporate in the top two inches of soil before planting using a finishing disc, field cultivator or similar implement capable of providing uniform two inch incorporation. Do not incorporate CORVUS Herbicide deeper than 2" or weed control may be reduced.

**Preplant/Preemergence Burndown**

When weeds are present at the time of treatment and prior to corn emergence, a tank mixture of CORVUS Herbicide (+/- DiFlexx® Herbicide) with COC or MSO is recommended for burndown of labeled weeds 6" or less in height. When weeds are greater than 6" in height or weeds not controlled by CORVUS Herbicide are present, the addition of a burndown herbicide (e.g., Liberty 280® SL Herbicide, paraquat, glyphosate, or 2, 4-D) is recommended. If giant ragweed, common cocklebur, henbit, Pennsylvania smartweed or purple deadnettle are present at the time of application, the addition of atrazine will improve control. Observe directions for use, precautions and restrictions, and adjuvants on the label of the burndown tank-mixed herbicide. When mixing with liquid nitrogen fertilizer or certain glyphosate formulations, substitute a non-ionic surfactant for oil concentrates.

**Preemergence**

Apply CORVUS Herbicide during planting (behind the planter after furrow closure) or after planting, but before weeds emerge. Failure to thoroughly close and firm the seed furrow may allow herbicide to directly contact the seed which can cause injury.

**Early Postemergence**

CORVUS Herbicide can be applied to corn in tank mixture with atrazine from spiking through the 2-leaf collar growth stage. Tank-mixtures with other herbicides or adjuvants are not recommended for early postemergence applications of CORVUS Herbicide to emerged corn as crop response symptoms including bleaching, leaf edge necrosis and stunting may result. Do not use COC or MSO with CORVUS Herbicide applied to emerged field corn.

Early postemergence applications of CORVUS Herbicide should be made in water as the carrier. Sprayable fluid fertilizer as an herbicide carrier for early postemergence applications in corn can typically cause corn injury up to and including tissue burn (necrosis). Sprayable fluid fertilizer as a carrier is not recommended for use with CORVUS Herbicide after crop emergence unless typical fertilizer burn symptoms on the crop are acceptable.

*(continued)*

4.0625"

COPY AREA

COPY AREA

7.375"

7.75"

4.50"

Do not apply tankmixtures of CORVUS Herbicide with organophosphate or carbamate insecticides to emerged corn. Foliar applications of an organophosphate or carbamate insecticides should not be made within 7 days of an application of CORVUS Herbicide or crop injury may result.

**TANK MIX INSTRUCTIONS**

CORVUS Herbicide may be used in tank mixtures with other herbicides for improved control of certain broadleaf and grass weeds in corn. Preplant/preemergence/early postemergence tank-mixes with CORVUS Herbicide include but are not limited to those listed. Refer to and follow all parts of the label of each tank-mix partner.

Tank-mix combinations may be used in either conventional, conservation tillage or no-till cropping systems and may be applied at the same timings as CORVUS Herbicide unless otherwise specified on this label or on the tank-mix partner's label.

Multiple tank mixtures are allowed unless otherwise specified by the respective product labels. Check all tank-mix product labels for proper rates and compatibilities for multiple tank-mixes.

**Possible Preplant/Preemergence Tank Mix Partners for Additional Weed Control**

Anthem®	DIFLEXX® Herbicide	LIBERTY® 280 Herbicide (LibertyLink traited corn only)	Surpass®
Anthem® ATZ	Glyphosate [including Roundup® and Touchdown® branded products (Roundup Ready traited corn only)]	Paraquat (including Gramoxone® branded products)	Verdict®
Atrazine (including Aatrex® branded products)	Harness®	Sharpen™ Herbicide	Zidua®
Dicamba	Harness® Xtra	Simizine/Princep	2, 4-D

**Possible Early Postemergence\* Tank Mix Partners For Additional Weed Control**

Atrazine (including Aatrex® branded products)	DIFLEXX® Herbicide		
---	--------------------	--	--

\*See instructions in APPLICATION TIMING section of this label for early postemergence tank mixtures, adjuvants and carrier solutions for directions on the use of tank mixtures with CORVUS Herbicide after crop emergence.

**SEQUENTIAL APPLICATION INSTRUCTIONS**

Sequential herbicide applications either before or following CORVUS Herbicide treatments may be used to control additional weeds. Herbicides to use sequentially with CORVUS Herbicide treatments include but are not limited to those listed. Refer to all parts of the individual product labels of herbicides used in sequence with CORVUS Herbicide.

COPY AREA

7.375"

7.75"

4.0625"

COPY AREA

4.50"

**Possible Sequential Herbicide Applications For Additional Weed Control**

Dicamba-containing herbicides (Status, Banvel, etc.)	Glyphosate [including Roundup® and Touchdown® branded products (Roundup Ready traited corn only)]	LAUDIS® Herbicide	
DIFLEXX® Herbicide	LIBERTY® 280 Herbicide (LibertyLink traited corn only)		

If CORVUS Herbicide has been applied preplant/preemergence and use of a HPPD herbicide (Laudis, Callisto, Armezon, Impact, etc.) is planned as a sequential postemergence application, always tankmix the postemergence HPPD herbicide with an additional effective mode of action herbicide(s).

**SEED/SOIL-APPLIED INSECTICIDE INTERACTIONS**

CORVUS Herbicide can be used in conjunction with Poncho® seed treatments and a variety of registered seed and soil-applied insecticides. Use of CORVUS Herbicide with soil and seed-applied insecticides on all corn hybrids should follow the recommendations in the table below. DO NOT USE CORVUS Herbicide in the same year as Counter® 15G, Counter® 20G, or any other organophosphate or carbamate soil-applied insecticides not specifically recommended.

Seed or Soil-Applied Insecticide	Use Pattern	Use of CORVUS Herbicide in the Same Year
Poncho®, Poncho®/Votivo®, Aztec®, Regent®, Force®, Chlorpyrifos (e.g. Lorsban®), Phorate (e.g. Thimet®), and Fonophos (e.g. Dyfonate®, Chlorethoxyfos (e.g. Fortress®), Bifenthrin (e.g. Capture®)	All	No use precautions
Terbufos (e.g. Counter® 15G, Counter® 20CR,) and other organophosphate or carbamate insecticides.	All	DO NOT USE

**PRECAUTIONS FOR USE**

- **Planting depth:** Corn seed should be planted a minimum of 1-1/2 inches deep and must be completely covered with soil and furrow firmed or reduced crop stand or injury may occur.
- **Effect of variable soils on use rate:** The proper use rate of CORVUS Herbicide is affected by several soil factors, including soil texture, organic matter, and soil pH. Soils which contain variations in one or more of these factors in a given area are termed variable soils and may be more likely to incur localized corn injury symptoms from an application of CORVUS Herbicide, especially in those localized areas containing a more coarse soil texture, a lower organic matter and/or a higher pH (alkaline/calcareous soil) than other areas of the same field. The user is responsible for selecting the appropriate rate of CORVUS Herbicide as specified in the table above that corresponds to all soils in the area of application.
- **Effect of adverse weather:** Following an application of CORVUS Herbicide, extended periods of cool/cold, wet conditions (cool/cold daytime/nighttime temperatures, saturated soil conditions, recurring rainfall events, etc.) during corn seed germination and/or early crop development period may result in temporary crop injury. Injury symptoms may appear as leaf tissue bleaching (whitening) and/or crop stunting. Corn plants usually recover from this injury without affecting yield.

(continued)

4.0625"

COPY AREA

COPY AREA

7.375"

7.75"

4.50"

COPY AREA

7.375"

- **Corn hybrids and certain male pollinators:** Corn hybrids and certain male pollinators within blended corn varieties vary in their response to CORVUS Herbicide. Not all hybrids or male pollinators within blended corn varieties have been tested for sensitivity to CORVUS Herbicide. You should consult with your seed provider, your local Bayer CropScience representative and/or other knowledgeable agricultural professionals for advice on tolerance of hybrids or varieties containing male pollinator lines before applying CORVUS Herbicide. If the tolerance of a hybrid or variety containing male pollinator lines is not known, you should apply CORVUS Herbicide to a small area to first determine if the hybrid is tolerant prior to spraying large acreages of that hybrid.

**RESTRICTIONS FOR USE**

- Application: Do not exceed maximum labeled rate for soil type. Spray overlaps produce areas of over application which increase the potential for crop damage.
- In corn, the following Corvus components must not exceed per acre per 365 day period from all sources: 0.04 pounds Thiencarbazone-methyl, 0.094 pounds Isoxaflutole, or 0.20 pounds Cyprosulfamide
- Do not use CORVUS Herbicide in the same season as certain soil-applied organophosphate or carbamate insecticides (refer to the 'SEED/SOIL-APPLIED INSECTICIDE INTERACTIONS' section of the label).
- Do not use CORVUS Herbicide on popcorn, or sweet corn.
- Do not irrigate CORVUS Herbicide into coarse soils at planting time when soils are saturated.
- Do not harvest field corn forage within 45 days of application of CORVUS Herbicide.
- Do not use COC or MSO with CORVUS Herbicide applied to emerged field corn.
- Do not apply tankmixtures of CORVUS Herbicide with organophosphate or carbamate insecticides to emerged corn.
- Do not apply solo HPPD inhibitor Postemergence herbicides to corn that has been treated with CORVUS Herbicide in the same growing season.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide storage**

Store in a cool, dry secured storage area.

**Pesticide disposal**

Dispose of wastes resulting from the use of this product on site or at an approved waste disposal facility.

**Container handling**

**Non-Seed Treatment Products in Non-Refillable Containers**

**Rigid, Non-refillable containers (equal to or less than 5 gallons)**

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix

*(continued)*

4.0625"  
COPY AREA

7.75"



4.50"

tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

**Rigid Non-refillable Containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)**

Non-refillable container. Do not reuse or refill this container. Refer to Bottom Discharge IBC information as follows.

**Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Autumn™ Super WDG Herbicide, Corvus® Herbicide, DiFlexx® Herbicide, Laudis® Herbicide, and Liberty® 280 SL Herbicide are registered trademarks of Bayer CropScience.

Aatrex® branded products, Callisto®, Gramoxone®, and Touchdown® are registered trademarks of a Syngenta Group Company.

Harness®, Harness Xtra® and Roundup® are registered trademark of Monsanto Company.

Impact® is a registered trademark of AMVAC.

Armezon®, Sharpen®, Status®, Verdict® and Zidua® are registered trademarks of BASF Corporation.

Banvel is a registered trademark of Arysta Life Sciences North America LLC.

Anthem® and Anthem® ATZ are registered trademarks of FMC Corporation.

Surpass® is a registered trademark of Dow AgroSciences.

COPY AREA


7.375"


7.75"

Bayer

4.0625"

COPY AREA






4.50"

COPY AREA  
7.375"

7.75"

18

4.0625"  
COPY AREA



BASE LABEL

5.50"

**RESTRICTED USE PESTICIDE**

May injure (phytotoxic) susceptible non-target plants.  
For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial and certified applicators must ensure that all persons involved in these activities are informed of the precautionary statements.

**CORVUS® Herbicide**

**GROUP 2/27 HERBICIDE**

**For:** weed control in field corn, seed corn and corn grown for silage in the states of: AR, AL, CO, DE, GA, IL, IN, IA, KS, KY, LA, MI, MN, MO, MS, MT, NE, NJ, NM, NC, ND, OH, OK, PA, SC, SD, TN, TX, VA, WI and WY.

In the states of CO, DE, KS, MD, MO, NJ, NM, SD, and WV use is only allowed under 24c registrations. A current 24c label must be in the possession of the user at the time of the pesticide application.

In the state of MN use is only allowed in accordance with the Minnesota Product Bulletin.

In the state of WI use is only allowed in accordance with the Wisconsin Product Bulletin.

<b>ACTIVE INGREDIENTS:</b> Thien carbazono-methyl: (Methyl 4-[[[4,5-dihydro-3-methoxy-4-methyl-5-oxo-1H-1,2,4-triazol-1-yl]carbonyl]amino]sulfonyl]-5-methyl-3-thiophenecarboxylate) .....	<b>7.60%</b>
Isoxaflutole [5-cyclopropyl-4-(2-methylsulfonyl-4-trifluoromethylbenzoyl) isoxazole] .....	<b>19.00%</b>
<b>OTHER INGREDIENTS:</b> .....	<b>73.40%</b>
	<b>TOTAL 100.00%</b>

Contains 0.75 pounds Thien carbazono-methyl per U.S. gallon  
Contains 1.88 pounds Isoxaflutole per U.S. gallon

EPA Reg. No. 264-1066

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours a Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

See additional precautionary statements and directions for use on label.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC  
ANIMALS  
CAUTION**

- Harmful if swallowed or absorbed through the skin.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing.

or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide storage**

Store in a cool, dry secured storage area.

**Pesticide disposal**

Dispose of wastes resulting from the use of this product on site or at an approved waste disposal facility.

**Container handling**

**Non-Seed Treatment Products in Non-Refillable Containers**

**Rigid, Non-refillable containers (equal to or less than 5 gallons)**

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment

Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
CORVUS is a registered trademark of Bayer.  
©2016 Bayer CropScience  
US61380343C 161219C 12/16

NET CONTENTS: 2 1/2 GALLONS

COPY AREA

7.50"



7.75"

0.875"

4.125"

0.50"

COPY AREA